

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

The Egyptian Heart Journal

journal homepage: www.elsevier.com/locate/ehj

Original Article

Masked uncontrolled hypertension: Prevalence and predictors

Ghada Youssef^{a,*}, Sherif Nagy^a, Ahmed El-gengehe^a, Amr Abdel Aal^b, Magdy Abdel Hamid^a^a Cairo University, Cairo, Egypt^b Helwan University, Cairo, Egypt

ARTICLE INFO

Article history:

Received 26 June 2018

Accepted 3 October 2018

Available online 22 October 2018

Keywords:

Ambulatory BP monitoring

Masked hypertension

Uncontrolled hypertension

ABSTRACT

Background: There are limited data on 'masked uncontrolled hypertension' (MUCH) in patients with treated and apparently well-controlled BP is unknown.**Objectives:** To define the prevalence and predictors of MUCH among hypertensive patients with controlled office blood pressure.**Methods:** One hundred ninety-nine hypertensive patients presented to the specialized hypertension clinics at two University Hospitals. All patients had controlled office blood pressure (less than 140/90 mmHg). Patients were assessed regarding history, clinical examination, and laboratory data. All patients underwent ambulatory blood pressure monitoring (ABPM) for 24 h, within a week after the index office visit. MUCH was diagnosed if average 24-h ABPM was elevated (systolic BP \geq 130 mmHg and/or diastolic BP \geq 80 mmHg) despite controlled clinic BP.**Results:** Sixty-six patients (33.2%) had MUCH according to 24-h ABPM criteria (mean age 53.5 ± 9.3 years, 60.6% men). MUCH was mostly caused by the poor control of nocturnal BP; with the percentage of patients in whom MUCH was solely attributable to an elevated nocturnal BP almost double that due to daytime BP elevation (57.3% vs. 27.1%, $P < 0.001$). The most common predictors of MUCH were smoking, DM and positive family history of DM.**Conclusion:** The prevalence of masked suboptimal BP control is high. Office BP monitoring alone is thus inadequate to ascertain optimal BP control because many patients have an elevated nocturnal BP. ABPM is needed to confirm proper BP control, especially in patients with high cardiovascular risk profile. Smoking, DM and positive family history of DM were the most common predictors of MUCH.© 2018 Egyptian Society of Cardiology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Masked hypertension (MH) is a term used to define people who have a normal seated clinic blood pressure (BP) but an elevated out-of-office BP, as determined by ambulatory BP monitoring (ABPM) or home BP monitoring (HBPM). Masked hypertension is

the opposite of the more commonly recognized 'white coat hypertension'. Patients with MH are now known to be at particularly high risk of developing cardiovascular disease (CVD) because they often remain undetected and untreated.¹

Most studies on the prevalence of MH have primarily focused on 'treatment naïve' patients, prior to the diagnosis of hypertension, and many of them based the measurements on HBPM or daytime ABPM, or were of small size.¹ This daytime definition of MH didn't include people whose sole abnormality is an elevation in nocturnal BP, which some studies suggest is the strongest predictor of CVD risk compared with daytime or 24-h mean pressures.² Furthermore, few studies have established the prevalence of the equivalent of MH, i.e. 'masked uncontrolled hypertension; MUCH', in patients with treated hypertension. MUCH is used to describe treated patients in whom BP levels are sub-optimally controlled according to ABPM, but who are considered controlled according to clinic BP targets by current treatment guidelines recommendations (<140/90 mmHg). Despite the recognized potential for clinic BP alone to both over- and under diagnose hypertension, to date,

Abbreviations: A/C ratio, Albumin/Creatinine ratio; ABPM, Ambulatory Blood Pressure Monitoring; ACEI, Angiotensin Converting Enzyme Inhibitor; ARBs, Angiotensin Receptor Blockers; BB, Beta Blockers; BMI, Body Mass Index; BP, Blood Pressure; CAD, Coronary Artery Disease; CCB, Calcium Channel Blocker; CKD, Chronic Kidney Disease; CV, Cardio-Vascular; CVD, Cardio-Vascular Diseases; DBP, Diastolic Blood Pressure; DM, Diabetes Mellitus; ECG, ElectroCardioGram; HBPM, Home Blood Pressure Monitoring; HDL Cholesterol, High-Density Lipoprotein Cholesterol; HTN, Hypertension; LDL Cholesterol, Low-Density Lipoprotein Cholesterol; LVH, Left Ventricular Hypertrophy; MH, Masked Hypertension; MUCH, Masked Uncontrolled Hypertension; NHP, National Hypertension Project; SBP, Systolic Blood Pressure; SD, Standard Deviation; TOD, Target Organ damage.

Peer review under responsibility of Egyptian Society of Cardiology.

* Corresponding author at: Fostatat New City, Misr Qadeema, Cairo, Egypt.

E-mail address: ghadayoussef@kasralainy.edu.eg (G. Youssef).<https://doi.org/10.1016/j.ehj.2018.10.001>

1110-2608/© 2018 Egyptian Society of Cardiology. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

few guidelines, as NICE 2011 guidelines have recommended the routine use of ABPM to monitor the quality of BP control because there are very little data on the quality of BP control in routine clinical practice.³

The aim of this study is to determine the prevalence and predictors of MUCH in hypertensive patients with controlled office BP.

2. Methods

This is a prospective, non-randomized, observational, cross sectional study that enrolled 199 HTN patients who presented to the specialized HTN clinics at two university hospitals. Patients were recruited from February 2016 to June 2017. Inclusion included hypertensive patients on regular antihypertensive treatment who had controlled office blood pressure readings (less than 140/90 mmHg and \leq 140/85 for diabetics, for at least two visits, one month apart).⁴ Excluded from the study were those with secondary hypertension, acute myocardial infarction, significant valvular heart disease, decompensated heart failure (New York Heart Association class III and IV), and pregnant ladies.

Patients gave informed consent about being included in this study. They underwent full clinical evaluation including cardiovascular risk factors assessment e.g. history of diabetes mellitus, smoking and their duration, current medications, family history of CV risk factors and current antihypertensive drugs (class and dosage).

Examination included; assessment of the body mass index (BMI) (Obesity is defined as BMI > 30 Kg/m²), waist circumference, supine heart rate, peripheral pulses as well as searching for signs of target organ damage.

Blood pressure measurement was done using a digital fully automated device (Omron-6 automated device).⁵ Patients were allowed to rest for 3–5 min before measurement. Three BP readings were taken, 1–3 min apart, the first one was omitted and the last two readings were averaged. Patients were allowed to stand unsupported for 2 min and then standing BP readings were recorded.

Laboratory workup included (Hemoglobin level, serum Creatinine, potassium, total cholesterol, low density lipoprotein, high density lipoprotein, and triglycerides, fasting blood sugar and uric acid). Fundus examination was asked for to detect significant hypertensive retinopathy (\geq grade II hypertensive retinopathy). Urine analysis was performed in all studied patients, those who had proteinuria underwent albumin creatinine (A/C) ratio. Patients with abnormal A/C ratio (defined as having albuminuria above 30 mg/dl) are considered to have proteinuria as a marker of target organ damage.⁶

Standard 12 lead ECG was done in all patients. Abnormalities as arrhythmias, premature beats, ischemic heart disease, conduction defects and left ventricular hypertrophy (LVH) were documented. Criteria for LVH diagnosis with ECG were followed.⁷

Target organ damage (TOD) including; LVH, carotid bruit, hypertensive retinopathy \geq grade II, peripheral arterial disease, and clinical CVD (coronary heart disease, congestive heart failure) were diagnosed, using the appropriate investigation, and were documented. Chronic renal disease was diagnosed when serum creatinine was >1.3 mg/dl and/or when proteinuria was present.

All patients underwent 24-hour ABPM. ABPM was conducted on the patient's non-dominant arm using Holter system Model DMS 300-4A⁸ with device set to measure the BP every half an hour in daytime and every hour during the night, according to the patient's sleep and awake times. The patients were asked to continue performing their normal routines but remain still during the measurements. Blood pressure measurement performed for all patients on all days of the working week. Average day, night, and 24-hour blood pressure and pulse rate of patients were collected.

Dipping (i.e. nocturnal blood pressure fall) has been categorized into four groups:(6) (a) normal dipping; where the ratio between mean night systolic and mean day systolic is (0.8–0.9), (b) no dipping; where the ratio is (0.9–1), (c) reverse dipping; the ratio is more than 1 and (d) extreme dipping; the ratio is less than 0.8.

Valid ABPM recordings had to fulfill a series of pre-established criteria, including successful recording of \geq 80% of systolic BP (SBP) and diastolic BP (DBP) during both the daytime and nocturnal periods, and at least one BP measurement per hour.

The primary aim is to detect the prevalence of MUCH which is defined as:(6) normal office BP and, (a) mean awake ABPM readings \geq 135/85 and/or (b) mean night ABPM readings \geq 120/70 and/or (c) mean average 24H ABPM readings \geq 130/80.

2.1. Statistical analysis

Quantitative variables were expressed as mean and standard deviation (SD), while qualitative variables were presented as numbers and percentages.

We divided the study patients into two groups; group 1 with normal office and normal mean 24 h ABPM, (controlled HTN) and group 2 with normal office and elevated mean 24 h ABPM, (masked uncontrolled HTN; MUCH).

We compared the two groups regarding demographics, risk factors, target organ damage and other parameters by means of Chi-square/Fisher exact test for categorical data, and student *t*-test for continuous data. Linear Regression analysis was used to detect predictors of MUCH. All statistical tests were 2 sided, and we judged a P-value of <0.05 to be significant. All analyses were carried out using SPSS 20.

3. Results

Demographic and clinical characteristics of patients are demonstrated in Table 1. Most patients were middle aged. About one third had diabetes mellitus (DM) and one third were current heavy cigarette smokers.

The data obtained from the Ambulatory BP analysis shows that about two-thirds of patients had non-dipping or reversed dipping patterns in nocturnal BP readings.

About one third of patients (n = 66, 33.2%) were diagnosed to have MUCH, according to 24-hours ABPM readings. Taking only day time ABPM, 54 (27.1%) patients had MUCH, while when using only nighttime BP, 114 (57.2%) patients had MUCH. Nighttime ABPM seems to cause greater impact on the high prevalence of MUCH found in 24-hours average ABPM analysis. Diagnosis of MUCH (in 24-hours average BP) is mainly due to combined elevation of both systolic and diastolic BP (48%) rather than elevated only systolic (30.3%) or only diastolic (21.2%) blood pressures.

Characteristically, patients with MUCH had more prevalence of cardiovascular risk factors; higher prevalence of DM, dyslipidemia, heart failure, smoking and higher prevalence of positive family history of HTN and DM. They also had inadequate response to standing as compared to the controlled HTN group.

Both groups showed comparable results regarding the prescribed antihypertensive medication, Fig. 1. The most frequently prescribed anti-hypertensive drug, in both groups, was beta blocker and the least prescribed was diuretic.

Laboratory workup of patients with MUCH is shown in Table 2. No characteristic laboratory difference was found between the 2 groups.

Linear regression analysis showed that the most significant predictors of MUCH were smoking, DM, and a positive family history of DM, Table 3.

Table 1
Clinical profile of patients with MUCH versus patients with controlled BP.

Variable	Patients with MUCH (n = 66), No (%)	Patients with controlled HTN (n = 133), No (%)	P value
Gender, Male	40 (60.6)	54 (40.6)	0.008
Age, years (Mean ± SD)	53.5 ± 9.3	53.8 ± 10.5	0.8
Non-employed	22 (33.3)	63 (47.4)	0.06
Illiterate	13 (19.7)	37 (27.8)	0.21
Associated comorbidities			
Diabetes mellitus	44 (66.7)	64 (48.1)	0.013
Dyslipidemia	29 (43.9)	27 (20.3)	<0.001
Known CKD	11 (16.7)	9 (6.8)	0.029
CAD	8 (12.1)	6 (4.5)	0.048
History of stroke	6 (9.1)	14 (10.5)	0.8
Heart Failure	4 (6.1)	5 (3.8)	0.5
Smokers	4 (6.1)	1 (0.8)	0.042
Family history of HTN	30 (45.5)	26 (19.5)	<0.001
Family history of DM	56 (84.4)	77 (57.9)	<0.001
Family history of DM	51 (77.3)	49 (36.8)	<0.001
Abnormal ECG	27 (40.9)	35 (26.3)	0.04
Abnormal fundus examination (n = 176)	12 (19.7)	13 (11.3)	0.13
TOD	37 (56.1)	38 (28.6)	<0.001
	Mean ± SD	Mean ± SD	
BMI	32.2 ± 6.4	31.0 ± 7.0	0.2
Obesity, Kg. no (%)	37 (56.1)	58 (43.6)	0.2
Waist circumference; cm			
Male	100.0 ± 8.9	98.7 ± 12.6	0.6
Female	103.3 ± 13.8	99.8 ± 14.9	0.3
Office BP measurements			
Supine SBP	130.0 ± 6.9	128.5 ± 7.7	0.2
Supine DBP	80.9 ± 4.7	79.6 ± 5.4	0.1
Standing SBP	134.2 ± 8.4	129.8 ± 9.0	0.001
Standing DBP	85.1 ± 8.0	83.3 ± 7.1	0.1
Supine heart rate			
ABPM measurements			
Daytime average SBP	138.7 ± 9.5	121.0 ± 8.5	<0.001
Daytime average DBP	83.7 ± 8.0	71.4 ± 7.1	<0.001
Nighttime average SBP	130.6 ± 11.1	113.2 ± 10.0	<0.001
Nighttime average DBP	77.1 ± 9.1	65.0 ± 8.4	<0.001
24-hours average SBP	135.8 ± 8.4	118.6 ± 7.5	<0.001
24-hours average DBP	81.6 ± 6.9	69.6 ± 6.4	<0.001
Dipping category			
Extreme dipping	2 (3.0)	1 (0.8)	0.3
Normal dipping	16 (24.2)	42 (31.6)	0.5
No dipping	34 (51.5)	67 (50.4)	0.9
Reversed dipping	14 (21.2)	23 (17.3)	0.5

BMI; Body mass index, BP; Blood pressure, CAD; Coronary artery disease, CKD; Chronic kidney disease, DBP; Diastolic blood pressure, DM; Diabetes mellitus, ECG; Electrocardiogram, HTN; Hypertension, MUCH; Masked uncontrolled hypertension, SBP; Systolic blood pressure, TOD; Target organ damage. Significant p values are marked in bold.




Fig. 1. Percentage of prescribed antihypertensive drugs in both study groups. MUCH; Masked uncontrolled hypertension; HTN; Hypertension, BB; Beta blockers, CCB; Calcium channel blockers, ACEI; Angiotensin enzyme inhibitor, ARBs; Angiotensin receptor blockers.

Table 2
Laboratory characteristics of MUCH patients versus controlled HTN patients.

Variable	Patients with MUCH (n = 66), Mean ± SD	Patients with controlled HTN (n = 133), Mean ± SD	P value
Serum creatinine	1.1 ± 0.5	1.1 ± 1.0	0.8
Fasting blood sugar	107.2 ± 30.8	100.6 ± 26.7	0.1
Total Cholesterol	172.0 ± 25.6	174.9 ± 34.2	0.6
LDL-Cholesterol	115.9 ± 21.9	114.2 ± 26.8	0.7
HDL-Cholesterol	46.9 ± 6.3	46.3 ± 8.8	0.7
Triglycerides	161.1 ± 32.0	154.8 ± 37.0	0.3
Uric acid	5.6 ± 1.3	6.4 ± 5.8	0.5
Serum potassium	4.2 ± 0.4	4.2 ± 0.3	0.9
Hemoglobin	13.1 ± 1.1	13.3 ± 1.2	0.2

HDL; High density lipoprotein, HTN; Hypertension, LDL; Low density lipoprotein, MUCH; Masked uncontrolled hypertension.

Table 3
Regression analysis of the predictors of MUCH.

Variable	Standardized Beta-coefficient	P value
Gender	0.039	0.638
DM	0.138	0.037
Dyslipidemia	0.056	0.404
CKD, diagnosed by elevated serum creatinine level	0.010	0.882
History of heart failure	0.090	0.165
Family history of HTN	0.043	0.574
Family history of DM	0.252	0.001
Smoking	0.165	0.046
TOD	0.112	0.126
Standing systolic BP	0.184	0.004

BP; Blood Pressure, CKD; Chronic Kidney Disease, DM; Diabetes Mellitus, HTN; Hypertension, TOD; Target Organ Damage.
Significant p values are marked in bold.

4. Discussion

Hypertension (HTN) is a chief public-health problem challenging both economically developed and developing countries as it is highly coupled with cardiovascular and kidney diseases.⁹ Data from the Egyptian National Hypertension Project (NHP; 1993–1995) showed that prevalence of HTN was 26.3% among Egyptian adults. Awareness rate among Egyptians was 37.5% with 23.9% of patients receiving anti-hypertensive medications and control rate of only 8%.¹⁰

For several years, BP measurement in the clinic was the golden standard for detection and diagnosis of clinical HTN and monitoring the beneficial effect of anti-hypertensive medications. With the introduction of ABPM and HBPM to clinical practice, new clinical terms for describing HTN were introduced. One of the underscored terms in clinical practice is MH which was first described by Pickering in 2002.¹¹ Despite the term was originally used for untreated hypertensive persons, later periodicals used the term to refer to patients with treated hypertension.¹²

The exact mechanism responsible for MH is not completely recognized. In order to understand the mechanism behind MH it was postulated that it may be the result of reduced office BP and/or increased ABPM. Lower office BP measurement may be due to white coat effect which is the difference between office and out of office BP and this effect is negative in patients with MH.¹³ Another reason that may attribute to lower office BP is the relation between diagnostic labeling as hypertensive and office BP, as it was found that the absence of diagnostic labeling as hypertensive was found to be associated with lower office BP.¹⁴ Smoking, alcohol consumption, physical activity and psychosocial factors (anxiety, interpersonal conflict and job stress) may all contribute to increase in ABPM.¹⁵

The prevalence of MH was reported to range from 8% to 49% with tendency to be higher in treated hypertensive.¹⁶ This discrepancy of MH prevalence is attributed to several factors as the characteristic of the population studied (general population vs. clinic-based population, treated HTN vs. HTN naïve, ethnic background) and ABPM criteria used to define MH (day time BP vs. 24 h BP) and the use of different BP thresholds for defining MH. Our study which is a clinic-based study on patients with treated HTN showed MUCH prevalence of 33.2%. In concordant to our results, the Spanish registry¹⁷ reported MUCH prevalence of 31.1% in Spanish patients with treated HTN. Similar prevalence of MUCH in treated HTN was reported by Pierdomenico et al.¹⁸ However, our results were higher than that reported by SHEAF study 9.4%¹⁹ and J-home study 19%²⁰ both of which used HBPM rather than ABPM for detection of MUCH.

Nighttime BP is known to be a strong predictor factor for total, cardiovascular, stroke and cardiac mortality.²¹ Elevated night time BP showed a great impact on the prevalence of MUCH in our study as 57.3% of the patients proved to have MUCH using only nighttime BP vs. 27.1% when only daytime BP was used. About 80% of patients reported marked discomfort with the device especially at night and they were awakened from sleep by cuff inflations. The resulting disturbed sleep rhythm may have altered sympathetic activity leading to nocturnal surge of BP.

Results from the Spanish registry¹⁷ showed lower prevalence of nocturnal HTN (24.3% vs 57.3% in our study).

One of the findings of this study is that patients with MUCH showed higher standing SBP compared to those with controlled BP. An inverse relationship between BP response to standing and the difference between clinic BP and daytime BP was documented before. Compared to patients with normal reaction to standing, patients with increased reaction showed higher levels of systolic and diastolic ABPM.²² Such data indicate that increased reactivity to standing is predictive of higher ABPM and explain the reason why patient with MUCH had higher standing SBP in our study.

This study showed that patients with MUCH had high prevalence of cardiovascular risk factors and TOD which is concordant to what reported by Pickering et al,²³ the Spanish registry¹⁷ and Japan home study²⁰ which signifies the importance of early detection and treatment of patients with MH since such patients are at increased risk of cardiovascular mortality and stroke.²⁴ Whether the high-risk profile is a consequent or merely an association to MUCH is not yet known.

Using multivariate analysis, smoking, DM and family history of DM were found to be the strongest predictor of MH in our study. Bromfield et al²⁵ also reported diabetes to be associated with a higher prevalence of masked daytime and isolated nocturnal uncontrolled hypertension among African Americans taking anti-hypertensive medication in the Jackson Heart Study. The Spanish registry¹⁷ showed that after multivariable adjustment, the odds ratio for masked 24-hour uncontrolled hypertension associated with diabetes taking antihypertensive medications was 1.25 (95% CI = 1.14–1.37).

5. Limitations

This study has several limitations that we have to address. It was performed on a small sample size of hypertensive patients and a study on a larger number of patients from different geographic regions across the country is needed to verify our findings. Also, our study doesn't reflect the general population as patients were recruited from specialized HTN clinics and multi-center population-based study may be required. The diagnosis of MUCH was based on a single AMBP recording and would have been better to repeat the ABPM to test the reproducibility of MUCH diagnosis.

6. Conclusion

More than one third of patients showed MUCH despite apparently well controlled office BP readings. Elevated nocturnal BP was acting as a major determinant of the presence of MUCH, a finding which cannot be detected by regular clinic measurements. Patients with MUCH showed a higher constellation of traditional cardiovascular risk factors and TOD, which imposes tight BP control in order to reduce future cardiovascular events. Our recommendation is to suspect MUCH in apparently controlled HTN patients with high risk profile and to order ABPM for these patients for better evaluation and management of HTN.

Conflicts of interest

The authors declare that none of them had any conflict of interests.

References

- Mancia G, Bombelli M, Seravalle G, Grassi G. Diagnosis and management of patients with white-coat and masked hypertension. *Nat Rev Cardiol*. 2011;8:686–693. Epub 2011/08/10.
- Verdecchia P, Angeli F, Mazzotta G, et al.. Day-night dip and early-morning surge in blood pressure in hypertension: prognostic implications. *Hypertension*. 2012;60:34–42. Epub 2012/05/16.
- Ritchie LD, Campbell NC, Murchie P. New NICE guidelines for hypertension. *BMJ*. 2011;343. Epub 2011/09/09 d5644.
- Mancia G, Fagard R, Narkiewicz K, et al.. 2013 ESH/ESC practice guidelines for the management of arterial hypertension. *Blood Press*. 2014;23:3–16. Epub 2013/12/24.
- OMRON. Available from: <<http://www.omron-healthcare.com/en/products/blood-pressure-monitoring>>.
- O'Brien E, Parati G, Stergiou G, et al.. European Society of Hypertension position paper on ambulatory blood pressure monitoring. *J Hypertens*. 2013;31:1731–1768. Epub 2013/09/14.
- Surawicz B, Nihilans T. *Chou's electrocardiography in clinical practice: adult and pediatric*. 6th ed. Saunders Elsevier; 2008.
- Chine Medical Device. DMS Holter Recorder Available from: <<http://www.chinamedevice.com/chinasuppliers/5070/DMS-Holter-Recorder-DMS300-3A-DMS300-4A-454330.html>>.
- Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet*. 2005;365(9455):217–223. Epub 2005/01/18.
- Ibrahim MM, Rizk H, Appel LJ, et al.. Hypertension prevalence, awareness, treatment, and control in Egypt. Results from the Egyptian National Hypertension Project (NHP). NHP Investigative Team. *Hypertension*. 1995;26:886–890. Epub 1995/12/01.
- Pickering TG, Davidson K, Gerin W, Schwartz JE. Masked hypertension. *Hypertension*. 2002;40:795–796. Epub 2002/12/07.
- Aksoy I, Deinum J, Lenders JW, Thien T. Does masked hypertension exist in healthy volunteers and apparently well-controlled hypertensive patients? *Neth J Med*. 2006;64:72–77. Epub 2006/03/21.
- Pickering TG, Eguchi K, Kario K. Masked hypertension: a review. *Hypertens Res*. 2007;30:479–488. Epub 2007/08/01.
- Spruill TM, Pickering TG, Schwartz JE, et al.. The impact of perceived hypertension status on anxiety and the white coat effect. *Ann Behav Med*. 2007;34:1–9. Epub 2007/08/11.
- Ogedegbe G. Causal mechanisms of masked hypertension: socio-psychological aspects. *Blood Press Monit*. 2010;15:90–92. Epub 2010/03/12.
- Papadopoulos DP, Makris TK. Masked hypertension definition, impact, outcomes: a critical review. *J Clin Hypertens (Greenwich)*. 2007;9:956–963. Epub 2007/11/30.
- Banegas JR, Ruilope LM, de la Sierra A, et al.. High prevalence of masked uncontrolled hypertension in people with treated hypertension. *Eur Heart J*. 2014;35:3304–3312. Epub 2014/02/06.
- Pierdomenico SD, Lapenna D, Bucci A, et al.. Cardiovascular outcome in treated hypertensive patients with responder, masked, false resistant, and true resistant hypertension. *Am J Hypertens*. 2005;18:1422–1428. Epub 2005/11/11.
- Bobrie G, Chatellier G, Genes N, et al.. Cardiovascular prognosis of “masked hypertension” detected by blood pressure self-measurement in elderly treated hypertensive patients. *JAMA*. 2004;291:1342–1349. Epub 2004/03/18.
- Obara T, Ohkubo T, Kikuya M, et al.. Prevalence of masked uncontrolled and treated white-coat hypertension defined according to the average of morning and evening home blood pressure value: from the Japan Home versus Office Measurement Evaluation Study. *Blood Press Monit*. 2005;10:311–316. Epub 2006/02/24.
- Dolan E, Stanton A, Thijs L, et al.. Superiority of ambulatory over clinic blood pressure measurement in predicting mortality: the Dublin outcome study. *Hypertension*. 2005;46:156–161. Epub 2005/06/09.
- Narkiewicz K, Piccolo D, Borella P, Businaro R, Zonzin P, Palatini P. Response to orthostatic stress predicts office-daytime blood pressure difference, but not nocturnal blood pressure fall in mild essential hypertensives: results of the Harvest trial. *Clin Exp Pharmacol Physiol*. 1995;22:743–747.
- Liu JE, Roman MJ, Pini R, Schwartz JE, Pickering TG, Devereux RB. Cardiac and arterial target organ damage in adults with elevated ambulatory and normal office blood pressure. *Ann Intern Med*. 1999;131:564–572. Epub 1999/10/16.
- Ohkubo T, Kikuya M, Metoki H, et al.. Prognosis of “masked” hypertension and “white-coat” hypertension detected by 24-h ambulatory blood pressure monitoring 10-year follow-up from the Ohasama study. *J Am Coll Cardiol*. 2005;46:508–515. Epub 2005/08/02.
- Bromfield SG, Shimbo D, Bertoni AG, Sims M, Carson AP, Muntner P. Ambulatory blood pressure monitoring phenotypes among individuals with and without diabetes taking antihypertensive medication: the Jackson Heart Study. *J Hum Hypertens*. 2016;30:731–736. Epub 2016/05/14.