

## Metabotyping and its role in nutrition research

Elaine Hillesheim<sup>1,2</sup> and Lorraine Brennan<sup>1,2\*</sup>

<sup>1</sup>UCD Institute of Food and Health, UCD School of Agriculture and Food Science, UCD, Belfield, Dublin 4, Republic of Ireland

<sup>2</sup>UCD Conway Institute of Biomolecular and Biomedical Research, UCD, Belfield, Dublin 4, Republic of Ireland

### Abstract

Personalised nutrition is at its simplest form the delivery of dietary advice at an individual level. Incorporating response to different diets has resulted in the concept of precision nutrition. Harnessing the metabolic phenotype to identify subgroups of individuals that respond differentially to dietary interventions is becoming a reality. More specifically, the classification of individuals in subgroups according to their metabolic profile is defined as metabotyping and this approach has been employed to successfully identify differential response to dietary interventions. Furthermore, the approach has been expanded to develop a framework for the delivery of targeted nutrition. The present review examines the application of the metabotype approach in nutrition research with a focus on developing personalised nutrition. Application of metabotyping in longitudinal studies demonstrates that metabotypes can be associated with cardiometabolic risk factors and diet-related diseases while application in interventions can identify metabotypes with differential responses. In general, there is strong evidence that metabolic phenotyping is a promising strategy to identify groups at risk and to potentially improve health promotion at a population level. Future work should verify if targeted nutrition can change behaviours and have an impact on health outcomes.

**Key words:** Cluster analysis: Metabotypes: Personalised nutrition: Targeted nutrition

### Introduction


Poor diet quality is a major contributor to chronic diseases such as type 2 diabetes, CVD and various cancers<sup>(1,2)</sup>. Despite the well-known association between dietary patterns and diseases, interventions to change dietary habits have had a limited impact on wellbeing and public health outcomes<sup>(3,4)</sup>. In recent years, the diverse inter-individual responses to interventions have become apparent and support the need for the development of strategies that are based upon the delivery of advice to the individual<sup>(5-9)</sup>. Concomitant with this, different strategies have emerged for delivering advice taking personal characteristics into account. Furthermore, studies have demonstrated that personalisation of dietary advice is more effective in promoting improvements in the dietary habits of individuals compared with the general healthy eating advice<sup>(10-12)</sup>.

Metabolomics is the study of small molecules in biological samples and is a powerful tool in the characterisation of individuals<sup>(13,14)</sup>. The set of metabolites in the human body, termed the metabolome, is the product of metabolic reactions influenced by endogenous, lifestyle and environmental factors<sup>(15,16)</sup>. Applications of metabolomics in nutrition research have expanded in recent years and it has the potential to contribute to the delivery of personalised nutrition<sup>(17)</sup>. Metabotypes are defined as groups of similar individuals based on combinations of specific metabolites. Thus, individuals within a metabotype

have similar metabolic profiles and those in different metabotypes have different profiles<sup>(17,18)</sup> (Fig. 1). Metabotypes are often defined using cluster analysis, such as *k*-means analysis and hierarchical cluster analysis<sup>(18)</sup>. Applications of metabotypes has identified differential response to interventions and have the potential of identifying optimal treatment strategies for individuals. For example, using serum metabolites Palau-Rodriguez *et al.*<sup>(19)</sup> identified two subgroups with different degrees of improvement in insulin resistance, total cholesterol (TC), HDL-cholesterol (HDL-C) and uric acid following bariatric surgery. Importantly, the metabolic changes in each cluster were independent of the baseline anthropometric/clinical parameters of the patients and the magnitude of weight loss. Another example identified metabotypes with different lipid responses to fenofibrate<sup>(20)</sup>. Similarly, in the field of nutrition science there are several examples of applications of metabotypes in healthy and subjects with chronic diseases for determining metabolically homogeneous subgroups with differential responses to dietary interventions<sup>(18)</sup>. However, the applications are not limited to intervention studies, with the metabotyping approach being developed for the delivery of targeted nutrition<sup>(21,22)</sup>. Given the rapid growth of this area, the objective is to review the research conducted on metabotypes related to nutrition research and to identify gaps where further work is needed.

**Abbreviations:** HDL-C, HDL-cholesterol; HOMA-IR, homeostatic model assessment for insulin resistance; HSFAM, high-SFA meal; IGF, insulin-like growth factor; MetS, metabolic syndrome; MMM, mixed Mediterranean-type meal; MMTT, mixed meal tolerance test; OGTT, oral glucose tolerance test; OLTT, oral lipid tolerance test; RCT, randomised controlled trial; TC, total cholesterol.

\* **Corresponding author:** Lorraine Brennan, email [lorraine.brennan@ucd.ie](mailto:lorraine.brennan@ucd.ie)



**Fig. 1.** Overview of the concept of metabotyping for the delivery of personalised nutrition. Intrinsic and extrinsic factors influence the metabolic phenotype of individuals. Groups of individuals with similar metabolic phenotypes are termed metabotypes.

**Metabolic phenotyping of longitudinal data to examine associations with cardiometabolic risk factors and diet-related diseases**

Longitudinal studies are important tools in the epidemiological setting to investigate the aetiology of a disorder and indicate risk factors or population groups that may be targeted as part of prevention strategies. In fact, within the metabolic phenotype approach, longitudinal studies offer the possibility to study subgroups of individuals (metabotypes) over a period of time and the potential to identify those at higher risk of disease development. A summary of studies examining longitudinal associations of metabotypes with cardiometabolic risk factors and diet-related diseases is presented in Table 1.

In order to identify risk profiles for the emergence of the metabolic syndrome (MetS), Ventura *et al.*<sup>(23)</sup> assessed a non-clinical sample of healthy non-Hispanic white girls (*n* 154) in a retrospective analysis with follow-up performed every 2 years from age 5 to 13 years old. Six risk factors for the MetS (waist circumference, systolic blood pressure, diastolic blood pressure, HDL-C, TAG and blood glucose) were used in cluster analysis to determine metabotypes at age 13 years. At age 5 years, the higher MetS risk group had the highest BMI relative to the other groups. Across childhood, both the higher MetS risk and the hypertension risk groups had significantly greater increases in weight and fat mass, while the higher MetS risk group had the highest daily sweetened beverage intake. Findings from this



**Table 1.** Summary of studies examining longitudinal associations of metabolotypes with cardiometabolic risk factors and diet-related diseases

Author	Objective	Study design	Study sample	Follow-up period	Variables and method for clustering	Main findings
Ventura <i>et al.</i> <sup>(23)</sup>	Describe risk profiles for the metabolic syndrome during adolescence	Retrospective longitudinal study	154 Non-clinical 13-year-old white girls in the USA	Every 2 years for 8 years	Six risk factors for the metabolic syndrome (waist circumference, SBP, DBP, and fasting HDL-C, TAG and glucose) clustered by mixture model	Four metabolotypes. At age 13 years, the higher metabolic syndrome risk group and the hypertension risk group had more family history of type 2 diabetes and obesity. Across childhood, the higher metabolic syndrome risk group and the hypertension risk group had greater increases in BMI and fat mass, as well as the former had the higher intake of sweetened beverages; a dyslipidaemia risk group had the lowest physical activity
Kirchberg <i>et al.</i> <sup>(24)</sup>	Identify predictive metabolotypes for childhood obesity	Prospective longitudinal study	154 Healthy, singleton, term and breastfed infants aged 6 months in the CHOP trial in Europe	6 years	21 Fasting plasma amino acids, sum of hexoses and 146 polar lipids (free carnitine, 40 acylcarnitines, 11 lyso PC, 91 PC, and 14 sphingomyelins) clustered by the Bayesian agglomerative method	Twenty metabolotypes. Only the four biggest clusters ( $n \geq 14$ ) were analysed and at 6 months of age cluster 3 had the lowest weight, height, free IGF-1 and IGF-BP3, and the highest IGF-BP2. The BMI z-score at 6 years of age tended to differ (unadjusted $P = 0.07$ ) among clusters, with cluster 3 presenting the highest median and large proportion of overweight/obese children
Riedl <i>et al.</i> <sup>(25)</sup>	Define metabolotypes of diet-related diseases	Prospective longitudinal study	1729 Adults aged 32–77 years in the population-based KORA F4 study in Germany	7 years	BMI and 33 fasting biochemical parameters clustered by $k$ -means cluster analysis	Three metabolotypes. At the baseline, cluster 3 showed the most unfavourable marker profile with the highest prevalence of cardiometabolic diseases. After the follow-up, disease incidence was higher in cluster 3 compared with clusters 2 and 1, respectively, for hypertension (41.2, 25.3, 18.2%), type 2 diabetes (28.3, 5.1, 2.0%), hyperuricaemia/gout (10.8, 2.3, 0.7%), dyslipidaemia (19.2, 18.3, 5.6%), all metabolic diseases (54.5, 36.8, 19.7%) and all CVD (6.3, 5.5, 2.3%) together

SBP, systolic blood pressure; DBP, diastolic blood pressure; HDL-C, HDL-cholesterol; CHOP, Childhood Obesity Project; PC, phosphatidylcholines; IGF-1, insulin-like growth factor 1; IGF-BP3, insulin-like growth factor-binding protein 3; IGF-BP2, insulin-like growth factor-binding protein 2.

study support the role of metabolotypes for identifying individuals at higher risk who could be targeted by clinicians as part of preventive healthcare.

Application of metabolotypes to baseline data in longitudinal studies can be very useful in defining at-risk groups which could be targeted for the prevention of undesirable health outcomes. The European Childhood Obesity Project (CHOP), using a Bayesian agglomerative clustering method on twenty-one plasma amino acids and 146 polar lipids, classified healthy infants ( $n$  154) of 6 months of age into twenty metabolotypes in order to predict later obesity risk<sup>(24)</sup>. Only the four biggest clusters ( $n \geq 14$ ) were analysed and at the baseline cluster 3 had the lowest weight, height, free insulin-like growth factor (IGF)-1 and IGF-binding protein 3, and the highest IGF-binding protein 2. The BMI  $z$ -score at 6 years of age tended to differ (unadjusted  $P = 0.07$ ) among clusters, with cluster 3 presenting the highest median and largest proportion of overweight/obese children. These results support the concept that even very young individuals can be clustered according to their inter-individual differences so that the clusters provide insight into later development and health and opportunities for developing more targeted and personalised intervention strategies.

Another notable example employing metabolotypes in a prospective cohort is the KORA F4 study in which 1729 adults aged 32 to 77 years were clustered based on BMI and thirty-three biochemical markers<sup>(25)</sup>. For each of the three metabolotypes identified, the current disease prevalence and the incidence in the follow-up cohort 7 years later was determined. The 'high-risk' cluster showed the most unfavourable biomarker profile with the highest BMI and prevalence of cardiometabolic diseases at the baseline as well as the highest incidence of hypertension, type 2 diabetes, hyperuricaemia/gout, dyslipidaemia, all metabolic diseases and all CVD together. This study provides strong evidence that metabolotyping is a robust approach for identifying groups of individuals that could be targeted for prevention strategies.

Overall, the derivation of metabolotypes in longitudinal studies to predict cardiometabolic risk factors and diet-related diseases is nascent. However, replication of the metabolotypes in other populations is a necessary next step. Notwithstanding this, the presented studies make a strong case for the metabolotype approach and highlight its potential in identifying groups that could benefit from targeted dietary advice.

### Metabolic phenotyping to investigate differential responses to dietary challenges and interventions

Differential responses to dietary interventions are becoming increasingly recognised. Concomitantly, metabolic phenotyping has emerged as a useful tool to examine responses to interventions. In the context of nutrition, health can be defined as the ability of an organism to adapt to challenges<sup>(26)</sup>. Challenge tests investigate the disturbance and restoration of homeostasis of an individual using a dietary challenge as a physiological stressor<sup>(27)</sup>. In combination with metabolomics, dietary challenges have been used to identify groups of subjects with distinct metabolic phenotypes/metabolotypes and unique responses.

Table 2 illustrates studies which focus specifically on differential responses of metabolotypes to dietary challenges and intervention studies.

Krishnan *et al.*<sup>(28)</sup> investigated the differential responses of metabolotypes to dietary challenges. The authors used low- and high-glycaemic index meals in a cross-over randomised trial with healthy overweight women ( $n$  24; aged 20–50 years) to identify response patterns that could provide insight into early subclinical glycaemic disruption. By using blood glucose, insulin and leptin responses to the challenges, individuals were clustered into three metabolotypes. While the most populated metabolotype presented little deviation from the expected response to the dietary challenges, the two minor metabolotypes were suggestive, one of sub-clinical insulin resistance and the other of hyperleptinaemia. In the Metabolic Challenge (MECHE) Study, healthy subjects ( $n$  214; aged 18–60 years) were randomised to one of three groups to receive oral glucose tolerance tests (OGTT) and/or oral lipid tolerance tests (OLTT) and four metabolotypes were identified based on their blood glucose response curves to the OGTT ( $n$  116)<sup>(29)</sup>. The cluster with the most adverse metabolic profile at baseline presented a reduced  $\beta$ -cell function and differential responses to insulin and C-peptide during the OGTT and OLTT, as well as to glucose and TAG during the OLTT, which characterises this metabolotype as at risk. The postprandial metabolic responses to different kinds of bread – refined rye bread, wholemeal rye bread and a control refined wheat bread – were investigated in a cross-over randomised controlled trial (RCT) with healthy postmenopausal women ( $n$  19; aged 61 (SD 4.8) years)<sup>(30)</sup>. The clustering of the fasting metabolic profile identified two distinct metabolotypes. Women with higher fasting concentrations of leucine and isoleucine and lower fasting concentrations of sphingomyelins and phosphatidylcholines had higher insulin responses despite similar glucose concentrations after all kinds of bread, suggesting higher insulin resistance. In a recent study with data from the NutriTech project, the response to the intervention was only evident following the classification of the individuals into metabolotypes<sup>(26)</sup>. Healthy subjects ( $n$  72; aged 59 to 64 years) were enrolled to a mixed meal tolerance test (MMTT) before and after 12 weeks targeting moderate weight loss (basal BMI 29.7 (SD 2.7) kg/m<sup>2</sup>). The intervention group ( $n$  40) consumed a diet that reduced energy intake by 20%, whereas subjects in the control group ( $n$  32) consumed an average European diet matched to their energy expenditure to maintain body weight. Two metabolotypes were reported based on the plasma concentration of metabolites (markers of lipolysis, fatty acid  $\beta$ -oxidation and ketogenesis) during the mixed meal challenge test. Before the intervention, individuals from metabolotype B ( $n$  36) showed slower glucose clearance, increased visceral fat volume, higher hepatic lipid concentrations, and a less healthy dietary pattern according to the urinary metabolomic profile when compared with individuals from metabolotype A. Following the weight loss (about 5.6 kg), only the individuals from metabolotype B showed positive changes in the glycaemic response to the MMTT. Since the metabolite differences found between metabolotypes A and B are all closely associated with insulin signalling, metabolotype B was considered to be prediabetic with a modestly impaired insulin action. Collectively, all these studies clearly demonstrate that the use of a metabolotype

**Table 2.** Summary of studies investigating differential responses of metabolotypes to meal challenges and dietary interventions

Author	Objective	Study design	Study sample	Dietary challenge(s)	Intervention	Variables and method for clustering	Main findings
Fiamoncini <i>et al.</i> <sup>(28)</sup>	Investigate the metabolic response of metabolotypes to an MMTT before and after weight loss	Metabolic challenge before and after a 12-week RCT	70 Healthy subjects (based on fasting glucose, insulin and blood pressure) aged 59–64 years in the NutriTech Study in Europe	Mixed-meal tolerance test (400 ml of high-energy drink with 33% carbohydrates, 59% lipids and 8% protein)	Control group: European diet for weight stability. Intervention group: supervised diet for weight loss	Response concentrations of plasma markers of lipolysis, fatty acid $\beta$ -oxidation and ketogenesis clustered by HCA	Two metabolotypes. At baseline, metabolotype B had slower glucose clearance, increased intra-abdominal adipose tissue mass, higher hepatic lipid concentrations, and a less healthy dietary pattern than metabolotype A. Following the weight loss (about 5.6 kg), only metabolotype B showed positive changes in the glycaemic response to the MMTT, with improvements in metabolites of amino acids, acylcarnitines and biochemical parameters Three metabolotypes. The two minor groups were one suggestive of sub-clinical insulin resistance and the other of hyperleptinaemia Four metabolotypes. Cluster 1 was at risk with the highest BMI, TAG, hsCRP, C-peptide, insulin and HOMA-IR and the lowest $VO_2$ max. Cluster 1 had a reduced $\beta$ -cell function and differential responses to insulin and C-peptide during OGTT and to insulin, glucose and TAG during OLTT Two metabolotypes. Subgroup B, with the lower fasting concentrations of sphingomyelins and diacyl-PC and the higher concentrations of BCAA, had the higher insulin responses to all kinds of bread, despite a similar glucose response to metabolotype A, suggesting higher insulin resistance
Krishnan <i>et al.</i> <sup>(29)</sup>	Identify metabolotypes of response to meals with different GI	Metabolic challenge in a cross-over randomised trial	24 Healthy premenopausal women aged 20–50 years in the USA	High-GI and low-GI meals preceded by a 3 d run-in diet matching the GI of the tested meal	Not tested	Response concentrations of blood glucose, insulin and leptin clustered by PCA	
Morris <i>et al.</i> <sup>(29)</sup>	Identify metabolotypes of response to an OGTT	Metabolic challenge in a randomised trial	116 Healthy subjects aged 18–60 years in the Metabolic Challenge (MECHE) Study in Ireland	75 g OGTT or an OLTT (54 g of lipids and 12 g of carbohydrates)	Not tested	Response curves of blood glucose to OGTT clustered by mixed model	
Moazzami <i>et al.</i> <sup>(30)</sup>	Investigate the metabolic response of metabolotypes to different types of bread	Metabolic challenge in a cross-over RCT	19 Healthy postmenopausal women (61 (SD 4.8) years) in Finland	Refined wheat, wholemeal rye and refined rye breads, providing 50 g of carbohydrate	Not tested	189 Fasting metabolites (21 amino acids, 17 biogenic amines, 47 acylcarnitines, 38 PC, 39 acyl-alkyl PC, 14 lyso PC, 15 sphingomyelins, and 1 hexose) clustered by O-PLS, HCA and PCA	



**Table 2** Continued

Author	Objective	Study design	Study sample	Dietary challenge(s)	Intervention	Variables and method for clustering	Main findings
Lacroix <i>et al.</i> <sup>(31)</sup>	Evaluate the endothelial and metabolic response of metabolotypes to complete meals	Metabolic challenge in a cross-over RCT	28 Healthy men aged 18–50 years in Canada	HSFAM and MMM	Not tested	Age, BMI, HOMA-IR, and fasting glucose, insulin, TC, LDL-C, HDL-C, and TAG clustered by HCA	Two metabolotypes. Group 1 had a higher BMI, HOMA-IR, and fasting insulin, TC, non-HDL-C, TAG and TAG:HDL-C, and a lower intake of fruits and vegetables. Following the MMM, the healthiest group (group 2) had a lower increase in TAG, with no difference in postprandial endothelial function. The HSFAM induced postprandial endothelial dysfunction only in group 1
Wang <i>et al.</i> <sup>(32)</sup>	Identify metabolotypes of response to dietary carotenoids	Cross-over 3-week trial	23 Healthy subjects aged 36–69 years in the USA	Not tested	Watermelon juice (20.1 mg/d lycopene + 2.5 mg/d carotene) and a second watermelon juice (40.2 mg/d lycopene + 5.0 mg/d carotene) or tomato juice (18.4 mg/d lycopene + 0.6 mg/d carotene)	Temporal response concentrations of plasma carotenoids (β-carotene, lycopene, phytoene and phytofluene) clustered by <i>k</i> -means cluster analysis	Five metabolotypes per carotenoid per intervention type. Strong or weak responders to each carotenoid were identified. Responses were associated with genetic variants of carotenoid-metabolising enzyme
Vázquez-Fresno <i>et al.</i> <sup>(33)</sup>	Investigate urinary changes in metabolotypes following red wine polyphenol intake	Cross-over 4-week RCT	57 High-risk subjects aged ≥55 years in Spain	Not tested	Red wine polyphenol intake (733 equivalents of gallic acid/d) in the form of dealcoholised wine	67 Fasting blood and urinary markers and 2 anthropometric parameters (BMI and waist:hip ratio) clustered by <i>k</i> -means cluster analysis	Four metabolotypes. Following the intervention, 4-hydroxyphenylacetate concentrations significantly increased in the healthier cluster compared with the higher-risk cluster, while glucose was higher in the higher-risk cluster compared with the healthier cluster; tartarate was higher for both clusters



**Table 2** Continued

Author	Objective	Study design	Study sample	Dietary challenge(s)	Intervention	Variables and method for clustering	Main findings
O'Sullivan <i>et al.</i> <sup>(34)</sup>	Identify metabolotypes of response to vitamin D supplementation in terms of the metabolic syndrome	Double-blind 4-week RCT	135 Healthy subjects aged 18–63 years in Ireland	Not tested	Group 1, 15 µg vitamin D <sub>3</sub> + 10 <sup>9</sup> CFU <i>Lactobacillus salivarius</i> ; Group 2, vitamin D + placebo probiotic; Group 3, placebo vitamin D + probiotic; and Group 4, placebo vitamin D + placebo probiotic	13 Fasting blood markers of the metabolic syndrome (leptin, resistin, adiponectin, IL-6, hsCRP, TNF-α, insulin, C-peptide, TC, TAG, NEFA, glucose, HOMA-IR) and 25(OH)D concentrations clustered by <i>k</i> -means cluster analysis	Five metabolotypes. Cluster 5, with lower serum 25(OH)D and higher concentrations of adipokines at baseline, showed significant improvements in insulin, HOMA-IR and hsCRP, as well as an inverse correlation between changes in serum 25(OH)D and glucose concentrations

MMTT, mixed meal tolerance test; RCT, randomised controlled trial; HCA, hierarchical cluster analysis; GI, glycaemic index; OGTT, oral glucose tolerance test; OLTT, oral lipid tolerance test; hsCRP, high-sensitivity C-reactive protein; HOMA-IR, homeostatic model assessment for insulin resistance; PCA, principal component analysis; OGTT, oral glucose tolerance test; OLTT, oral lipid tolerance test; hsCRP, high-sensitivity C-reactive protein; HOMA-IR, homeostatic model assessment for insulin resistance; PC, phosphatidylcholines; BCAA, branched-chain amino acids; HSFAM, high-SFA meal; MMM, mixed Mediterranean-type meal; O-PLS, orthogonal partial least squares; TC, total cholesterol; LDL-C, LDL-cholesterol; HDL-C, HDL-cholesterol; CFU, colony-forming units; 25(OH)D, 25-hydroxyvitamin D.

approach in conjunction with meal challenges has the ability to characterise individuals into meaningful subgroups which could receive targeted nutrition advice to lower the individual disease risk<sup>(30)</sup>.

In contrast to other studies that used the responses to challenges to form clusters, Lacroix *et al.*<sup>(31)</sup> used only fasting metabolic data in a cross-over RCT designed to evaluate the metabolic and vascular effects of a high-SFA meal (HSFAM) and a mixed Mediterranean-type meal (MMM). Age, BMI, glycaemic and lipid parameters were used to cluster healthy men (*n* 28; 18–50 years) into two metabolotypes at baseline. Compared with the healthiest group, the less healthy group showed significantly higher BMI, insulin and homeostatic model assessment for insulin resistance (HOMA-IR), in addition to a less favourable lipid profile and a lower intake of fruit and vegetables (dietary pattern score = 5.12 (sd 1.7) *v.* 3.9 (sd 1.4)). Following the meal challenges, the less healthy group experienced a greater significant increase in TAG with MMM and endothelial dysfunction with HSFAM, in comparison with the healthier group. The MMM did not significantly alter postprandial endothelial function in both groups. The authors concluded that the less healthy group would benefit even more from consuming meals representative of a Mediterranean-type diet given its non-deleterious endothelial properties, indicating the potential of cluster techniques to individualise dietary advice.

Application of the metabolotype approach has also encompassed dietary interventions that did not involve meal challenges. Wang *et al.*<sup>(32)</sup> in a controlled cross-over study with healthy subjects (*n* 23; aged 36–69 years) identified groups of individuals with differing plasma carotenoid responses to carotenoid-rich beverages. Following 3 weeks of daily intake of watermelon juice (20 mg lycopene, 2.5 mg β-carotene, *n* 23; 40 mg lycopene, 5 mg β-carotene, *n* 12) or tomato juice (18 mg lycopene, 0.6 mg β-carotene, *n* 10), cluster analysis applied to weekly carotenoid responses identified groups of individuals with differential responses. This, in turn, was used to classify individuals as strong responders or weak responders to the carotenoid intake. These findings demonstrate that subgroups of individuals can have differential responses to interventions which could be harnessed in the future to give more precise dietary advice. With respect to employing a metabolotype approach for dietary interventions in clinical populations or disease risk factors, two studies are noteworthy. In a sample of high-risk cardiovascular subjects (*n* 57; aged ≥55 years) a 4-week cross-over RCT identified differential responsiveness to red wine polyphenols<sup>(33)</sup>. At baseline, fasting blood and urinary metabolites and anthropometric parameters were used to cluster individuals in four metabolotypes, including a higher-risk cluster and a healthier cluster. Following 28 d of dealcoholised red wine intake (polyphenol content = 733 equivalents of gallic acid/d), concentrations of urinary 4-hydroxyphenylacetate significantly increased in the healthier cluster compared with the higher-risk cluster, indicating a differential response in this cluster. In a double-blind 4-week RCT with healthy subjects (*n* 135; aged 18–63 years), the effect of vitamin D supplementation (15 mg vitamin D<sub>3</sub> per d) to improve markers of the MetS was only visible after the classification of the sample into metabolotypes<sup>(34)</sup>. The vitamin D supplementation significantly increased the

**Table 3.** Summary of studies developing targeted dietary advice solutions for metabolotypes through the decision tree approach

Author	Study sample	Variables and method for clustering	Clusters' biomarker characterisation	Design of decision trees	Validation of decision trees	Main findings
O'Donovan <i>et al.</i> <sup>(22)</sup>	875 Subjects aged 18–90 years in the Irish National Nutrition Survey in Ireland	Fasting TAG, TC, HDL-C and glucose clustered by <i>k</i> -means cluster analysis	Cluster 1 ( <i>n</i> 274) had high TC, cluster 2 ( <i>n</i> 423) had adequate concentrations of all biomarkers, and cluster 3 ( <i>n</i> 178) had high TAG, TC and glucose	One decision tree by cluster. Dietary advice was based on the biochemical cluster's characteristics and branches for BMI, waist circumference and blood pressure	Comparison with individual-based approach manually compiled and delivered by a dietitian ( <i>n</i> 99)	Three decision trees with 12 possible messages each, which are the combination of 20 possible types of advice. An average agreement of 89% (range 20–100%) was found between the targeted advice and the individual-based approach with 69% of the participants presenting an agreement of 100%
O'Donovan <i>et al.</i> <sup>(35)</sup>	1354 Subjects ≥18 years in the Food4Me Study in seven European countries	27 Fasting metabolic markers (TC, fatty acids and carotenoids) clustered by <i>k</i> -means cluster analysis	Cluster 1 ( <i>n</i> 326) had the highest TC and <i>trans</i> -fatty acids and the lowest omega-3 index; cluster 2 ( <i>n</i> 433) had the highest omega-3 index and total carotenoids and the lowest total saturated fat; and cluster 3 ( <i>n</i> 595) had the lowest TC and highest stearic acid	Two decision trees by cluster. The first was based on biomarkers (TC, total saturated fat, omega-3 index and carotenoids) with branches for TC, BMI and waist circumference. The second was based on the individual intakes of five nutrients (salt, Fe, Ca, folate and fibre)	Comparison with personalised dietary advice based on phenotypic features and delivered by nutritionists ( <i>n</i> 180)	A wide set of messages raised from the combination of two decision trees and ranged from 2 to 6 per participant. An average agreement of 82% was found between the targeted advice and the individual-based approach, with an average agreement of 83, 74 and 88% for clusters 1, 2 and 3, respectively

TC, total cholesterol; HDL-C, HDL-cholesterol.

serum 25-hydroxyvitamin D in comparison with the placebo group, but there was no effect of supplementation on the measured markers of the MetS. Based on thirteen fasting blood biomarkers, one cluster characterised by low concentrations of vitamin D and higher concentrations of adipokines showed a significant decrease in insulin, HOMA-IR scores and C-reactive protein and an inverse relationship between the change in serum vitamin D and glucose. Collectively, these examples clearly present how comprehensive phenotyping may identify subgroups of individuals that can benefit from specific dietary interventions.

The metabolotype approach represents a tool through which we can start to understand individual responses to interventions. The ultimate goal will be to harness this information to deliver personalised nutrition.

### Harnessing the metabolotype approach to deliver targeted nutrition

To the best of our knowledge, there are only two published examples of a framework for the delivery of personalised nutrition using a metabolotype approach (Table 3).

In 2015, O'Donovan *et al.*<sup>(22)</sup> proposed a framework based on metabolotyping using four commonly measured fasting markers of metabolic health (TAG, TC, HDL-C and glucose). Application of

the approach in 875 adults resulted in three metabolotypes. Individuals in cluster 1 (*n* 274) had high TC concentrations, individuals in cluster 2 (*n* 423) had adequate concentrations of all four biomarkers, and individuals in cluster 3 (*n* 178) had the most unfavourable metabolic profile with high concentrations of TAG, TC and glucose and the lowest concentration of HDL-C. Targeted dietary advice was developed for each metabolotype incorporating characteristics of the metabolotype and personal traits. In order to test the reliability of the approach to deliver personalised dietary advice, the targeted approach was compared with an individual-based approach manually compiled and delivered by a dietitian for a random sample of participants (*n* 99). An excellent agreement of 89% (range 20–100%) was found between the methods, considering the dietary advice given with the targeted approach in relation to those given with the individual-based approach. The most important strength of this study is the fact that for clustering individuals only four biomarkers of metabolic health routinely measured were used. Furthermore, the approach generated a limited number of decision trees with simple and clear messages which allow the automation of the delivery of personalised dietary advice to individuals who are not high priority dietetic patients or where the access to a dietitian is limited. All these features make the proposed framework easily transferable to a clinical or primary care setting.

Development of this approach for a more diverse population was achieved in a proof-of-concept format with data from seven



European countries<sup>(35)</sup>. Twenty-seven fasting metabolic markers measured in finger-prick blood samples, including cholesterol, individual fatty acids and carotenoids, were clustered into three metabotypes. Individuals in cluster 1 ( $n$  326) had the highest TC and circulating *trans*-fatty acids and the lowest omega-3 index, so this cluster was therefore considered the metabolically unhealthy cluster. Cluster 2 ( $n$  433) was labelled the healthy group as individuals in this metabotype had the highest average omega-3 index and total carotenoid concentrations and the lowest total SFA. Individuals in cluster 3 ( $n$  595) had the lowest average TC and highest levels of stearic fatty acid. Decision trees with targeted dietary advice were developed on the metabolic markers (TC, total SFA, omega-3 index and carotenoids), demographics and five key nutrients (salt, Fe, Ca, folate and fibre). The targeted approach was compared with the messages delivered by nutritionists as part of the Food4Me study ( $n$  180) to participants receiving personalised dietary advice. An average match of 82% at the level of delivery of the same dietary message was found and the agreement was also good by cluster, with an average match of 83% for cluster 1, 74% for cluster 2 and 88% for cluster 3. These results, obtained in a European population from seven countries with diverse cultures and dietary intakes, confirm the metabotype approach as a robust approach to the delivery of targeted dietary advice and its applicability in different populations.

### Conclusions and future directions

While metabotyping emerged initially to distinguish individuals with and without diet-related diseases, it has rapidly developed to identify those at metabolic risk and interrogate responses to dietary interventions. With a heightened interest in inter-individual variation in response to interventions, the approach presents an unbiased method of identifying differential responses. The ultimate goals will be to harness the approach for the delivery of personalised nutrition. However, further work is needed in understanding the biological mechanisms underlying the differential responses. We need detailed studies examining the underlying biology responsible for the different metabotypes and deciphering the role of genetics and the microbiome will be important future steps. Building this evidence base will be important for the further development of the metabotype concepts.

The framework comprising the metabotypes and decision trees represents a model for the delivery of personalised nutrition. However, there is a paucity of data demonstrating the impact of this approach on metabolic health parameters. Future studies are warranted to demonstrate that the approach is effective in changing behaviours and health outcomes.

### Acknowledgements

E. H. has a grant from the Brazilian Federal Agency for Support and Evaluation of Graduate Education (CAPES) as part of the Full Ph.D. Abroad Program (process no. 88881.174061/2018-01). L. B. is supported by a European Research Council (ERC) grant (no. 647783). CAPES and ERC had no role in the design, analysis or writing of this article.

E. H. and L. B. contributed to the conception and design of the review, E. H. drafted the manuscript, and E. H. and L. B. edited the manuscript.

There are no conflicts of interest.

### References

- Ezzati M & Riboli E (2013) Behavioral and dietary risk factors for noncommunicable diseases. *N Engl J Med* **369**, 954–964.
- Schulze MB, Martínez-González MA, Fung TT, *et al.* (2018) Food based dietary patterns and chronic disease prevention. *BMJ* **361**, k2396.
- Browne S, Minozzi S, Bellisario C, *et al.* (2019) Effectiveness of interventions aimed at improving dietary behaviours among people at higher risk of or with chronic non-communicable diseases: an overview of systematic reviews. *Eur J Clin Nutr* **73**, 9–23.
- Ordovas JM, Ferguson LR, Tai ES, *et al.* (2018) Personalised nutrition and health. *BMJ* **361**, k2173.
- Curran AM, Horner K, O'Sullivan V, *et al.* (2019) Variable glycaemic responses to intact and hydrolyzed milk proteins in overweight and obese adults reveal the need for precision nutrition. *J Nutr* **149**, 88–97.
- Minihane AM, Khan S, Leigh-Firbank EC, *et al.* (2000) ApoE polymorphism and fish oil supplementation in subjects with an atherogenic lipoprotein phenotype. *Arterioscler Thromb Vasc Biol* **20**, 1990–1997.
- Mathias MG, Coelho-Landell CA, Scott-Boyer MP, *et al.* (2018) Clinical and vitamin response to a short-term multi-micronutrient intervention in Brazilian children and teens: from population data to interindividual responses. *Mol Nutr Food Res* **62**, e1700613.
- Manach C, Milenkovic D, Van de Wiele T, *et al.* (2017) Addressing the inter-individual variation in response to consumption of plant food bioactives: towards a better understanding of their role in healthy aging and cardiometabolic risk reduction. *Mol Nutr Food Res* **61**, 1600557.
- Healey GR, Murphy R, Brough L, *et al.* (2017) Interindividual variability in gut microbiota and host response to dietary interventions. *Nutr Rev* **75**, 1059–1080.
- Celis-Morales C, Livingstone KM, Marsaux CF, *et al.* (2017) Effect of personalized nutrition on health-related behaviour change: evidence from the Food4Me European randomized controlled trial. *Int J Epidemiol* **46**, 578–588.
- Curtis PJ, Adamson AJ & Mathers JC (2012) Effects on nutrient intake of a family-based intervention to promote increased consumption of low-fat starchy foods through education, cooking skills and personalised goal setting: the Family Food and Health Project. *Br J Nutr* **107**, 1833–1844.
- Wright JL, Sherriff JL, Dhaliwal SS, *et al.* (2011) Tailored, iterative, printed dietary feedback is as effective as group education in improving dietary behaviours: results from a randomised control trial in middle-aged adults with cardiovascular risk factors. *Int J Behav Nutr Phys Act* **8**, 43.
- Brennan L (2016) Metabolomics in nutrition research – a powerful window into nutritional metabolism. *Essays Biochem* **60**, 451–458.
- de Toro-Martin J, Arsenault BJ, Després JP, *et al.* (2017) Precision nutrition: a review of personalized nutritional approaches for the prevention and management of metabolic syndrome. *Nutrients* **9**, E913.
- Beger RD, Dunn W, Schmidt MA, *et al.* (2016) Metabolomics enables precision medicine: “A White Paper, Community Perspective”. *Metabolomics* **12**, 149.



16. Nicholson JK, Connelly J, Lindon JC, *et al.* (2002) Metabonomics: a platform for studying drug toxicity and gene function. *Nat Rev Drug Discov* **1**, 153–161.
17. Brennan L (2017) Use of metabotyping for optimal nutrition. *Curr Opin Biotechnol* **44**, 35–38.
18. Riedl A, Gieger C, Hauner H, *et al.* (2017) Metabotyping and its application in targeted nutrition: an overview. *Br J Nutr* **117**, 1631–1644.
19. Palau-Rodriguez M, Tulipani S, Marco-Ramell A, *et al.* (2018) Metabotypes of response to bariatric surgery independent of the magnitude of weight loss. *PLOS ONE* **13**, e0198214.
20. van Bochove K, van Schalkwijk DB, Parnell LD, *et al.* (2012) Clustering by plasma lipoprotein profile reveals two distinct subgroups with positive lipid response to fenofibrate therapy. *PLOS ONE* **7**, e38072.
21. O'Donovan CB, Walsh MC, Gibney MJ, *et al.* (2016) Can metabotyping help deliver the promise of personalised nutrition? *Proc Nutr Soc* **75**, 106–114.
22. O'Donovan CB, Walsh MC, Nugent AP, *et al.* (2015) Use of metabotyping for the delivery of personalised nutrition. *Mol Nutr Food Res* **59**, 377–385.
23. Ventura AK, Loken E & Birch LL (2006) Risk profiles for metabolic syndrome in a nonclinical sample of adolescent girls. *Pediatrics* **118**, 2434–2442.
24. Kirchberg FF, Grote V, Gruszfeld D, *et al.* (2018) Are all breastfed infants equal? Clustering metabolomics data to identify predictive risk clusters for childhood obesity. *J Pediatr Gastroenterol Nutr* **68**, 408–415.
25. Riedl A, Wawro N, Gieger C, *et al.* (2018) Identification of comprehensive metabotypes associated with cardiometabolic diseases in the population-based KORA Study. *Mol Nutr Food Res* **62**, e1800117.
26. Fiamoncini J, Rundle M, Gibbons H, *et al.* (2018) Plasma metabolome analysis identifies distinct human metabotypes in the postprandial state with different susceptibility to weight loss-mediated metabolic improvements. *FASEB J* **32**, 5447–5458.
27. Vis DJ, Westerhuis JA, Jacobs DM, *et al.* (2014) Analyzing metabolomics-based challenge tests. *Metabolomics* **11**, 50–63.
28. Krishnan S, Newman JW, Hembrooke TA, *et al.* (2012) Variation in metabolic responses to meal challenges differing in glycemic index in healthy women: is it meaningful? *Nutr Metab (Lond)* **9**, 26.
29. Morris C, O'Grada C, Ryan M, *et al.* (2013) Identification of differential responses to an oral glucose tolerance test in healthy adults. *PLOS ONE* **8**, e72890.
30. Moazzami AA, Shrestha A, Morrison DA, *et al.* (2014) Metabolomics reveals differences in postprandial responses to breads and fasting metabolic characteristics associated with postprandial insulin demand in postmenopausal women. *J Nutr* **144**, 807–814.
31. Lacroix S, Des Rosiers C, Gayda M, *et al.* (2016) A single Mediterranean meal does not impair postprandial flow-mediated dilatation in healthy men with subclinical metabolic dysregulations. *Appl Physiol Nutr Metab* **41**, 888–894.
32. Wang TT, Edwards AJ & Clevidence BA (2013) Strong and weak plasma response to dietary carotenoids identified by cluster analysis and linked to  $\beta$ -carotene 15,15'-monooxygenase 1 single nucleotide polymorphisms. *J Nutr Biochem* **24**, 1538–1546.
33. Vázquez-Fresno R, Llorach R, Perera A, *et al.* (2016) Clinical phenotype clustering in cardiovascular risk patients for the identification of responsive metabotypes after red wine polyphenol intake. *J Nutr Biochem* **28**, 114–120.
34. O'Sullivan A, Gibney MJ, Connor AO, *et al.* (2011) Biochemical and metabolomic phenotyping in the identification of a vitamin D responsive metabotype for markers of the metabolic syndrome. *Mol Nutr Food Res* **55**, 679–690.
35. O'Donovan CB, Walsh MC, Woolhead C, *et al.* (2017) Metabotyping for the development of tailored dietary advice solutions in a European population: the Food4Me study. *Br J Nutr* **118**, 561–569.