



Original article

Healthy dietary pattern and their corresponding gut microbiota profile are linked to a lower risk of type 2 diabetes, independent of the presence of obesity



D.A. Díaz-Rizzolo ^a, B. Kostov ^b, M. López-Siles ^c, A. Serra ^a, C. Colungo ^b,
L. González-de-Paz ^b, M. Martínez-Medina ^c, A. Sisó-Almirall ^{b, d}, R. Gomis ^{a, d, e, f, g, *}

^a Diabetes and Obesity Research Laboratory, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS) – Hospital Clinic of Barcelona, Barcelona, Spain

^b Primary Care Research Group, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Primary Care Centre Les Corts, CAPSBE, Barcelona, Spain

^c Laboratory of Molecular Microbiology, Biology Department, Universitat de Girona, Girona, Spain

^d University of Barcelona, Barcelona, Spain

^e Centro de Investigación Biomédica en Red de Diabetes y Enfermedades Metabólicas Asociadas (CIBERDEM), Spain

^f Department of Endocrinology and Nutrition, Hospital Clinic of Barcelona, Barcelona, Spain

^g Faculty of Health Sciences, Universitat Oberta de Catalunya, Barcelona, Spain

ARTICLE INFO

Article history:

Received 17 August 2018

Accepted 21 February 2019

Keywords:

Diet
Microbiota
Diabetes
Nutrition
Prevention
Obesity

SUMMARY

Background: Prediabetes and old age are both high risk factors for developing Type 2 Diabetes (T2D), while obesity is one of the most important factors triggering the disease. Nutritional interventions are the most effective tool for preventing T2D, as they improve different biochemical and anthropometric outcomes and growth-promoting/inhibiting gut microbiota populations. However, to date there are no specific dietary recommendations to stop the development of T2D in elderly groups, for whom hypocaloric diets and other commonly used weight-loss programs could be considered dangerous. The objective of our study, thus, was to understand the impact of dietary patterns on T2D risk as related to gut microbiota profile in obese and non-obese elderly prediabetic subjects.

Methods: A cross-sectional study was performed in 182 subjects ≥ 65 years old with prediabetes, divided into obese (OB) or non-obese (NOB) subgroups, and their risk of developing T2D was measured according to FINDRISK score and biochemical parameters. Also, clusters into different dietary patterns in each group by PCA analysis was related with gut microbiota, which was analyzed from stool samples by qPCR. The creation of clusters was used to re-evaluate T2D risk.

Results: OB was at higher risk of developing T2D and showed worse metabolic outcomes. Unhealthier and healthier dietary pattern clusters were observed for both OB (OB-6 and OB-5 respectively) and NOB (NOB-2 and NOB-3 respectively) groups. Results obtained from the gut microbiota showed that only Prevotella was higher in NOB, but when comparisons were made between clusters, a clear relation with dietary pattern was observed; showing in healthier dietary clusters a decrease in Prevotella, an increase of Faecalibacterium prausnitzii and an increase in lactic acid bacteria. T2D risk was greater in the obese group between unhealthier dietary clusters. No difference between healthier dietary clusters was observed.

Conclusion: A healthy dietary pattern and the growth-promoting beneficial and growth-inhibiting disadvantageous gut microbiota populations linked to it provide protection against the development of T2D in an obese population with advanced age and preDM.

© 2019 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. Diabetes and Obesity Research Laboratory, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS) – Hospital Clinic of Barcelona, Barcelona, Spain.

E-mail address: ramon.gomis@idibaps.org (R. Gomis).

1. Introduction

Prediabetes (preDM) is defined as a state that precedes type 2 diabetes (T2D), which, together with its complications, is a major cause of early death in most countries [1].

Moreover, preDM, characterized by fasting blood glucose levels of 100–125 mg/dL [2], has been recognized as clinically relevant due to its association with metabolic syndrome and cardiovascular disease (CVD) [3]. Alongside T2D and obesity, preDM has become prevalent worldwide [1]. The natural history of preDM predicts that up to 70% of individuals with this condition will develop T2D, with the annual conversion ratio in prediabetic population around 10% [4,5]. However, the number of new-onset T2D in individuals ≥ 65 years old particularly stands out, increasing 4.5-fold as compared to 3-fold in total population [6].

The prevalence of T2D increases with age but also with the presence of obesity [7], as the risk of developing T2D at 65 years old increases >25% with a BMI ≥ 30 kg/m² [8].

Clinical guidelines recommend weight loss for the prevention of T2D through lifestyle modifications but fail to include specifications for old age [2]. The effects of weight loss in the elderly have been under debate, as some epidemiological studies have shown that it leads to increased mortality [9,10]. This is probably due to the unintentional weight loss of concretely fat-free or lean mass loss that also occurs [11], which is associated with various health problems. Among these is increased insulin resistance [12] and, with it, an increased risk of developing T2D.

Generally, nutritional interventions used for weight loss are based on hypocaloric diets that can lead to malnutrition in the elderly [13]. A more interesting intervention in this age group would be one based on changing dietary patterns instead of focusing on caloric restrictions, but first it is essential to know which dietary patterns are present, information that is currently missing for this population.

Meanwhile, in addition to the known effects that a specific diet can produce on different clinical parameters such as glycaemia, impact on gut microbiota has also been established. Diets provide a range of growth-promoting and growth-inhibiting factors for specific bacterial phylotypes that ferment different substrates [14]. Unhealthy dietary habits have been found to produce dysbiosis, which, in turn, can affect metabolism and increase susceptibility to different diseases. Gut microbiota has been suggested to play an important role in the development of T2D, in conjunction with or independently of obesity [15].

We hypothesize that elderly obese-prediabetic subjects are at higher risk of developing T2D than non-obese ones but that a healthy diet will be linked to a better gut microbiota profile, therefore, obese subjects who show beneficial dietary and gut microbiota pattern will have a lower risk of developing T2D.

2. Methods

This study was a cross-sectional study developed in three primary care centers in Barcelona (CAP Les Corts, Borrell and Casanova) from 2014 to 2016.

2.1. Participants

Subjects included, from whom informed consent was obtained for experimentation and study was performed in accordance with The Code of Ethics of the World Medical Association, were males and females ≥ 65 years old with glucose levels between 100 and 125 mg/dL and without a previous diagnosis of diabetes. Subjects were excluded from the study if they were taking antidiabetic

drugs or had previously received any nutritional education to improve their blood glucose levels or were taking antibiotics.

Based on the clinical records of the primary care centers, a list of 929 patients who met the inclusion criteria was created. 403 were contacted at random, of which 195 were cited for a screening appointment. A final total of 182 subjects were selected for inclusion in the study.

All participants provided written informed consent, and the study protocol was approved by the Ethics Committee of Hospital Clinic de Barcelona.

2.2. Measures

- **Anthropometrics:** Subjects were weighed and measured without clothing and shoes. BMI was calculated as weight (kg)/height (m)², and waist circumference was measured at the midpoint between the last rib and the iliac crest and hip circumference at the widest point of the gluteus.
- **Body composition** was measured using Lunar iDXA body composition (GE Healthcare) according to the standard operating protocol.
- **Blood pressure** was taken after individuals sat quietly for 5 min in a clinical examination room. The mean of three different measurements, obtained every 3 min using an OMRON M6 AC sphygmomanometer, was recorded.
- **Fasting blood samples** were collected by a nurse from each primary care center and were sent, refrigerated at 4 °C, to the central laboratory of Hospital Clinic i Provincial de Barcelona, where they were analyzed by the Biomedical Diagnosis Centre (CDB).
- **T2D risk:** The Finnish Diabetes Risk Score (FINDRISK) questionnaire was completed according of the creation groups of “low”, “medium”, “high” and “very high” risk of developing T2D.
- **Lifestyle habits:** Physical activity was estimated using VREM, a short questionnaire adapted from the Minnesota Leisure Time Physical Activity Questionnaire for individuals of advanced age. Nutritional patterns were measured using a 3-day dietary record revised corrected by a nutritionist and analyzed by the DIAL nutritional calculation program.
- **Gut microbiota:** Genomic DNA was extracted from fecal samples using the PSP[®] Spin Stool DNA Plus kit (Stratec, Berlin, Germany). Fecal samples were collected by the volunteers, immediately frozen and bring to the laboratory in less than 4 hours and were kept at –80°C until analysis. Gut microbiota composition was determined by quantifying the abundance of target bacterial indicators by quantitative PCR (qPCR), namely *Firmicutes*, *Bacteroidetes*, *Faecalibacterium prausnitzii* (*F. prausnitzii*), *Escherichia coli* (*E. coli*), *Prevotella*, and lactic acid bacteria. These indicators were chosen based on previous association with T2D or preDM. Total bacteria quantification was also performed in order to calculate relative abundances of the indicators and to normalize data. 16S rRNA targeted primers, probes, standards for quantification (16S rRNA gene inserted in a pCR[®]4-TOPO[®] cloning plasmid and linearized with NotI), and PCR conditions are detailed in [Supplementary Table S2](#). Standard curves, inhibition controls and non-template controls were included in each run, and samples were run in duplicate. A 7500 Real Time PCR system was used for all quantifications. Data is given as the Log₁₀ of the ratio between 16S rRNA gene copies of the target microorganism and total bacterial 16S rRNA genes detected in the same sample.
- **Statistical analysis:** Descriptive data are presented as the mean and standard deviation (SD) or median and interquartile ranges (IQRs) for continuous outcomes, or number and percentage (%) for categorical outcomes. Non-parametric Mann-

Whitney U test was used for the comparison of continuous outcomes when normality and equality of variance could not be assumed. Student's t test was used for the rest of continuous outcomes and Chi-square test for categorical outcomes. Nutrient pattern analysis was performed using Principal Component Analysis (PCA) based on the intake of 169 nutrients divided into monounsaturated, polyunsaturated, saturated fatty acids, cholesterol and carbohydrates (starches and sugars). In order to capture variability of nutrient intake independently, two separate PCA were performed for obese and non-obese subjects. False discovery rate (FDR) correction was used for multiple comparisons. All significance tests were 2-tailed, and values of $P < 0.05$ were considered significant. All analyses were conducted using the R V.3.0.3 for Windows statistical software package.

3. Results

Our recruited subjects were 49.7% female and 50.3% male. Their mean age was 71.5 ± 5.4 years, while the mean duration of PreDM was 4.7 ± 3.4 years.

T2D risk was calculated, with 86.18% of subjects showing a “high” or “very high” risk of developing the disease, according to FINDRISK test. Since one of the most important risk factors was body weight, participants were divided into two groups according to whether they were obese (OB) or not (NOB), with 45.6% of them presenting a BMI ≥ 30 kg/m².

Clinical characteristics of both groups were similar, with the exception of antithrombotic drug intake, which was more frequent in the OB group, along with systolic (SBP) and diastolic (DBP) blood pressure, which were also higher in the OB group. (Table 1).

Table 1
Characteristics from each group.

	NOB (n=99)	OB (n=83)	P-value
Sex, % female	43.4	56.6	0.104
Age, year	71.15 (4.59)	71.67 (6.04)	0.33
Personal history			
Evolution IFG, years	4.77 (3.36)	4.78 (3.46)	0.976
Family history DM, % cases	36.4	43.4	0.417
Hypertension, % cases	56.6	71.1	0.062
CVD, % cases	12.1	22.9	0.084
Dyslipidemia, % cases	63.6	51.8	0.144
Drugs			
Antihypertension, % consumers	58.6	72.3	0.076
Antithrombotic, % consumers	21.2	38.6	0.016
Cholesterol-lowering drugs, % consumers	49.5	37.3	0.135
Gastric protector, % consumers	24.2	24.1	1
Thyroid hormone therapy, % consumers	8.1	15.7	0.173
Antidepressant, % consumers	16.2	24.1	0.249
Bronchodilators, % consumers	4	8.4	0.354
Opioid analgesics, % consumers	10.1	14.5	0.503
Other, % consumers	62.6	74.7	0.114
Toxic habits			
Smoking, % cases	6.1	8.4	0.366
Previous smoking, % cases	52.5	42.2	
Occasional drinking, % cases	58.8	53	0.177
Daily drinking, % cases	24.7	19.3	
Blood pressure			
Systolic blood pressure, mmHg	131.76 (16.32)	140.69 (19.74)	0.002
Diastolic blood pressure, mmHg	77.98 (9.53)	81.84 (9.52)	0.009
Blood chemistry			
HbA1c, %	6.02 (0.43)	6.21 (0.47)	0.006
Glucose, mg/dL	104.04 (12.76)	110.21 (13.8)	0.002
Insuline, mU/L	13.97 (5.92)	19.39 (8.49)	<0.001
Total cholesterol, mg/dL	197.23 (34.11)	202.45 (31.79)	0.29
Cholesterol- HDL, mg/dL	49.94 (11.25)	52.39 (14.51)	0.212
Cholesterol-LDL, mg/dL	125.69 (29.4)	127.95 (26.55)	0.594
Tryglicerides, mg/dL	113.4 (60.12)	116.45 (47.21)	0.703
AdipoQ, µg/mL	10.89 (5.45)	11.15 (5.72)	0.762
Anthropometrics			
Weight, kg	73.06 (9.92)	86.74 (11.83)	<0.001
BMI, kg/m ²	26.62 (2.06)	33.54 (3.65)	<0.001
Waist, cm	95.2 (8.91)	107.51 (10.17)	<0.001
Hips, cm	102.42 (5.78)	114.75 (8.76)	<0.001
Percentage approximation ideal weight, %	120.63 (9)	151.81 (16.75)	<0.001
Body composition	NOB (n=51)	OB (n=37)	P-value
Centile fat mass, index	85.25 (18.5)	94.38 (7.64)	0.002
Fat mass total, %	37.8 (5.05)	44.44 (6.17)	<0.001
Fat mass total, gr	26263.12 (4691.9)	36422.38 (5538.48)	<0.001
Fat mass, torso %	42.31 (4.87)	49.69 (5.83)	<0.001
Fat mass in abdomen, %	47.68 (5.63)	54.58 (6.05)	<0.001
Fat mass in hips, %	40.42 (7.93)	47.61 (9.57)	<0.001
Lean mass, total %	56.69 (6.12)	53.40 (5.98)	<0.001

Values are expressed as mean (SD) for quantitative data, and mean % of subjects for frequency variables. Statistically significant differences are highlighted in boldface. No differences were observed in the general characteristics of the group in terms of age, sex, personal history or toxic habits. Regarding the consumption of drugs, only antithrombotics showed to be higher in the OB group. OB also demonstrated greater blood pressure, anthropometrics measures, fat mass and fat distribution in abdomen, torso and hips. Worse blood chemistry values were observed in OB, with higher HbA1c, glucose, insulin. NOB= non-obese, OB=obese, IFG= impaired fasting glycemia, DM= diabetes mellitus, HbA1c=standardized glycosylated hemoglobin NGS, AdipoQ= adiponectine, BMI= body mass index.

As expected, OB showed higher weight, body mass index (BMI) and waist and hip circumference. Ideal weight was calculated by Lorentz formula, and the percentage of approximation to it was more distant for OB (Table 1).

Total fat amount and the accumulation of fat in different parts of the body, as well as the centile in which they were classified, were higher in the OB than in NOB group. On the other hand, OB showed a lower percentage of lean mass (Table 1).

The mean of the percentage risk of developing diabetes according to FINDRISK showed a significant difference ($p < 0.001$) between groups: 40.17 ± 8.45 in the OB group and 32.40 ± 12.20 in the NOB group. Specifically, the percentage of individuals classified as having a “high” or “very high” risk of developing T2D was different ($p < 0.001$): 100% of OB participants and 77.78% of NOB participants.

OB was associated with significantly higher glucose, HbA1c and insulin (Table 1). When homeostatic model assessment for insulin resistance (HOMA-IR) was calculated, OB showed a higher index than NOB (5.37 ± 2.73 and 3.68 ± 1.85 respectively, $p < 0.001$), and for beta-cell function (HOMA- β), OB also showed a stronger compensatory effect of pancreatic function than NOB (155.06 ± 70.59 and 127.5 ± 45.81 respectively, $p = 0.003$).

3.1. Lifestyle habits

The percentage of sedentariness, classified according FAO/WHO/UNU [16], in the OB group was higher than in the NOB group, at 61.2% and 40.65%, respectively ($p = 0.01$). Moreover, OB expended a lower energy expenditure practicing physical activity than the NOB group, as calculated by METS (272.1 ± 184.7 and 330.2 ± 192.4 respectively, $p = 0.05$) and by Kcal (721.7 ± 554.9 and 537.4 ± 386 respectively, $p = 0.02$).

Harris-Benedict formula was used to evaluate daily energy expenditure with adjustments made for elderly and overweight or obese conditions, as described previously [17]. Total daily energy expenditure was lower in OB as compared to NOB (1167.71 ± 201.62 and 1402.11 ± 187.50 respectively, $p < 0.001$).

Analyses of 3-day dietary records revealed that the energy intake was similar in NOB and OB groups (1992.9 ± 564.6 and 1958.44 ± 503.9 Kcal/day, respectively), however, although both groups ingested more Kcal than they expended per day, OB showed a greater excess because their daily energy expenditure was lower (Fig. 1).

As for total calories consumed (TCC), both groups followed a diet rich in lipids (>35% of TCC) and poor in carbohydrates (<45% of TCC). Although both groups consumed a low percentage of carbohydrates, they exceeded for the recommendation for simple sugars (>10% of TCC) [18,19]. Furthermore, in terms of lipid consumption, both groups consumed an excess of saturated fatty acids (SFA) (>7% of TCC) but a correct proportion of unsaturated fatty acids (UFA) to polyunsaturated fatty acids (PUFA) (<10% of TCC), and monounsaturated fatty acids (MUFA) (10–20% of TCC) [2].

Differences between groups also appear in nutrient consumption (Table 2). NOB showed a lower intake of cholesterol and a higher intake of PUFA, especially omega 3 fatty acids (PUFA ω 3); in particular, NOB consumed more linolenic acid but also more linoleic acid, an omega 6 fatty acid (PUFA ω 6). For their part, OB showed less protein consumption (gr/Kg weight).

Other nutrients were considered in the analyses (data not shown), revealing a better quality diet in NOB due to a higher consumption of micronutrients with antioxidant properties, such as phytosterols ($p = 0.038$), available organic acid ($p = 0.011$), luteine ($p = 0.034$), citric acid ($p = 0.015$) and tocopherols ($p < 0.005$).




Fig. 1. Energy balance from each group comparing energy intake recorded from dietary records and energy expended, calculated taking into consideration basal energy expenditure and physical activity. While no difference in energy intake was observed, the NOB group expended more energy than OB. The difference between energy expended and energy ingested was different in each group and also between groups, being higher in OB. NOB= on-obese, OB=obese.

3.1.1. Clusters

Nutrient pattern analysis by means of PCA showed that participants were distributed among six clusters (Supplementary Figure S1).

From them, four different dietary patterns were clearly identified on the first principal planes (Fig. 2), with two different subpatterns observed in each group. NOB showed, on the one hand, subpattern NOB-2, based on amino acids (A) and SFA and MUFA (B), and, on the other, subpattern NOB-3 based on PUFA, in particular PUFA ω 3 (C), antioxidant nutrients (D) and based on minerals and purines (E). On the contrary, the OB group showed subpattern OB-5 based on foods rich in PUFA (F), vitamin B and some minerals (G), antioxidant nutrients (H) and based on other minerals and purines (I). The remaining subpattern of this group, OB-6, was characterized by amino acids (J), other antioxidant nutrients (K), B-group

Table 2
Nutrient intake differences between groups.

	NOB (n=86)	OB (n=64)	P-value
Carbohydrates, kcal % diet	39.42 (6.61)	38.8 (6.42)	0.553
Simple sugar, gr	78.26 (27.67)	80.04 (29.65)	0.963
Starch, gr	86.53 (33)	87.11 (34.61)	0.967
Soluble fiber, gr	4.96 (1.92)	4.69 (1.98)	0.341
Insoluble fiber, gr	9.52 (4.15)	8.85 (3.77)	0.301
Lipids, kcal % diet	40.88 (5.99)	40.12 (6.02)	0.599
Cholesterol, gr	278.13 (148.15)	308.1 (115.51)	0.03
SFA, %	11.82 (2.49)	11.5 (2.6)	0.367
MUFA, %	19.3 (4.02)	19.43 (4.14)	0.527
PUFA, %	6.28 (1.79)	5.61 (1.42)	0.008
PUFA ω 6, gr	11.01 (5.04)	9.71 (4.06)	0.115
Linoleic acid C18:2, %	4.82 (1.49)	4.35 (1.3)	0.031
PUFA ω 3, gr	2.8 (1.61)	2.23 (1.26)	0.012
Linolenic acid C18:3, %	0.82 (0.43)	0.67 (0.28)	0.003
Proteins, kcal diet	17.04 (2.97)	17.86 (3.24)	0.22
Proteins, gr/Kg weight	1.07	0.97	0.004

Values are expressed as mean (SD). Statistically significant differences are highlighted in boldface.

No differences were observed in the consumption of carbohydrates and total lipids. Only for one of three macronutrients, OB showed a lower consumption of proteins considering gr of protein/Kg of weight. The OB group also demonstrated a higher intake of cholesterol and less PUFA, linoleic acid, PUFA ω 3 and linolenic acid. NOB= non-obese, OB=obese, SFA= saturated fatty acids, MUFA= monounsaturated fatty acids, PUFA=polyunsaturated fatty acids, ω 6= omega 6, ω 3=omega.




Fig. 2. PCA analysis of different nutrients found in each dietary pattern explaining the 30% of NOB and the 51% of OB in dimensions 1 and 2. NOB= on-obese, OB=obese. Groups of nutrients: A: Alanine, Serine, Arginine, Glycine, Proline, Tyrosine, Histidine, Isoleucine, Lysine, Methionine, Threonine, Tryptophane, Phenylalanine, Leucine, Valine. B: Palmitoleic C16:1, Palmitic C16:0, Oleic C18:1 cis 9, Lipids, MUFA, SFA. C: PUFA, PUFA ω 6, PUFA ω 3, Linoleic C18:2. D: Vegetable fiber, Vit. E, Vit. K, TRAP, Oxalic acid, Tocopherol total, α -Tocopherol. E: Nickel, Chrome, Fluor, Magnesium, Potassium, Adenine, Guanine. F: Linolenic C18:3, PUFA ω 3. G: Calcium, Vit. B2, Iron. H: Vit K, TRAP, Oxalic acid, Vegetable fiber. I: Nickel, Chrome, Fluor, Magnesium, Potassium, Adenine, Guanine. J: Alanine, Serine, Arginine, Glycine, Proline, Tyrosine, Histidine, Isoleucine, Lysine, Methionine, Threonine, Tryptophane, Phenylalanine, Leucine, Valine. K: Fiber, Insoluble fiber, Soluble fiber, Vit. E, Tocopherol total, α -Tocopherol, Folate. L: Vit. B1, Vit. B9, Vit. B7, Vit. B3, Coline. M: PUFA, PUFA ω 6, Linoleic C18:2, SFA, Stearic C18:0, FA cis, Palmitoleic C16:1, Palmitic C16:0, Oleic C18:1 cis 9, Lipids, MUFA, SFA. N: Carbohydrates, Simple sugar.

vitamins (L), and PUFA, specifically omega 6, but also SFA and MUFA (M) and carbohydrates, especially simple sugars (N). The two remaining clusters (NOB-1 and OB-4), did not show any clear dietary pattern when focusing on 6 dimensions. Clusters NOB-3 and OB-5 were similar in nutrient composition, while NOB-2 and OB-6 also showed very similar patterns.

3.2. Microbiota

Prevotella relative abundance was three times higher in NOB than in OB subjects ($p = 0.047$). No statistically significant differences were found in the abundance of the other bacterial groups analyzed comparing obese and non-obese subjects (Supplementary Table S1).

3.2.1. Clusters

Differences for microbiota population were observed according to the four previously described clusters of dietary patterns (Table 3).

When comparing these four clusters, no significant differences were found. However, in NOB-2, *Prevotella* load tended to be higher

than in any of the other clusters ($p = 0.052$), and also a trend towards differences in *F.prausnitzii* abundance were observed between clusters ($p = 0.059$) (Table 3 and Supplementary Figure S2). Within OB, those in the OB-5 cluster had a higher load of both lactic acid bacteria ($p = 0.053$) and *F. prausnitzii* ($P = 0.042$) than subjects in cluster OB-6. In contrast, within NOB, the NOB-2 cluster tended to have a higher load of *Prevotella* than cluster NOB-3, but without reaching statistical significance ($P = 0.073$).

More similarities were observed between clusters NOB-3 and OB-5 and between clusters NOB-2 and OB-6 than between clusters within the same group (NOB-2 and NOB-3, OB-5 and OB-6).

3.3. T2D risk

3.3.1. Clusters

Once we observed the differences among the 4 clusters in terms of dietary patterns and their effects on gut microbiota, we studied if these differences might play a role in the risk of developing T2D.

The risk percentage of developing T2D varied among the four clusters ($p = 0.005$), however, the differences did not come from comparisons between clusters within the same group (NOB-2 and

Table 3
Comparison of microbiota populations between clusters selected in dietary patterns described.

	NOB -2 (n=22)	NOB -3 (n=5)	OB -5 (n=8)	OB -6 (n=24)	P-value ^a	P-value ^b	P-value ^c
Firmicutes	206 [126.7–370.2]	199.1 [34.33–210.1]	193.1 [135.5–227.5]	146.1 [102.9–478.3]	0.960	0.625	0.902
Bacteroidetes	42.17 [15.09–63.51]	54.06 [34.29–67.62]	26.34 [20.06–35.53]	37.61 [17.06–66.31]	0.585	0.454	0.418
<i>Prevotella</i>	2.261 [0.181–10.91]	0.026 [0.025–0.055]	0.035 [0.022–0.703]	0.052 [0.014–1.927]	0.052	0.073	0.902
Lactic acid bacteria	0.077 [0.026–0.235]	0.366 [0.006–0.586]	0.31 [0.068–19.56]	0.056 [0.027–0.162]	0.292	0.819	0.053
<i>Faecalibacterium prausnitzii</i>	28.78 [11.78–61.96]	93.6 [23.39–108.2]	104.1 [54.44–189.9]	26.69 [11.36–79.25]	0.059	0.204	0.042
<i>Escherichia coli</i>	0.188 [0.06–2.323]	0.127 [0.072–1.306]	0.872 [0.328–10.32]	0.141 [0.033–1.862]	0.550	0.672	0.177
Firmicutes/Bacteroidetes ratio	7.73 [3–11.38]	3.11 [1.3–5.81]	7.48 [3.76–19.085]	6.25 [2.065–18.3]	0.406	0.111	0.539
<i>Faecalibacterium prausnitzii</i> / <i>Escherichia coli</i> ratio	101.04 [26.68–619.1]	106.98 [71.64–184.43]	213.02 [34.715–264.74]	175.59 [11.28–1525.895]	0.922	0.536	0.825

Values are concentrations expressed as median [IQRs]. Statistically significant differences are highlighted in boldface.

No differences when comparing the four clusters were observed, with the exception of two differences found in the significance level limit for *Prevotella* and *Faecalibacterium prausnitzii*. Also, only one difference was observed in *Prevotella* when comparing clusters from the NOB group. A significant difference was seen in *Faecalibacterium prausnitzii* and a small one in lactic acid bacteria when comparing clusters from the OB group.

NOB= non-obese, OB=obese.

^a Comparison among 4 clusters.

^b Comparison between clusters 2 and 3.

^c Comparison between clusters 5 and 6.

NOB-3 clusters with $p = 0.916$, OB-5 and OB-6 clusters with $p = 1000$).

We then looked at the individuals classified as having a “high” or “very high” risk of T2D and compared the clusters that were similar in dietary patterns. We observed that NOB-3 and OB-5 presented a similar T2D risk classification ($p = 0.19$), therefore, despite the difference in BMI, there was no greater risk of developing T2D in the obese group. In contrast, comparing OB-6 to NOB-2 showed more individuals in this “high” and “very high” T2D risk classification ($p = 0.003$), which means that obesity plays an important role in T2D risk when the quality of diet and intestinal microbiota are poorer.

In agreement with these results, we observed that insulinemia, as well as the compensation of β -cell function measured by HOMA- β , was not different between NOB-3 and OB-5 but was higher in OB-6 with respect to NOB-2 ($p = 0.001$, $p = 0.005$ respectively).

4. Discussion

Obesity is associated with an array of health problems and its prevalence clearly shows a gradual increase of impaired glucose metabolism, a finding that supports the important role of obesity in the development of T2D [11]. One particular study showed that an increase in BMI raised the incidence of T2D by 25% in the general population [12].

On the contrary, the relative risk of many diseases associated with a greater BMI declines with age [20]. Nevertheless, absolute mortality and health complications associated with obesity increase linearly with a corresponding increase in BMI until the age of 75 years [20]. Beyond the age of 80 years, the association between BMI and mortality becomes weakened, primarily because the elderly may have a low body weight due to unintentional weight loss, a common complication of many serious diseases, which could confound the interpretation of weight loss effects on mortality, and also due to sarcopenia, a decrease in lean mass which often accompanies functional limitations and poor health outcomes [21].

Obesity, even so, has a negative impact on older persons with an impaired health status [22]. In particular, we found that T2D risk also increases with obesity in the elderly, along with a worsening of many parameters such as levels of glucose, HbA1c and insulin, but also HOMA-IR and HOMA- β . With these results, we can also relate an increased risk of developing other diseases associated with T2D which are driven by the combination of insulin resistance and a pro-inflammatory state [23], such as CVD [24].

The relation between energy intake and expenditure is an important determinant of obesity. Therefore, the increase in total fat mass that occurs with aging must be attributable to an increase in energy intake, a decrease in energy expenditure, or both.

Physical activity reduces with age, and it has been estimated that it accounts for about one-half of the decrease in total energy expenditure that occurs with aging [25]. Therefore, it is likely that a decrease in energy expenditure is an important contributor to the gradual increase in body fat observed with age. Both groups in our study were characterized by lower energy expenditure than found in a group of people ≥ 60 years old described in a previous study [26]. Also, we found that the OB group performed less physical activity than the NOB group, in accordance with previous results in which BMI was found to be inversely related to measured physical performance in older persons [27].

In the last decade, sedentary behavior has emerged as a new risk factor for health [28].

Findings from a study in the UK reported that objectively measured sedentary time increased with age [29]. Indeed, our study OB group spent more time performing sedentary activities,

which has been associated with higher plasma glucose [26]. In addition, excess body fat mass and a BMI of ≥ 30 in older individuals have been associated with physical dysfunction and are predictive of a decline in functional status and future disability [30].

The elderly prediabetic population in our study showed an elevated food intake in relation to daily energy expenditure. The calories they ingested were higher than in a previous study performed in the same region, Catalonia-Spain, in healthy individuals aged between 65 and 75 years [31]. In addition, according to the American Diabetes Association (ADA), healthy low-calorie eating patterns should be recommended for patients at a high risk of developing T2D [2].

The percentage of TCC from carbohydrates ingested by our population did not exceed recommendations [18,19]. Nevertheless, the percentage of TCC from simple sugars was above the recommendations [18,19], which is described that, independently the mechanisms by which the sugar intake might promote weight gain by increasing energy consumption to an extent that exceeds energy output and distorts energy balance [32], the increased consumption of simple sugar increase hyperinsulinaemia [33]. In addition, total fiber was less than previously described in same region population [30,34], and soluble fiber in particular was below the recommendations [35,36].

The percentage of TCC from lipids was higher than the recommendations [34–36], but recent evidence suggests that quality of fats is prior than quantity [37]. In this regard, dietary patterns characterized by a low intake of SFA and a higher intake of UFA are protective and promote a beneficial effect on insulin sensitivity [38]. In our study, comparing to recommendations, for both groups ingested SFA was greater [36], but amounts of MUFA and PUFA were close to it [35]. Specifically, the intake of PUFA $\omega 3$, in particular, linolenic acid, which have been associated to an improvement to cardiovascular health [39,40], was higher in the NOB group. Meanwhile, the intake of cholesterol was greater in the OB group, exceeding the recommendations [35], which has been associated with an increase in the incidence of CVD.

The quantity and also the percentage of TCC from proteins were correct in both OB and NOB groups, but when taking body weight into consideration, the OB showed lower intake than NOB group. A modest increase in protein assists in achieving and maintaining weight reduction and also helps to minimize loss of lean mass [36].

As for micronutrients, different parameters related to antioxidant power consumed were lower for the OB group. Among them, less intake of tocopherol, which a cross-sectional study found to have an inverse correlation with fasting plasma insulin concentrations [41], luteine, which the same study found inversely correlated with insulin resistance [42], and others such as organic and citric acids, which are very present in plant-based foods. In fact, green leafy vegetables, which are the most rich in these nutrients, demonstrated a capacity to reduce the risk of developing T2D, as an increased consumption of them was associated with a decrease in the incidence of T2D [42].

Subsequently, two different dietary patterns were observed in each group. Subgroups NOB-2 and OB-6 showed similar nutrient intake patterns rich in amino acids and B-group vitamins, SFA and MUFA, but pattern from the subgroup OB-6 was characterized to involved also the consumption of PUFA $\omega 6$, carbohydrates, and, especially, simple sugar. Another similar nutrient intake pattern was observed between subgroups NOB-3 and OB-5 characterized by the consumption of PUFA $\omega 3$, antioxidant nutrients, minerals and purines, but while the former was rich in PUFA $\omega 6$, the latter was rich in B-group vitamins and some minerals.

The effect of obesity on elderly prediabetic subjects seems to not have an impact on gut microbiota composition. Only *Prevotella* load, previously associated with Mediterranean diet and plant-based

diets [43] and with a polyphenols-rich diet [44], was higher in NOB. This is related to a higher consumption of PUFA and antioxidant nutrients such as polyphenols [45], which corresponded to the differences observed in nutrient intake between NOB and OB.

Previous studies have documented changes in the gut microbiota taken from subjects with obesity or certain dietary patterns and analysed for bacterial load indicators. Since no major shifts in gut microbiota have been observed in our cohort, our results could suggest that in these population groups nutrient intake has a stronger effect than obesity in modulating the composition of gut microbiota.

Interestingly, differences were observed between clusters of patients based on dietary patterns. *Prevotella* was found to be especially elevated in the NOB-2 cluster following a more unhealthy diet. These bacteria have been found in elevated concentrations in T2D patients [46] and in prediabetic populations following a diet rich in fat and protein [47]. But, on the other hand, this genus has been positively correlated with a higher consumption of oleic acid and inversely correlated with animal protein and animal fats [48]. Concretely, *Prevotella* is linked with the Mediterranean diet probably due to consuming plant-foods rich in lipids of the MUFA type, such as olive oil and nuts, while the amino acid and SFA dietary patterns probably indicate a non-animal origin, thereby creating similarities with the dietary patterns found in the NOB-2 group.

Prevotella appeared even slightly higher in the OB-6 cluster, but the reason for this could also be due to a diet rich in carbohydrates and simple sugars [49], suggesting that high proportions of these bacteria can lead to obesity in high-caloric Western diets [50]. This could explain the lower load of lactic acid bacteria found in this cluster in comparison with OB-5, due to the dietary patterns of OB-6 should be linked with Western diets rich in animal protein and fat and low in fibre which have demonstrated a decrease in these bacteria load [51]. In contrast, the OB-5 cluster shows a very high load of lactic acid bacteria, which has been related with a fish-oil rich diet in mice [52] and can be linked to the dietary pattern of this cluster through PUFA ω 3. A high load of this bacterial group has also been observed in a lactose-rich diet [53], which can be also linked to the presence of vitamin B2 and specific minerals such as calcium and iron, among others, in this cluster. Different lactic acid bacteria have been related with an improvement in glucose metabolism and a reduction in chronic inflammation [54] and have been found in reduced quantities in patients with T2D and obesity [55]. Finally, *F. prausnitzii* was higher in NOB-3 and OB-5. The dietary patterns of both subgroups were rich in PUFA ω 3, antioxidant nutrients and were compounded for the same minerals and purines. These results can be linked with previously studies where an increase in these bacteria was found due to a diet rich in fruit, legumes and some vegetables [56–58]. Also, *F. prausnitzii* has been demonstrated to decrease with hydrolysed complex carbohydrates [59], which could translate into the carbohydrates appearing in the dietary pattern of OB-6 that come from simple carbohydrates found in ultra-processed foods [60]. A possible association between changes in this species of bacteria and insulin sensitivity has been previously suggested [61–63], and depletion of this species has been reported in patients with T2D [64–66]. Supporting the influence of *F. prausnitzii* on T2D, a study analysing *F. prausnitzii* profiles in obese subjects with and without T2D has suggested that differences in this species phylotypes may lead to differences in inflammatory status in the host, thus having an influence on disease development [65]. Furthermore, its anti-inflammatory properties have been consistently demonstrated [66].

We have shown in our study that individuals from a specific group of elderly people with preDM and obesity present a higher risk of developing T2D and worse metabolic outcomes. However, in

those individuals who present a healthier dietary pattern and gut microbiota, the presence of obesity does not appear to modify insulinemia or the compensatory function of β cells, nor does it increase the risk of developing T2D.

Therefore, we can conclude that a healthy dietary pattern and the growth-promoting beneficial and growth-inhibiting disadvantageous gut microbiota populations linked to it provide protection against the development of T2D in an obese population with advanced age and preDM.

Clinical trial

This study was registered in clinicaltrials.gov with identifier number: NCT03557541.

Statement of authorship

DA Díaz-Rizzolo and R Gomis designed the study. DA Díaz Rizzolo and C Colungo performed clinical visits. DA Díaz Rizzolo and A Serra collected samples and codified data. M López-Siles and M Martínez-Medina were responsible for microbiota extraction and determination. B Kostov Formal analysis participated in all data analysis. DA Díaz-Rizzolo, A Sisó-Almirall and R Gomis were responsible for interpretation the results. All of the authors participated in the critical revision of the manuscript. All authors gave final approval of the version to be submitted.

Funding sources

This research received a grant from Recercaixa 2013 and used funds from the Fundación AstraZeneca Chair in Diabetes Innovation at IDIBAPS.

Acknowledgements

The authors wish to thank: the nurses from the three primary care centers collaborating on this study, Ms. Eva Sanchez Romero for calling potential participants, Ms. Lucía Daniela Rizzolo Brime for assistance in introducing dietary records, Dr. Vanessa Blanc and Ms. Marta Gonzalez from Dentaïd for providing *Prevotella* intermedia NCTC 13070, and Ms. Ariadna Peñafiel for assistance in qPCR analysis. We also wish to thank Kimberly Katte for editorial assistance. The authors wish to thank the Fundación AstraZeneca Chair in Diabetes Innovation at IDIBAPS for its support for the present work. Finally, we thank CIBERDEM, an initiative of the Instituto de Salud Carlos III, also the support of the Secretaria d'Universitats i Recerca del Departament d'Empresa i Coneixement de la Generalitat de Catalunya.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.clnu.2019.02.035>.

References

- [1] International Diabetes Federation. IDF diabetes atlas 2017. 8th ed. 2017 Available from: www.diabetesatlas.org.
- [2] American Diabetes Association. Standards of medical care in diabetes—2018. *Diabetes Care* 2018;41(Suppl. 1).
- [3] SM Grundy. Pre-diabetes, metabolic syndrome and cardiovascular risk. *JACC (J Am Coll Cardiol)* 2012;635–43.
- [4] Forouhi NG, Luan J, Hennings S, Wareham NJ. Incidence of Type 2 diabetes in England and its association with baseline impaired fasting glucose: the Ely study 1990–2000. *Diabet Med* 2007;24(2):200–7.

- [5] Nathan DM, Davidson MB, DeFronzo RA, Heine RJ, Henry RR, Pratley R, et al., American Diabetes Association. Impaired fasting glucose and impaired glucose tolerance: implications for care. *Diabetes Care* 2007;30(3):753–9.
- [6] Narayan KMV, Boyle JP, Geiss LS, Saaddine JB, Thompson TJ. Impact of recent increase in incidence on future diabetes burden. *Diabetes Care* 2006;29(9):2114–6.
- [7] Han TS, Tajar A, Lean MEJ. Obesity and weight management in the elderly. *Br Med Bull* 2011;97(1):169–96.
- [8] Narayan KMV, Boyle JP, Thompson TJ, Gregg EW, Williamson DF. Effect of BMI on lifetime risk for diabetes in the U.S. *Diabetes Care* 2007;30(6):1562–6.
- [9] Lissner L, Odell PM, D'Agostino RB, Stokes J, Kreger BE, Belanger AJ, et al. Variability of body weight and health outcomes in the framingham population. *N Engl J Med* 1991;324(26):1839–44.
- [10] Blair SN. Evidence for success of exercise in weight loss and control. *Ann Intern Med* 1993;119(7 Pt 2):702–6.
- [11] Heymsfield SB, Cristina Gonzalez MC, Shen W, Redman L, Thomas D. Weight loss composition is one-fourth fat-free mass: a critical review and critique of this widely cited rule. *Obes Rev* 2014;15(4):310–21.
- [12] Fukushima Y, Kurose S, Shinno H, Thu HC, Takao N, Tsutsumi H, et al. Importance of lean muscle maintenance to improve insulin resistance by body weight reduction in female patients with obesity. *Diabetes Metab J* 2016;40(2):147–53.
- [13] Darmon P, Kaiser MJ, Bauer JM, Sieber CC, Pichard C. Restrictive diets in the elderly: never say never again? *Clin Nutr* 2010;29(2):170–4.
- [14] Graf D, Di Cagno R, Fåk F, Flint HJ, Nyman M, Saarela M, et al. Contribution of diet to the composition of the human gut microbiota. *Microb Ecol Health Dis* 2015;26:10.
- [15] Brown K, DeCoffe D, Molcan E, Gibson DL. Diet-induced dysbiosis of the intestinal microbiota and the effects on immunity and disease. *Nutrients* 2012;4(8):1095–119.
- [16] FAO/WHO-OMS/UNU Expert Consultation Report. Energy and protein requirements. Technical report series 724. Ginebra: WHO/OMS; 1985.
- [17] Guía de Práctica Clínica sobre Diabetes tipo 2. Madrid: plan Nacional para el SNS del MSC. Agencia de Evaluación de Tecnologías Sanitarias del País Vasco; 2008. Guías de Práctica Clínica en el SNS: OSTEBA N° 2006/08. Grupo de trabajo de la Guía de Práctica Clínica sobre Diabetes tipo 2.
- [18] U.S. Department of Agriculture, U.S. Department of Health and Human Services. Dietary guidelines for Americans. 8th ed. Washington, DC: U.S. Government Printing Office; 2015–2020. [Accessed 8 March 2016].
- [19] EFSA. Panel on dietetic products, nutrition and allergies (NDA). Scientific opinion on dietary reference values for carbohydrates and dietary fiber. *EFSA J* 2010;8(3):1462.
- [20] Stevens J, Cai J, Pamuk ER, Williamson DF, Thun MJ, Wood JL. The effect of age on the association between body-mass index and mortality. *N Engl J Med* 1998;338:1–7.
- [21] Stenholm S, Harris TB, Rantanen T, Visser M, Kritchevsky SB, Ferrucci L. Sarcopenic obesity – definition, etiology and consequences. *Curr Opin Clin Nutr Metab Care* 2008;11:693–700.
- [22] Blaum CS, Xue QL, Michelon E, Semba RD, Fried LP. The association between obesity and the frailty syndrome in older women: the Women's Health and Aging studies. *J Am Geriatr Soc* 2005;53:927–34.
- [23] Feingold KR, Grunfeld C. Diabetes and dyslipidemia. In: Feingold KR, Anawalt B, Boyce A, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2019 Jan 3. 2000. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK305900/>.
- [24] Carr DB, Utzschneider KM, Hull RL, Kodama K, Retzlaff BM, Brunzell JD, et al. Intra-abdominal fat is a major determinant of the national cholesterol education program adult treatment panel III criteria for the metabolic syndrome. *Diabetes* 2004;53(8):2087–94.
- [25] Tzankoff SP, Norris AH. Effect of muscle mass decrease on age-related BMR changes. *J Appl Physiol* 1977;43:1001e6.
- [26] Gennuso KP, Gangnon RE, Matthews CE, Thraen-Borowski KM, Colbert LH. Sedentary behavior, physical activity, and markers of health in older adults. *Med Sci Sports Exerc* 2013;45:1493.
- [27] Apovian CM, Frey CM, Rogers JZ, McDermott EA, Jensen GL. Body mass index and physical function in obese older women. *J Am Geriatr Soc* 1996;44:1487e8.
- [28] Katzmarzyk PT. Physical activity, sedentary behavior, and health: paradigm paralysis or paradigm shift? *Diabetes* 2010;59:2717–25.
- [29] Davis MG, Fox KR, Hillsdon M, Sharp DJ, Coulson JC, Thompson JL. Objectively measured physical activity in a diverse sample of older urban UK adults. *Med Sci Sports Exerc* 2001;43:647–54.
- [30] Davison KK, Ford ES, Cogswell ME, Dietz WH. Percentage of body fat and body mass index are associated with mobility limitations in people aged 70 and older from NHANES III. *J Am Geriatr Soc* 2002;50:1802e9.
- [31] Serra-Majem L, Ribas-Barba L, Salvador G, Jover L, Raidó B, Ngo J, et al. Trends in energy and nutrient intake and risk of inadequate intakes in Catalonia, Spain (1992–2003). *Publ Health Nutr* 2007;10:1354–67.
- [32] Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *Br J Med* 2013;346:e7492.
- [33] Johnson RJ, Perez-Pozo SE, Sautin YY, Manitius J, Sanchez-Lozada LG, Feig DL, et al. Hypothesis: could excessive fructose intake and uric acid cause type 2 diabetes? *Endocr Rev* 2009;30:96–116.
- [34] Molina-Montes E, Sanchez-Cantalejo E, Martinez C, Contreras JM, Molina E, Sánchez MJ. Compliance with dietary and nutrient recommendations in the European prospective investigation into cancer and nutrition (EPIC)-Granada cohort at recruitment. *Nutr Hosp* 2012;27:572–82.
- [35] Serra-Majem L, J. aranceta; SENC working group on nutritional objectives for the Spanish population. Spanish society of community nutrition. Nutritional objectives for the Spanish population. Consensus from the Spanish society of community nutrition. *Publ Health Nutr* 2001;4(6A):1409–13.
- [36] Joslin Diabetes Center & Joslin Clinic. Clinical nutrition guideline for overweight and obese adults with type 2 diabetes, prediabetes or those at high risk for developing type 2 diabetes 2011.
- [37] Wang DD, Li Y, Chiuve SE, Stampfer MJ, Manson JE, Rimm EB, et al. Association of specific dietary fats with total and cause-specific mortality. *JAMA Intern Med* 2016;176:1134–45.
- [38] Riserus U, Willett WC, Hu FB. Dietary fats and prevention of Type 2 diabetes. *Prog Lipid Res* 2009;48(1):44–51.
- [39] Connor WE. Importance of n-3 fatty acids in health and disease. *Am J Clin Nutr* 2000;71(Suppl. 1):171S–5S.
- [40] Djoussé L, Pankow JS, Eckfeldt JH, Folsom AR, Hopkins PN, Province MA, et al. Relation between dietary linolenic acid and coronary artery disease in the national heart, lung, and blood institute family heart study. *Am J Clin Nutr* 2001;74:612–9.
- [41] Facchini FS, Humphreys MH, DoNascimento CA, Abbasi F, Reaven GM. Relation between insulin resistance and plasma concentrations of lipid hydroperoxides, carotenoids, and tocopherols. *Am J Clin Nutr* 2000;72:776–9.
- [42] Carter P, Gray LJ, Troughton J, Khunti K, Davies MJ. Fruit and vegetable intake and incidence of Type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ* 2010;341:c4229.
- [43] DeFilippis F, Pellegrini N, Vannini L, Jefery IB, La Storia A, Laghi L, et al. High-level adherence to a Mediterranean diet beneficially impacts the gut microbiota and associated metabolome. *Gut* 2016 Nov;65(11):1812–21.
- [44] Queipo-Ortuno MI, Boto-Ordóñez M, Murri M, Gomez-Zumaquero JM, Clemente-Postigo M, Estruch R, et al. Influence of red wine polyphenols and ethanol on the gut microbiota ecology and biochemical biomarkers. *Am J Clin Nutr* 2012;95:1323–34.
- [45] Natchochiy SM, Redman EK. Mediterranean diet and cardio protection: the role of nitrite, polyunsaturated fatty acids, and polyphenols. *Nutrition* 2011;27:733–44.
- [46] Leite AZ, Rodrigues NC, Gonzaga MI, Paiolo JCC, de Souza CA, Stefanutto NAV, et al. Detection of increased plasma interleukin-6 levels and prevalence of *Prevotella copri* and *Bacteroides vulgatus* in the feces of type 2 diabetes patients. *Front Immunol* 2017;8:1107.
- [47] Egshatyan L, Kashtanova D, Popenko A, Tkacheva O, Tyahkt A, Alexeev D, et al. Gut microbiota and diet in patients with different glucose tolerance. *Endocr Connect* 2016;5(1):1–9.
- [48] Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, et al. Linking long-term dietary patterns with gut microbial enterotypes. *Science* 2011;334(6052):105–8.
- [49] Durbán A, Abellán JJ, Jiménez-Hernández N, Latorre A, Moya A. Daily follow-up of bacterial communities in the human gut reveals stable composition and host-specific patterns of interaction. *FEMS Microbiol Ecol* 2012;81:427–37.
- [50] Durbán A, Abellán JJ, Latorre A, Moya A. Effect of dietary carbohydrate restriction on an obesity-related *Prevotella*-dominated human fecal microbiota. *Metagenomics* 2013;2:1–4.
- [51] Reddy BS, Weisburger JH, Wynder EL. Effects of high risk and low risk diets for colon carcinogenesis on fecal microflora and steroids in man. *J Nutr* 1975;105:878–84.
- [52] Caesar R, Tremaroli V, Kovatcheva-Datchary P, Cani PD, Bäckhed F. Crosstalk between gut microbiota and dietary lipids aggravates WAT inflammation through TLR signaling. *Cell Metabol* 2015;22:658–68.
- [53] Francavilla R, Calasso M, Calace L, Siragusa S, Ndagijimana M, Vernocchi P, et al. Effect of lactose on gut microbiota and metabolic outcome of infants with cow's milk allergy. *Pediatr Allergy Immunol* 2012;23:420–7.
- [54] Stenman Lk, Waget A, Garret C, Klopp P, Burcelin R, Lahtinen S. Potential probiotic *Bifidobacterium animalis* ssp. *lactis* 420 prevents weight gain and glucose intolerance in diet-induced obese mice. *Benef Microbes* 2014;5:437–45.
- [55] Xu X, Hui H, Cai D. Differences in fecal *Bifidobacterium* species between patients with type 2 diabetes and healthy individuals. *Nan Fang Yi Ke Da Xue Xue Bao* 2012;32(531–533):564.
- [56] Fernando WM, Hill JE, Zello GA, Tyler RT, Dahl WJ, Van Kessel AG. Diets supplemented with chickpea or its main oligosaccharide component raffinose modify faecal microbial composition in healthy adults. *Benef Microbes* 2010;1(2):197–207.
- [57] Hooda S, Boler BM, Seroo MC, Brulc JM, MA Staeger, Boileau TW, et al. 454 pyrosequencing reveals a shift in fecal microbiota of healthy adult men consuming polydextrose or soluble corn fiber. *J Nutr* 2012;142:1259–65.
- [58] Lopez-Siles M, Khan TM, Duncan SH, Harmsen HJ, Garcia-Gil LJ, Flint HJ. Cultured representatives of two major phylogroups of human colon *Faecalibacterium prausnitzii* can utilize pectin, uronic acids, and host-derived substrates for growth. *Appl Environ Microbiol* 2012;78:420–8.
- [59] Jia W, Whitehead RN, Griffiths L, Dawson C, Waring RH, Ramsden DB, et al. Is the abundance of *Faecalibacterium prausnitzii* relevant to Crohn's disease? *FEMS Microbiol Lett* 2010;310:138–44.

- [60] Monteiro C, Cannon G, Moubarac J, Levy R, Louzada M, Jaime P. The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Publ Health Nutr* 2018;21(1):5–17.
- [61] Khan MT, Nieuwdorp M, Backhed F. Microbial modulation of insulin sensitivity cell metabolism 2014;20(5):753–60.
- [62] Vrieze A, Out C, Fuentes S, Jonker L, Reuling I, Kootte RS. Impact of oral vancomycin on gut microbiota, bile acid metabolism, and insulin sensitivity. *J Hepatol* 2014;60(4):824–31.
- [63] Karlsson FH, Tremaroli V, Nookaew I, Bergstrom G, Behre CJ, Fagerberg B, et al. Gut metagenome in European women with normal, impaired and diabetic glucose control. *Nature* 2013;498(7452):99–103.
- [64] Furet J, Kong L, Tap J, Poitou C, Basdevant A, Bouillot J, et al. Differential adaptation of human gut microbiota to bariatric surgery-induced weight loss. *Diabetes* 2010;59:3049–57.
- [65] Hippe B, Remely M, Aumueller E, Pointner A, Magnet U, Haslberger AG. Faecalibacterium prausnitzii phylotypes in type two diabetic, obese, and lean control subjects. *Benef Microbes* 2016;7(4):511–7.
- [66] Miquel S, Martín R, Rossi O, Bermúdez-Humarán LG, Chatel JM, Sokol H, et al. Faecalibacterium prausnitzii and human intestinal health. *Curr Opin Microbiol* 2013;16(3):255–61.