



# Metformin Should Not Be Used to Treat Prediabetes

Mayer B. Davidson

*Diabetes Care* 2020;43:1983–1987 | <https://doi.org/10.2337/dc19-2221>

Based on the results of the Diabetes Prevention Program Outcomes Study (DPPOS), in which metformin significantly decreased the development of diabetes in individuals with baseline fasting plasma glucose (FPG) concentrations of 110–125 vs. 100–109 mg/dL (6.1–6.9 vs. 5.6–6.0 mmol/L) and A1C levels 6.0–6.4% (42–46 mmol/mol) vs. <6.0% and in women with a history of gestational diabetes mellitus, it has been suggested that metformin should be used to treat people with prediabetes. Since the association between prediabetes and cardiovascular disease is due to the associated nonglycemic risk factors in people with prediabetes, not to the slightly increased glycemia, the only reason to treat with metformin is to delay or prevent the development of diabetes. There are three reasons not to do so. First, approximately two-thirds of people with prediabetes do not develop diabetes, even after many years. Second, approximately one-third of people with prediabetes return to normal glucose regulation. Third, people who meet the glycemic criteria for prediabetes are not at risk for the microvascular complications of diabetes and thus metformin treatment will not affect this important outcome. Why put people who are not at risk for the microvascular complications of diabetes on a drug (possibly for the rest of their lives) that has no immediate advantage except to lower subdiabetes glycemia to even lower levels? Rather, individuals at the highest risk for developing diabetes—i.e., those with FPG concentrations of 110–125 mg/dL (6.1–6.9 mmol/L) or A1C levels of 6.0–6.4% (42–46 mmol/mol) or women with a history of gestational diabetes mellitus—should be followed closely and metformin immediately introduced only when they are diagnosed with diabetes.

The Diabetes Prevention Program (DPP) studied the effect of an intensive lifestyle intervention and metformin on the development of diabetes in a cohort of people with an increased risk for diabetes (termed prediabetes). After a mean of 2.8 years of follow-up, 31% fewer metformin-treated individuals developed diabetes than individuals in the control group (1). Eighty-six percent of members of the metformin and placebo groups agreed to be followed and entered the Diabetes Prevention Program Outcomes Study (DPPOS). The placebo was discontinued and metformin (850 mg b.i.d.) was unmasked and continued. The 15-year follow-up results in the DPPOS metformin-treated group recently showed significantly less development of diabetes in participants with higher baseline fasting plasma glucose (FPG) concentrations (110–125 vs. 100–109 mg/dL [6.1–6.9 vs. 5.6–6.0 mmol/L]) (2), in those with A1C levels 6.0–6.4% (42–46 mmol/mol) vs. <6.0%, and in women with a history of gestational diabetes mellitus (2). An accompanying editorial (3) invited arguments discussing whether people meeting the criteria for prediabetes should be treated with metformin. Since 33.9% of the population over 18 years of age in the U.S., 84.1 million people, have prediabetes (4), use of metformin to treat them would increase drug costs considerably for payers as well as for many individuals. This Perspective will argue against doing so.

Charles R. Drew University, Los Angeles, CA  
Corresponding author: Mayer B. Davidson,  
[mayerdavidson@cdrewu.edu](mailto:mayerdavidson@cdrewu.edu)

Received 5 February 2020 and accepted 28 April 2020

© 2020 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

See accompanying article, p. 1988.

It is instructive to review the history of diagnosing prediabetes. Before 1979, there were six different criteria for diagnosing diabetes. In that year, the National Diabetes Data Group (NDDG) published a single set of criteria for the diagnosis (FPG  $\geq 140$  mg/dL [7.8 mmol/L] or 2-h glucose concentration on an oral glucose tolerance test [OGTT]  $\geq 200$  mg/dL [11.1 mmol/L]) based on three prospective studies in subjects who had a baseline OGTT and were evaluated for diabetic retinopathy 3 to 8 years later (5). They also opined that individuals whose 2-h glucose value was  $\geq 140$  to 199 mg/dL (7.8 to 11.0 mmol/L) had impaired glucose tolerance (IGT), which indicated an increased risk for developing diabetes. No FPG criterion for diagnosing prediabetes was offered.

The NDDG criteria for diagnosing diabetes were not equally sensitive. Although 95% of all persons with an FPG concentration  $\geq 140$  mg/dL (7.8 mmol/L) also had a 2-h glucose concentration  $\geq 200$  mg/dL (11.1 mmol/L) on the OGTT, this level of concordance was not seen with all persons who had a 2-h glucose concentration  $\geq 200$  mg/dL (11.1 mmol/L). Only one-quarter to one-half of these individuals also had an FPG  $\geq 140$  mg/dL (7.8 mmol/L) (6). The American Diabetes Association (ADA) convened an Expert Committee to address this imbalance (7). Based on an analysis by the Expert Committee of the third National Health and Nutrition Examination Survey (NHANES III) and several other published studies, the new FPG criterion for diagnosing diabetes was set at  $\geq 126$  mg/dL (7.0 mmol/L), which yielded the same prevalence of diabetes as did a 2-h glucose value on the OGTT of  $\geq 200$  mg/dL (11.1 mmol/L). Since no studies defining a normal FPG concentration were known, the often-stated normal glucose value of  $< 110$  mg/dL (6.1 mmol/L) used by clinical laboratories was adopted. The FPG range of 110–125 mg/dL (6.1–6.9 mmol/L) was termed impaired fasting glucose (IFG) and joined IGT to diagnose prediabetes.

However, again there was an imbalance. Many fewer people with IFG subsequently developed diabetes compared with those who had IGT. The ADA convened another meeting of the Expert Committee to address this issue (8,9). They analyzed four populations and determined that lowering the IFG criterion to 100–125 mg/dL (5.6–6.9 mmol/L)

would greatly reduce the predictive discrepancy between IGT and IFG for the subsequent development of diabetes.

In 2008, an invited expert panel (IEP) recommended that diabetes could be diagnosed by an A1C level of  $\geq 6.5\%$  (48 mmol/mol) and also suggested that values of 6.0–6.4% (42–46 mmol/mol) required close follow-up and testing (10). In response, the ADA, the European Association for the Study of Diabetes, and the International Diabetes Federation appointed an International Expert Committee that agreed with the invited expert panel regarding the diagnosis of diabetes (if the A1C level were confirmed) (11). However, that committee also opined that because of the progressive continuum of risk of increasing glycemia below the diagnostic levels of diabetes for the subsequent development of diabetes, it was inappropriate to define a specific prediabetes risk group. The ADA subsequently adopted the recommended A1C level for diagnosing diabetes but also included an A1C criterion of 5.7–6.4% (39–46 mmol/mol) for prediabetes (12). The lower bound of the prediabetes criteria was based on modeling the estimated composite risk of developing diabetes and cardiovascular disease (CVD) using cross-sectional data from the 2005–2006 NHANES (13). However, the glycemia of prediabetes is not independently associated with CVD (14–21). Furthermore, in people who experience an acute coronary syndrome, the outcomes (length of hospital stay, 28-day readmission rate, acute pulmonary edema, 12-month recurrent acute coronary syndrome, or mortality) are no different between those with prediabetes (A1C 5.7–6.4% [39–46 mmol/mol]) or with A1C levels  $< 5.7\%$  (39 mmol/mol) (22). Rather, the association between prediabetes and CVD is due to the other risk factors for CVD that people meeting the glycemic criteria for prediabetes also have. Restricting the modeling to only the risk for developing diabetes might have influenced the prediabetes A1C criterion.

The World Health Organization (WHO) accepted the 1997 ADA IFG criterion of FPG 110–125 mg/dL (6.1–6.9 mmol/L) for prediabetes (23) but not the 2003 ADA IFG criterion of FPG 100–125 mg/dL (5.6–6.9 mmol/L) (24). Regarding the A1C criteria, the WHO adopted the ADA A1C criterion of  $\geq 6.5\%$  (48 mmol/mol) for diagnosing diabetes (if confirmed) but

stated that there was insufficient evidence to decide on A1C values  $< 6.5\%$  (48 mmol/mol) (25).

The Diabetes Canada 2018 Clinical Practice Guidelines recommended the criteria for prediabetes as IFG concentrations of 110–125 mg/dL (6.1–6.9 mmol/L) or A1C levels of 6.0–6.4% (42–46 mmol/mol) (26).

Although numerous studies have shown that glycemia is not an independent risk factor for CVD (14–21), it certainly is for the development of diabetes. However, there is no obvious threshold; the risk starts to increase beginning at FPG concentrations of 82–87 mg/dL (4.6–4.8 mmol/L) and progresses in a curvilinear fashion (27–29). For instance, the risk with the WHO IFG criterion of 110–125 mg/dL (6.1–6.9 mmol/L) is 2.1- to 11.3-fold higher than with the lower bound of the ADA IFG criterion of 100–109 mg/dL (5.6–6.0 mmol/L) (14,30,31). Similarly, the risk with the A1C IEP criterion of 6.0–6.4% (42–46 mmol/mol) is 2.0- to 6.5-fold higher than with the lower bound of the ADA A1C criterion of 5.7–5.9% (39–41 mmol/mol) (14,31).

Claims have been made that treating people with prediabetes with antihyperglycemic drugs (metformin, thiazolidinediones [TZD],  $\alpha$ -glucosidase inhibitors, glucagon-like peptide 1 agonists, basal insulin) has delayed or even prevented the development of diabetes. This is a misinterpretation of the situation. These drugs have simply treated a level of glycemia lower than the diagnostic criteria for diabetes retarding its increase to the level at which a diagnosis of diabetes would occur. After these drugs were discontinued, the prevalence of diabetes in treated individuals mirrored that in the placebo group.

An argument has been made that the difference between the placebo and metformin groups in the DPP only decreased from 31% to 25% 1–2 weeks (mean 11 days) after discontinuing metformin (a period of time that encompassed more than five half-lives of the drug), indicating that metformin caused a long-lasting, fundamental change in the pathophysiology of prediabetes (32). However, the time course of action of a drug is much more related to its tissue biologic effects than to the pharmacokinetics of its concentration in the blood. It is well established that it takes 2–4 weeks for both metformin and sulfonylureas to

exert their maximal effects when started (33–35). Although the author could find no studies examining the time course of the effect of metformin wearing off, it takes 2–4 weeks for the effect of a sulfonylurea (tolazamide) to completely dissipate (33). The facts that in the 1- to 2-week period in which metformin was discontinued 64% more subjects who had received metformin developed diabetes than those who had received a placebo (32) and that in the DPPOS the incidence of developing diabetes was similar in the original three groups of the DPP (36) strongly suggest that metformin does not fundamentally change the pathophysiology of prediabetes.

Troglitazone, a TZD that was removed from the market because of hepatic toxicity, was used for a mean of 0.9 years in the DPP (37). During this period, diabetes incidence was reduced by 75% compared with placebo but the incidence was identical to placebo after troglitazone was discontinued. In the DREAM (Diabetes Reduction Assessment With Ramipril and Rosiglitazone Medication) study, in which rosiglitazone was the TZD, 60% fewer persons developed diabetes compared with the placebo group (38). In those who had not developed diabetes during the intervention period, the rate of development of diabetes was the same in both groups during the 2- to 3-month washout period after both rosiglitazone and its placebo were discontinued (39) and 1.6 years later (40). The Outcome Reduction With Initial Glargine Intervention (ORIGIN) study compared people with CVD risk factors who also had IFG, IGT, or early type 2 diabetes and who were given either glargine insulin or placebo (41). In those who did not have diabetes at baseline, 30% and 35% developed diabetes in the glargine and placebo groups, respectively, approximately 3 months after the study ended. The pathophysiologic abnormalities of insulin resistance and progressive  $\beta$ -cell dysfunction that characterize prediabetes were not fundamentally altered by these drug treatments (42,43), which explains the lack of any long-term effects when these medications were discontinued (44).

Even so, should metformin treatment be offered to individuals whose glycemic parameters are near the diagnosis for diabetes, i.e., those with IGT or whose FPG meets the WHO IFG criterion of 110–

125 mg/dL (6.1–6.9 mmol/L) or the IEP A1C criterion of 6.0–6.4% (42–46 mmol/mol)? There are three arguments against this. First, approximately two-thirds of people with prediabetes do not develop diabetes, even after many years. In the placebo arm of the DPPOS, 65% of participants had not developed diabetes 5.7 years after the DPP had ended (45). In the Framingham Offspring Study, 69% of the cohort with prediabetes had not developed diabetes 27–30 years later (46). In people >60 years of age with prediabetes who were followed for 12 years in the Swedish National Study on Aging (47), 23% died and 13% developed diabetes. Even if all of the individuals who died had developed diabetes before doing so (highly unlikely), this would still leave 64% who did not develop diabetes.

Second, approximately one-third of people with prediabetes return to normal glucose regulation (NGR). In the DREAM study, 30% of the participants in the placebo arm returned to NGR during the 3.0 years of the study (38). After the study ended, the percent of participants who returned to NGR 1.6 years later was 38% in the placebo arm and 42% in the rosiglitazone arm (40). In the DPPOS, 24% of individuals in the placebo arm returned to NGR 5.7 years after the DPP ended (45). In a Korean population, 36% of people with prediabetes returned to NGR within 10 years (48). Even in the older population of the Swedish National Study on Aging, 23% returned to NGR (47). It is unknown how many of the 23% who died might have returned to NGR. In the Whitehall II Cohort Study (49), in which the ADA criteria were used to diagnose prediabetes, of those with IFG or IGT or diagnosed with A1C levels, 45%, 37%, and 17%, respectively, returned to NGR in 5 years. Finally, in a Cochrane Database systemic review of 47 studies of prediabetes, return to NGR ranged from 33% to 59% within 1–5 follow-up years and from 17% to 42% within 6–11 years of follow-up (50).

Third, as described previously, the diagnostic criteria for diabetes were selected because the risk for microvascular complications increased beyond that level of glycemia. Metformin, the preferred initial drug for treating patients with diabetes, is started to lower glycemia to levels that are not associated with this risk. Five studies (51–55) have shown that the development or progression of retinopathy and microalbuminuria over a

6- to 10-year period was almost nil if A1C levels were kept below 7.0% (53 mmol/mol). So, given that two-thirds of people with prediabetes do not develop diabetes over many years (45–47), and in approximately one-third glycemia returns to normal (40,45,47–50), why put people who are not at risk for the microvascular complications of diabetes when prediabetes is diagnosed on a drug (possibly for the rest of their lives) that has no immediate advantage except to lower subdiabetes glycemia to even lower levels? The authors of the Cochrane Database systemic review (50) also concluded that “practitioners should be careful about the potential implications of any active intervention for people ‘diagnosed’ with [intermediate hyperglycemia].”

This Perspective is not arguing against the benefit of delaying the development of diabetes. Rather, it is pointing out that the benefit of delay achieved with medication must be weighed against the potential adverse effects of the drug, its cost, and the important fact that a large number of people with the diagnosis of prediabetes will not develop diabetes and metformin would be of no benefit for them. The argument is that lifestyle interventions, especially weight loss in overweight and obese individuals, should be pursued rather than use of a medication.

It seems more prudent to identify individuals at the highest risk for developing diabetes—i.e., those fulfilling the WHO FPG criterion of 110–125 mg/dL (6.1–6.9 mmol/L) or those meeting the IEP A1C criterion of 6.0–6.4% (42–46 mmol/mol) or women with a history of gestational diabetes mellitus—in order to follow them closely and immediately introduce metformin when their glycemia meets the criteria for diabetes (if confirmed). Meanwhile, these individuals should be intensely counseled on lifestyle interventions to reduce the risk of developing diabetes, and the risk factors for CVD should be aggressively addressed. Although the ADA (56) and the Endocrine Society (57) recommend metformin for treatment of prediabetes, apparently most clinicians agree with the views described above because currently only 1–4% of people with prediabetes are given metformin (58,59).

---

**Duality of Interest.** No conflicts of interest relevant to this article were reported.

## References

- Knowler WC, Barrett-Connor E, Fowler SE, et al.; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403
- Diabetes Prevention Program Research Group. Long-term effects of metformin on diabetes prevention: identification of subgroups that benefited most in the Diabetes Prevention Program and Diabetes Prevention Program Outcomes Study. *Diabetes Care* 2019;42:601–608
- Cefalu WT, Riddle MC. More evidence for a prevention-related indication for metformin: let the arguments resume! *Diabetes Care* 2019;42:499–501
- Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2020. U.S. Department of Health and Human Services. Accessed 2 February 2020. Available from <https://www.cdc.gov/diabetes/data/statistics-report>
- National Diabetes Data Group. Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes* 1979;28:1039–1057
- Davidson MB. Clinical irrelevance of the current diagnostic criteria for abnormal carbohydrate metabolism in asymptomatic individuals. *Curr Opin Endocrinol Diabetes* 2005;12:437–443
- The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 1997;20:1183–1197
- Genuth S, Alberti KG, Bennett P, et al.; Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Follow-up report on the diagnosis of diabetes mellitus. *Diabetes Care* 2003;26:3160–3167
- Davidson MB. Correction to the 2010 report on the diagnosis and classification of diabetes. *Diabetes Care* 2010;33:e57
- Saudek CD, Herman WH, Sacks DB, Bergenstal RM, Edelman D, Davidson MB. A new look at screening and diagnosing diabetes mellitus. *J Clin Endocrinol Metab* 2008;93:2447–2453
- International Expert Committee. International Expert Committee report on the role of the A1C assay in the diagnosis of diabetes. *Diabetes Care* 2009;32:1327–1334
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2010;33(Suppl. 1):S62–S69
- Ackermann RT, Cheng YJ, Williamson DF, Gregg EW. Identifying adults at high risk for diabetes and cardiovascular disease using hemoglobin A1c. National Health and Nutrition Examination Survey 2005–2006. *Am J Prev Med* 2011;40:11–17
- Davidson MB, Kahn RA. A reappraisal of prediabetes. *J Clin Endocrinol Metab* 2016;101:2628–2635
- Vistisen D, Witte DR, Brunner EJ, et al. Risk of cardiovascular disease and death in individuals with prediabetes defined by different criteria: the Whitehall II Study. *Diabetes Care* 2018;41:899–906
- Bancks MP, Ning H, Allen NB, et al. Long-term absolute risk for cardiovascular disease stratified by fasting glucose level. *Diabetes Care* 2019;42:457–465
- Kiviniemi AM, Lepojärvi ES, Tulppo MP, et al. Prediabetes and risk for cardiac death among patients with coronary artery disease: the ARTEMIS Study. *Diabetes Care* 2019;42:1319–1325
- Hubbard D, Colantonio LD, Tanner RM, et al. Prediabetes and risk for cardiovascular disease by hypertension status in black adults: the Jackson Heart Study. *Diabetes Care* 2019;42:2322–2329
- Welsh C, Welsh P, Celis-Morales CA, et al. Glycated hemoglobin, prediabetes, and the links to cardiovascular disease: data from UK Biobank. *Diabetes Care* 2020;43:440–445
- Tang O, Matsushita K, Coresh J, et al. Mortality implications of prediabetes and diabetes in older adults. *Diabetes Care* 2020;43:382–388
- Petrie MC, Verma S, Docherty KF, et al. Effect of dapagliflozin on worsening heart failure and cardiovascular death in patients with heart failure with and without diabetes. *JAMA* 2020;323:1553–1568
- Mahendran DC, Hamilton G, Weiss J, et al. Prevalence of pre-existing dysglycaemia among inpatients with acute coronary syndrome and associations with outcomes. *Diabetes Res Clin Pract* 2019;154:130–137
- World Health Organization. *Definition, Diagnosis and Classification of Diabetes Mellitus and Its Complications: Report of a WHO Consultation. Part 1: Diagnosis and Classification of Diabetes Mellitus*. Geneva, World Health Organization, 1999
- World Health Organization, International Diabetes Federation. *Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycaemia: Report of a WHO/IDF Consultation*. Geneva, World Health Organization, 2006
- World Health Organization. Report of a World Health Organization Consultation. Use of glycated haemoglobin (HbA1c) in the diagnosis of diabetes mellitus. *Diabetes Res Clin Pract* 2011;93:299–309
- Punthakee Z, Goldenberg R, Katz P; Diabetes Canada Clinical Practice Guidelines Expert Committee. Definition, classification and diagnosis of diabetes, prediabetes and metabolic syndrome. *Can J Diabetes* 2018;42(Suppl. 1):S10–S15
- Tirosh A, Shai I, Tekes-Manova D, et al.; Israeli Diabetes Research Group. Normal fasting plasma glucose levels and type 2 diabetes in young men. *N Engl J Med* 2005;353:1454–1462
- Nichols GA, Hillier TA, Brown JB. Normal fasting plasma glucose and risk of type 2 diabetes diagnosis. *Am J Med* 2008;121:519–524
- Brambilla P, La Valle E, Falbo R, et al. Normal fasting plasma glucose and risk of type 2 diabetes. *Diabetes Care* 2011;34:1372–1374
- Schmidt MI, Bracco PA, Yudkin JS, et al. Intermediate hyperglycaemia to predict progression to type 2 diabetes (ELSA-Brasil): an occupational cohort study in Brazil. *Lancet Diabetes Endocrinol* 2019;7:267–277
- Ezquerro-Lázaro I, Cea-Soriano L, Giraldez-García C, et al.; PREDAPS Study Group. Lifestyle factors do not explain the difference on diabetes progression according to type of prediabetes: results from a Spanish prospective cohort of prediabetic patients. *Diabetes Res Clin Pract* 2019;153:66–75
- Diabetes Prevention Program Research Group. Effects of withdrawal from metformin on the development of diabetes in the diabetes prevention program. *Diabetes Care* 2003;26:977–980
- Boden G, Master RW, Gordon SS, Shuman CR, Owen OE. Monitoring metabolic control in diabetic outpatients with glycosylated hemoglobin. *Ann Intern Med* 1980;92:357–360
- DeFronzo RA, Goodman AM; The Multicenter Metformin Study Group. Efficacy of metformin in patients with non-insulin-dependent diabetes mellitus. *N Engl J Med* 1995;333:541–549
- Kahn SE, Haffner SM, Heise MA, et al.; ADOPT Study Group. Glycemic durability of rosiglitazone, metformin, or glyburide monotherapy. *N Engl J Med* 2006;355:2427–2443
- Knowler WC, Fowler SE, Hamman RF, et al.; Diabetes Prevention Program Research Group. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009;374:1677–1686
- Knowler WC, Hamman RF, Edelstein SL, et al.; Diabetes Prevention Program Research Group. Prevention of type 2 diabetes with troglitazone in the Diabetes Prevention Program. *Diabetes* 2005;54:1150–1156
- DREAM (Diabetes REduction Assessment with ramipril and rosiglitazone Medication) Trial Investigators; Gerstein HC, Yusuf S, Bosch J, et al. Effect of rosiglitazone on the frequency of diabetes in patients with impaired glucose tolerance or impaired fasting glucose: a randomised controlled trial. *Lancet* 2006;368:1096–1105
- DREAM Trial Investigators. Incidence of diabetes following ramipril or rosiglitazone withdrawal. *Diabetes Care* 2011;34:1265–1269
- DREAM On (Diabetes Reduction Assessment with Ramipril and Rosiglitazone Medication Ongoing Follow-up) Investigators. Long-term effect of rosiglitazone and/or ramipril on the incidence of diabetes. *Diabetologia* 2011;54:487–495
- Gerstein HC, Bosch J, Dagenais GR, et al.; ORIGIN Trial Investigators. Basal insulin and cardiovascular and other outcomes in dysglycemia. *N Engl J Med* 2012;367:319–328
- Retnakaran R, Qi Y, Harris SB, Hanley AJ, Zinman B. Changes over time in glycemic control, insulin sensitivity, and  $\beta$ -cell function in response to low-dose metformin and thiazolidinedione combination therapy in patients with impaired glucose tolerance. *Diabetes Care* 2011;34:1601–1604
- RISE Consortium. Lack of durable improvements in  $\beta$ -cell function following withdrawal of pharmacological interventions in adults with impaired glucose tolerance or recently diagnosed type 2 diabetes. *Diabetes Care* 2019;42:1742–1751
- Haw JS, Galaviz KI, Straus AN, et al. Long-term sustainability of diabetes prevention approaches: a systematic review and meta-analysis of randomized clinical trials. *JAMA Intern Med* 2017;177:1808–1817
- Perreault L, Pan Q, Mather KJ, Watson KE, Hamman RF, Kahn SE; Diabetes Prevention Program Research Group. Effect of regression from prediabetes to normal glucose regulation on long-term reduction in diabetes risk: results from the Diabetes Prevention Program Outcomes Study. *Lancet* 2012;379:2243–2251
- Echouffo-Tcheugui JB, Niiranen TJ, McCabe EL, et al. Lifetime prevalence and prognosis of

- prediabetes without progression to diabetes. *Diabetes Care* 2018;41:e117–e118
47. Shang Y, Marseglia A, Fratiglioni L, et al. Natural history of prediabetes in older adults from a population-based longitudinal study. *J Intern Med* 2019;286:326–340
48. Hwang Y-C, Cho I-J, Jeong I-K, Ahn KJ, Chung HY. Factors associated with regression from prediabetes to normal glucose tolerance in a Korean general population: a community-based 10-year prospective cohort study. *Diabet Med* 2018;35:1544–1551
49. Vistisen D, Kivimäki M, Perreault L, et al. Reversion from prediabetes to normoglycaemia and risk of cardiovascular disease and mortality: the Whitehall II cohort study. *Diabetologia* 2019;62:1385–1390
50. Richter B, Hemmingsen B, Metzendorf MI, Takwoingi Y. Development of type 2 diabetes mellitus in people with intermediate hyperglycaemia. *Cochrane Database Syst Rev* 2018;10:CD012661
51. Nathan DM, Genuth S, Lachin J, et al.; Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977–986
52. Ohkubo Y, Kishikawa H, Araki E, et al. Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus: a randomized prospective 6-year study. *Diabetes Res Clin Pract* 1995;28:103–117
53. Krolewski AS, Laffel LMB, Krolewski M, Quinn M, Warram JH. Glycosylated hemoglobin and the risk of microalbuminuria in patients with insulin-dependent diabetes mellitus. *N Engl J Med* 1995;332:1251–1255
54. Tanaka Y, Atsumi Y, Matsuoka K, Onuma T, Tohjima T, Kawamori R. Role of glycemic control and blood pressure in the development and progression of nephropathy in elderly Japanese NIDDM patients. *Diabetes Care* 1998;21:116–120
55. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837–853
56. American Diabetes Association. Standards of medical care in diabetes—2014. *Diabetes Care* 2014;37(Suppl. 1):S14–S80
57. Rosenzweig JL, Bakris GL, Berglund LF, et al. Primary prevention of ASCVD and T2DM in patients at metabolic risk: an Endocrine Society\* Clinical Practice Guideline. *J Clin Endocrinol Metab* 2019;104:3939–3985
58. Moin T, Li J, Duru OK, et al. Metformin prescription for insured adults with prediabetes from 2010 to 2012: a retrospective cohort study. *Ann Intern Med* 2015;162:542–548
59. Tseng E, Yeh HC, Maruthur NM. Metformin use in prediabetes among U.S. adults, 2005–2012. *Diabetes Care* 2017;40:887–893