


Cell Host & Microbe

An early-life microbiota metabolite protects against obesity by regulating intestinal lipid metabolism

Graphical abstract



Authors

Catherine D. Shelton, Elizabeth Sing, Jessica Mo, ..., Nicholas O. Markham, Min Liu, Mariana X. Byndloss

Correspondence

mariana.x.byndloss@vumc.org

In brief

Shelton et al. determine that early-life antibiotics exacerbate diet-induced obesity by disrupting interactions between the gut microbiota and the small intestine epithelium. Their study identifies that a *Lactobacillus*-derived metabolite, phenylactic acid, regulates intestinal PPAR- γ to limit fat accumulation, revealing a mechanism by which the early-life microbiota protects against metabolic dysfunction.

Highlights

- Early-life exposure to antibiotics and a HF diet exacerbates obesity
- Loss of small intestinal Lactobacillaceae leads to increased adiposity
- Antibiotics and a HF diet exacerbate adiposity via depletion of intestinal PPAR- γ
- *Lactobacillus*-derived phenylactic acid protects against antibiotic-induced obesity



Article

An early-life microbiota metabolite protects against obesity by regulating intestinal lipid metabolism

Catherine D. Shelton,¹ Elizabeth Sing,¹ Jessica Mo,¹ Nicolas G. Shealy,¹ Woongjae Yoo,^{1,2} Julia Thomas,¹ Gillian N. Fitz,³ Pollyana R. Castro,^{1,4} Tara T. Hickman,⁵ Teresa P. Torres,¹ Nora J. Foegeding,¹ Jacob K. Zieba,¹ M. Wade Calcutt,⁶ Simona G. Codreanu,⁷ Stacy D. Sherrod,⁷ John A. McLean,⁷ Sun H. Peck,^{5,8,9,10} Fan Yang,^{3,11} Nicholas O. Markham,^{9,10,12,13} Min Liu,¹⁴ and Mariana X. Byndloss^{1,13,15,16,17,18,*}

¹Department of Pathology, Microbiology, and Immunology, Vanderbilt University Medical Center, Nashville, TN 37232, USA

²Department of Life Sciences, Pohang University of Science and Technology (POSTECH), Pohang 37673, Republic of Korea

³Department of Cell and Developmental Biology, Vanderbilt University School of Medicine, Nashville, TN 37232, USA

⁴Laboratory of Immunoinflammation, Department of Genetics, Evolution, Microbiology, and Immunology, Institute of Biology, University of Campinas, Campinas, São Paulo 12083-862, Brazil

⁵Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, TN 37232, USA

⁶Mass Spectrometry Research Center and Department of Biochemistry, Vanderbilt University School of Medicine, Nashville, TN 37232, USA

⁷Center for Innovative Technology and Department of Chemistry, Vanderbilt University, Nashville, TN 37232, USA

⁸Department of Biomedical Engineering, Vanderbilt University School of Engineering, Nashville, TN 37232, USA

⁹Department of Veterans Affairs, Tennessee Valley Healthcare System, Nashville, TN, USA

¹⁰Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

¹¹Department of Pediatrics, Vanderbilt University Medical Center, Nashville, TN 37232, USA

¹²Epithelial Biology Center, Vanderbilt University Medical Center, Nashville, TN, USA

¹³Vanderbilt Institute of Infection, Immunology, and Inflammation, Vanderbilt University Medical Center, Nashville, TN 37232, USA

¹⁴Department of Pathology and Molecular Medicine, Metabolic Diseases Institute, University of Cincinnati College of Medicine, Cincinnati, OH 45237, USA

¹⁵Vanderbilt Digestive Disease Center, Vanderbilt University Medical Center, Nashville, TN 37232, USA

¹⁶Vanderbilt Microbiome Innovation Center, Vanderbilt University, Nashville, TN 37235, USA

¹⁷Howard Hughes Medical Institute, Vanderbilt University Medical Center, Nashville, TN 37232, USA

¹⁸Lead contact

*Correspondence: mariana.x.byndloss@vumc.org

<https://doi.org/10.1016/j.chom.2023.09.002>

SUMMARY

The mechanisms by which the early-life microbiota protects against environmental factors that promote childhood obesity remain largely unknown. Using a mouse model in which young mice are simultaneously exposed to antibiotics and a high-fat (HF) diet, we show that *Lactobacillus* species, predominant members of the small intestine (SI) microbiota, regulate intestinal epithelial cells (IECs) to limit diet-induced obesity during early life. A *Lactobacillus*-derived metabolite, phenyllactic acid (PLA), protects against metabolic dysfunction caused by early-life exposure to antibiotics and a HF diet by increasing the abundance of peroxisome proliferator-activated receptor γ (PPAR- γ) in SI IECs. Therefore, PLA is a microbiota-derived metabolite that activates protective pathways in the small intestinal epithelium to regulate intestinal lipid metabolism and prevent antibiotic-associated obesity during early life.

INTRODUCTION

The early-life gut microbiota plays a crucial role in promoting health.^{1,2} In contrast to an adult microbiota, the infant microbiota is characterized by lower bacterial abundance and diversity.^{3–5}

As an infant grows, their microbiota becomes more complex and resembles that of an adult by ages 3–5 years old.⁴ During this developmental window, the early-life microbiota is uniquely sensitive to perturbations by environmental factors, such as an obesogenic high-fat (HF) diet or antibiotic treatment.^{6,7} Disruption to the early-life gut microbiota is hypothesized to have last-

ing impacts on a child's health. For instance, early-life antibiotic exposure is correlated with increased weight gain and body mass index (BMI) in children.^{8,9} Previous research using mouse models corroborated human studies and established a causal link between changes in the gut microbiota composition due to early-life antibiotic exposure and increased adiposity.^{10,11} However, the mechanisms by which the gut microbiota exerts its protective effects in the context of childhood obesity are largely unknown.

An unappreciated aspect of the link between early-life antibiotic use and the risk for obesity is the impact of dietary habits



on the microbiota. Diet is a key determinant of microbiota composition. For instance, an obesogenic Western-style HF diet perturbs the microbiota and elevates the risk for metabolic diseases.^{12,13} In the United States, children have an elevated intake of saturated fats¹⁴ and a high percentage of therapeutic antibiotic usage.¹⁵ However, how concurrent exposure to an HF diet and antibiotics alters the risk for childhood obesity remains unknown. Thus, a complete understanding of how early-life antibiotic treatment increases adiposity requires an investigation into the compounding effects of diet and antibiotics on the gut microbiota and host metabolism.

Both the gut microbiota and diet regulate the function of the intestinal epithelium. During consumption of a Western-style obesogenic HF diet, the small intestine (SI) epithelium adapts to contend with the increase in dietary fat.¹⁶ Specifically, the SI epithelium increases its ability to metabolize fats, protecting the host from features of metabolic dysfunction.^{16–18} Concurrently, the SI microbiota changes upon exposure to an HF diet, which impacts the function of the SI,¹⁹ as the SI microbiota serves as an important regulator of intestinal lipid absorption, metabolism, and secretion.^{19–21} Furthermore, recent work has determined that specific members of the SI microbiota can inhibit lipid secretion and thereby limit serum triglycerides during consumption of an HF diet.^{22,23} Therefore, early-life antibiotics may promote obesity through perturbation to microbiota-epithelium interactions that protect against metabolic dysfunction caused by an HF diet.

In this manuscript, we investigate how early-life microbiota disruption promotes HF diet-induced obesity. We simultaneously exposed young mice to antibiotics and an HF diet and determined that antibiotic treatment leads to greater adiposity and metabolic dysfunction during consumption of an HF diet. Concurrent exposure to antibiotics and an HF diet depleted *Lactobacillus* species from the SI microbiota and experiments in colonized gnotobiotic identified the loss of *Lactobacillus* species as the driver of excess fat accumulation in our model. Characterization of the intestinal epithelium in mice given antibiotics and an HF diet revealed dysregulation of genes related to lipid metabolism, leading to increased triglycerides in the SI epithelium and serum. We propose that disruption to intestinal lipid metabolism leads to increased adiposity through depletion of peroxisome proliferator-activated receptor γ (PPAR- γ), a regulator of lipid metabolism, in the SI epithelium. Untargeted metabolomics revealed that antibiotics and an HF diet reduced the abundance of the *Lactobacillus*-derived metabolite, phenyllactic acid (PLA), in the SI lumen. We show that PLA upregulated intestinal PPAR- γ and protected against metabolic dysfunction caused by early-life exposure to antibiotics and an HF diet. Therefore, PLA is a microbiota-derived metabolite that activates protective pathways in the SI epithelium to prevent obesity during early life.

RESULTS

Concurrent exposure to antibiotics and an HF diet during early life promotes long-lasting metabolic dysfunction

To investigate whether early-life concurrent exposure to detrimental environmental factors (e.g., obesogenic HF diet and an-

tibiotics) have a synergistic effect in promoting obesity, we first established a mouse model (Figure 1A) in which young (3-week-old) mice were exposed to either a low-fat (LF) (10%) or HF (60%) diet with or without low-dose penicillin (LDP) treatment (6.67 mg/L).¹⁰ Mice were maintained on their respective diets and antibiotics for 5 weeks until they reached sexual maturity. After 5 weeks, mice exposed to an HF diet and LDP gained significantly more weight (Figure 1B) and abdominal fat (Figure 1C) than mice fed an HF diet alone, despite equal food consumption (Figure S1A). Furthermore, no difference in weight gain (Figure 1B) or abdominal fat (Figure 1C) was observed between mice fed an LF diet and mice given an LF diet and LDP, indicating that antibiotic exposure alone was insufficient to promote additional adiposity in our model. To assess whether the increased weight gain in mice exposed to an HF diet and LDP was due to overall body composition changes (e.g., lean mass or bone mineral density), we performed dual-energy X-ray absorptiometry (DEXA) scanning in animals exposed to an HF diet or HF diet and LDP (Figures S1B and S1C). DEXA analysis revealed no differences in lean weight (Figure S1B) or bone mineral density (Figure S1C) between experimental groups, suggesting that an HF diet and LDP specifically increase fat accumulation. Mice treated with LDP and given an HF diet also developed additional features of metabolic dysfunction, including increased fasting blood glucose (Figure 1D) and greater fat accumulation in hepatocytes (Figures 1E and 1F) compared with HF diet-fed mice.

To determine if the metabolic dysfunction persisted after antibiotic treatment ended, mice exposed to either an HF diet and LDP or an HF diet alone were given only an HF diet for an additional 5 weeks (Figure 1G). HF diet and LDP-exposed mice gained more weight (Figure 1H) and abdominal fat (Figure 1I) than mice only given an HF diet for 10 weeks, despite no longer receiving LDP and consuming equal amounts of food throughout the experiment (Figure S2A). Mice treated with LDP and fed an HF diet also sustained elevated fasting blood glucose 3 weeks after antibiotics were removed (Figure 1J). Differences between experimental groups were not significant at 10 weeks, when some mice exposed to an HF diet alone started to show increased fasting glucose levels.²⁴ These data suggest that early-life exposure to an HF diet and LDP causes increased fasting blood glucose (Figure 1D) and that glucose levels remain elevated in this experimental group even weeks after antibiotics are removed (Figure 1J). Consumption of an HF diet was required for LDP to promote lasting adiposity, as mice switched onto an LF diet after 5 weeks of exposure to an HF diet and LDP no longer gained additional weight or adiposity (Figures S2B–S2D). Moreover, mice treated with LDP and fed an LF diet were also given an LF diet alone for 5 additional weeks, and no differences in weight gain or abdominal fat were observed (Figures S2E–S2G). Collectively, these data reveal that concurrent early-life exposure to antibiotics and an HF diet leads to early-onset and lasting HF-dependent obesity and metabolic dysfunction through unknown mechanisms.

Depletion of *Ligilactobacillus murinus* from the SI microbiota increases fat accumulation

As disruption of the early-life microbiota by antibiotics was previously shown to promote adiposity,¹¹ we hypothesized that simultaneous exposure to antibiotics and an HF diet altered




Figure 1. Early-life exposure to low-dose penicillin (LDP) and a high-fat (HF) diet promotes metabolic dysfunction

(A and G) Schematics of the experimental models and the groups used.

(B–D) (B) Weight gain, (C) abdominal fat (g), and (D) fasting glucose levels of mice in each group after 5 weeks.

(E) Representative images of oil red O-stained sections of liver from HF diet and HF diet + LDP-treated mice. Scale bar represents 100 μ m.

(F) Combined oil red scores from livers of mice given an HF diet or an HF diet + LDP.

(H) Weight gain of mice in each group over the course of the 10-week experiment.

(I) Abdominal fat (g) measured after the 10-week diet and antibiotic manipulations.

(J) Fasting glucose levels were measured after 8 and 10 weeks on an HF diet.

(B–D, F, I, and J) Each dot represents one animal. Bars represent geometric mean. (H) Dots represent mean \pm standard error of the mean. (B and C) $n = 12$ mice/group. Data representative of two independent cohorts. (D, F, and H–J) $n = 9$ mice/group. Data representative of one independent cohort. (B–D, I, and J) * $p < 0.05$; ** $p < 0.01$; *** $p < 0.005$ using an unpaired two-tailed Student's t test. (F) * $p < 0.05$ using Mann-Whitney test. (H) * $p < 0.05$; ** $p < 0.01$ using multiple unpaired two-tailed Student's t tests.

the gut microbiota to accelerate fat deposition. Antibiotic treatment increased metabolic dysfunction after a 5-week treatment only in young mice fed an obesogenic HF diet (Figures 1B–1H), pointing to a contribution of dietary lipids to our phenotypes. As the SI is the primary site for dietary fat absorption and metabolism,¹⁹ we decided to investigate changes in the SI microbiota during concurrent early-life exposure to an HF diet and antibiotics. 16S rRNA analysis of the distal SI microbiota revealed that diet was the most predominant driver of compositional changes between the four experimental groups (Figures S3A–S3C), consistent with previously published research.²⁵ As antibiotics promoted adiposity only when young mice consumed an

HF diet (Figure 1C), we focused on differences between the SI microbiota of mice given an HF diet or an HF diet and LDP and not mice that were fed an LF diet or an LF diet and LDP. Relative to the HF diet-only group, exposure to LDP led to significant depletion of bacteria belonging to the Lactobacillaceae family, the most abundant family in their SI microbiota (Figures 2A, 2B, S3C, and S3D). Interestingly, *in vitro* growth of *Lactobacillus* species indicated sensitivity to the combination of penicillin and lipids or the combination of penicillin and bile salts (elevated in the intestine during consumption of an HF diet²⁶) (Figures S3F and S3G), providing an explanation for the depletion of *Lactobacillus* spp. during exposure to penicillin and an HF diet. While




Figure 2. *Ligilactobacillus murinus* protects against adiposity during early-life consumption of an HF diet

(A) Heatmap showing differential abundance of the 5 most abundant families in the small intestine microbiota of mice after 5 weeks of treatment. (B) Relative abundance of the genus *Lactobacillus* in the small intestine microbiota as determined by 16S rRNA sequencing. (C) The abundance of *L. murinus* in the small intestine as determined by qPCR. (D and E) (D) Weight gain and (E) abdominal fat of mice given an HF diet and LDP alone or gavaged with a penicillin resistant strain of *L. murinus* (*L. murinus* Pen^R) after 5 weeks. (F) A schematic representation of the gnotobiotic experiment. (G) *Lactobacillus* abundance (colony-forming unit [CFU]/gram) in small intestine (SI) after 5 weeks on an HF. Dotted line indicates limit of detection for *Lactobacillus* species in this experiment. (H and I) (H) Weight gain and (I) abdominal fat (g) of mice after 5 weeks. Each dot represents one animal. Bars represent geometric mean. (A) n = 5 mice/group; (B) n = 9 mice/group; (C) n = 7–8 mice/group; (D and E) n = 12 mice/group; (G and H) n = 8 mice/group. Data representative of two independent cohorts. *p < 0.05; **p < 0.01; ***p < 0.005; ****p < 0.001 using an unpaired two-tailed Student's t test.

Lactobacillus levels recovered in HF diet and LDP-exposed mice when they were switched onto an HF diet alone, their weight gain still persisted (Figures 1H, S3H, and S3I), supporting the hypothesis that short-term microbiota perturbations can promote lasting metabolic changes.¹¹

We next investigated if the loss of *Lactobacillus* species promoted metabolic dysfunction in mice exposed to antibiotics and an HF diet. Quantitative PCR (qPCR) revealed that the predominant *Lactobacillus* species present in the SI of untreated

mice was *Ligilactobacillus murinus* (*L. murinus*) (Figure S3E). Moreover, we determined that *L. murinus* was significantly depleted from the SI microbiota of mice given LDP and an HF diet (Figure 2C). Therefore, we generated a strain of *L. murinus* resistant to penicillin (*L. murinus* Pen^R) and used this strain to restore the presence of *L. murinus* in the intestinal lumen by re-colonizing a subset of mice exposed to an HF diet and LDP with *L. murinus* Pen^R. Mice given *L. murinus* Pen^R gained less weight (Figure 2D) and abdominal fat (Figure 2E) than mice




Figure 3. Repeated treatment with clinical doses of penicillin (CDP) and a high-fat diet promote increased adiposity

(A) A schematic of the experimental model and the groups used.

(B–D) (B) Weight gain, (C) percent abdominal fat, and (D) fasting glucose levels of mice after 5 weeks.

(E) *Lactobacillus* abundance (colony-forming unit [CFU]/gram) in the feces of treated mice over the course of the 5-week experiment.

(F) *Lactobacillus* abundance in the small intestine was determined after 5 weeks.

(A–D and F) Each dot represents one animal. Bars represent geometric mean. $n = 12$ mice/group. (E) Dots indicate geometric mean of abundances from six mice. * $p < 0.05$; ** $p < 0.01$; **** $p < 0.001$ using an unpaired two-tailed Student's *t* test.

exposed to an HF diet and LDP alone. Due to the transient colonization seen with *L. murinus* Pen^R (data not shown), we performed an additional experiment using a defined microbiota strategy in gnotobiotic mice. Young (3-week-old) germ-free (GF) mice were colonized with either a defined microbiota containing *L. murinus* or lacking *L. murinus* (Figure 2F) and were then maintained on an HF diet for 5 weeks. Mice colonized with a microbiota containing *L. murinus* (Figure 2G) gained less weight (Figure 2H) and accumulated less abdominal fat (Figure 2I) than mice given a microbiota without *L. murinus*. Together, these data suggest that *L. murinus* protects young mice from adiposity during the consumption of an HF diet.

Our model utilizes chronic, low doses of antibiotics,²⁷ an exposure that largely results from dietary sources.²⁸ However, children can also be treated with therapeutic doses of antibiotics during infections, which our LDP treatment does not replicate. To address if therapeutically relevant concentrations of antibiotics combined with an HF diet led to similar physiological effects as LDP, we exposed young mice to three clinical doses of penicillin (CDP, 0.167 g/L) (in combination with an LF or HF diet) over 5 weeks (Figure 3A). Consistent with LDP, concurrent exposure to an HF diet and clinical doses of antibiotics caused marked weight gain (Figure 3B), as well as increased abdominal fat (Figure 3C), and fasting blood glucose (Figure 3D) in mice

compared with mice given only an HF diet. As seen in our LDP model, CDP depleted *Lactobacillus* from the microbiota of young mice (Figures 3E and 3F), supporting the role of *Lactobacillus* depletion from the SI microbiota as a driver of the synergistic effect of clinical and subclinical antibiotic exposure and HF diet in promoting obesity. To further investigate how *L. murinus* exerts its anti-obesity effects, we chose to utilize our HF diet and LDP model, as mice simultaneously exposed to an HF diet and LDP had sustained *Lactobacillus* depletion from their small intestinal microbiota (Figures 1I, 1J, and S3H).

Early-life antibiotics disrupt intestinal lipid metabolism during the consumption of an HF diet

A mechanism by which disruptions to the gut microbiota can lead to metabolic dysfunction is through perturbation of intestinal epithelium function.^{29,30} The intestinal epithelium contributes to a healthy host metabolism through numerous mechanisms, including regulation of lipid metabolism, uptake, and secretion as well as gut hormone production.³¹ Therefore, changes in the early-life gut microbiota due to antibiotic exposure and an HF diet may contribute to childhood obesity by disrupting the activity of intestinal epithelial cells (IECs). To determine if the combination of an HF diet and antibiotics affected the intestinal epithelium, we examined sections of the ileum after 5 weeks of




Figure 4. Exposure to early-life antibiotics alters the response of the intestinal epithelium to a high-fat diet

(A) Hematoxylin and eosin-stained sections of the ileum from mice after 5 weeks of exposure to an LF or HF diet ± LDP. (B) Histopathology scores of ileum sections. Scale bar represents 200 μ m. (C and D) RNA sequencing analysis of epithelial cells isolated from the ileum (IECs) of mice after 5 weeks of exposure to an HF diet or an HF diet + LDP. (C) Volcano plot of genes significantly (an adjusted p value < 0.05) upregulated (red) and downregulated (blue) in mice exposed to an HF diet + LDP compared with mice given only an HF diet. (D) Enriched gene ontology (GO) categories in IECs of mice fed an HF diet + LDP treatment. (E) Epithelial transcripts of the indicated genes measured by qPCR in mice given an HF diet or an HF diet and LDP. (F) Fecal fat absorption was measured after exposure to an HF diet or HF diet and LDP for 5 weeks. (G–I) Mice exposed to an HF diet or an HF diet and LDP for 5 weeks were fasted and (G) triglyceride abundance in IECs and (H) neutral lipids in the serum were measured. (I) An oral fat tolerance test was performed in mice given an HF diet or an HF diet and LDP for 5 weeks. Serum triglyceride concentration was measured 2 h after the bolus of olive oil. (J and K) (J) Abdominal fat (g) and (K) fasting triglycerides of mice given an HF diet or an HF diet and LDP for 2 weeks. (B and E) Bars represent mean \pm standard error of the mean (SEM). (F–K) Bars represent geometric mean. Each dot represents one animal. (B) n = 6 mice/group. Data representative of one independent cohort; (C and D) n = 3 mice/group. (E) n = 14 mice/group. Data representative of two independent cohorts. (F–H) n = 5–6 mice/group. Data representative of one independent cohort. (I) n = 12 mice/group. Data representative of one independent cohort. (J and K) n = 9 mice/group. Data representative of one independent cohort. (B) *p < 0.05; **p < 0.01 using a one-tailed Mann-Whitney test. (E–K) *p < 0.05; **p < 0.01; ***p < 0.005; ****p < 0.0001 using an unpaired two-tailed Student's t test.

exposure to either an LF or HF diet with or without concurrent treatment with LDP (Figure 4A). Histopathological analysis of sections revealed that the individual impact of either LDP treatment or an HF diet synergized in perturbing the intestinal epithelium in mice exposed to both LDP and an HF diet (Figure 4B). To gain a deeper understanding of how exposure to an HF diet and LDP altered the function of the intestinal epithelium, we next performed RNA sequencing of distal SI IECs in mice given an HF diet

or an HF diet and LDP. We determined that concurrent exposure to an obesogenic HF diet and antibiotics altered the expression of approximately 200 genes (compared with mice fed an HF diet alone) belonging to diverse pathways (Figures 4C and 4D). Analysis of gene pathways altered in HF diet and LDP-treated mice revealed perturbed expression of transcripts related to lipid and fatty acid metabolism (Figure 4D). qPCR analysis of SI IECs confirmed RNA sequencing results and showed a




Figure 5. Loss of intestinal PPAR- γ signaling promotes increased adiposity due to exposure to LDP and an HF diet

(A) Expression of *Pparg* in the ileum epithelium of HF diet or HF diet + LDP-treated mice measured by qPCR.
 (B) PPAR- γ abundance in the epithelium was quantified by scoring stained sections of the ileum.
 (C and D) Representative images of PPAR- γ staining in (C) HF diet-fed mice and (D) HF diet-fed + LDP-treated mice. Scale bar represents 200 μ m.
 (E) 3-week-old *Pparg*^{fl/fl}*Villin*^{cre/+} mice (*Pparg* ^{Δ IEC}) and littermate control *Pparg*^{fl/fl}*Villin*^{-/-} mice (*Pparg*^{fl/fl}) were fed an HF diet for 10 weeks and weight gain was determined weekly.
 (F and G) (F) Abdominal fat (% total body weight) and (G) *Lactobacillus* abundance (colony-forming unit [CFU]/gram) in the small intestine (SI) of *Pparg* ^{Δ IEC} and *Pparg*^{fl/fl}, measured after the 6-week diet and antibiotic manipulations.
 (H–L) Intestinal epithelial cells were infected with *L. murinus* at an MOI of 100. (H) Amount of BODIPY C₁₂ secreted by treated Caco-2 cells was measured 4 h after addition of lipid micelles. (I) Immortalized mouse small intestinal epithelial (MSIE) cells were mock-treated or treated with *E. coli* Mt1B1, *S. xyloso* 33-ERD13C, or *L. murinus*, before expression of *Pparg* was measured using qPCR. (J–L) MSIE cells were treated with *L. murinus* (L) or left untreated (K) and incubated for 16 h. Cells were subsequently stained with anti-PPAR- γ and phalloidin to stain for F-actin. (J) PPAR- γ nuclear intensity was quantified using ImageJ; 15 cells were selected from two images from three independent experiments.
 (M–O) Germ-free mice were colonized with a defined microbiota with or without *L. murinus* and placed on an HF diet for 5 weeks. (M) PPAR- γ abundance in the epithelium was quantified by scoring stained sections of the ileum. Representative images of PPAR- γ staining in mice colonized with (O) or without (N) *L. murinus*. Scale bar represents 200 μ m.

(legend continued on next page)

significant difference in the expression of genes related to lipid digestion (*Angptl4*) and absorption (*Cd36*), trafficking (*Fabp1* and *Fabp2*), and secretion (*Mttp* and *ApoB*) (Figure 4E).^{32–37} Despite increased *Cd36* expression, mice that were given an HF diet and LDP did not absorb more fat from the diet compared with mice that were only fed an HF diet (Figure 4F). However, we measured significantly higher triglycerides in the SI IECs of fasted HF diet and LDP-treated mice (Figure 4G). We also determined that mice exposed to an HF diet and LDP had significantly more lipids in their fasted serum than mice given only an HF diet (Figure 4H). Thus, we reasoned that early-life treatment with LDP perturbed intestinal lipid metabolism, leading to increased lipid transport from the intestine into circulation. To investigate this hypothesis, we injected mice given an HF diet or HF diet and LDP with the lipoprotein lipase inhibitor tyloxapol and then challenged the mice with a bolus of olive oil. Mice exposed to an HF diet and LDP had significantly more serum triglycerides 2 h after the olive oil gavage (Figure 4I), suggesting that their elevated serum lipids (Figure 4H) result from greater intestinal lipid secretion.

To determine if the elevated serum lipids in mice exposed to an HF diet and LDP preceded their increased adiposity, we investigated differences in physiology after only 2 weeks on an HF diet and LDP. Despite a reduction in *Lactobacillus* species in the SI (Figure S4A), mice given an HF diet and LDP did not gain more weight (Figure S4B) or accumulate more abdominal fat (Figure 4J) after only 2 weeks of exposure. However, HF diet and LDP-treated mice had significantly greater triglycerides in their fasted serum (Figure 4K). To understand if elevated triglycerides in the serum coincided with perturbation of intestinal lipid metabolism after 2 weeks of exposure to an HF diet and LDP, we measured the expression of the same genes that were altered after 5 weeks in mice given an HF diet and LDP (Figure 4E). Interestingly, expression of *Angptl4* was decreased after exposure to an HF diet and LDP for 2 weeks, but no significant differences were observed in the expression of the other lipid metabolism genes (Figure S4F). Our data suggest that exposure to an HF diet and LDP dysregulates intestinal lipid metabolism, leading to elevated serum triglycerides, promoting excess weight gain and adiposity.

Activation of intestinal PPAR- γ by *L. murinus* protects against fat accumulation during consumption of an HF diet

We next investigated how an HF diet and LDP disrupted intestinal lipid metabolism, specifically focusing on mechanisms leading to reduced *Angptl4* in SI IECs, as this preceded the development of metabolic dysfunction. In the intestine, PPAR- γ , a nuclear receptor, regulates *Angptl4* expression^{38,39} as well as lipid metabolism in the SI.⁴⁰ Importantly, PPAR- γ abundance and activity in colonic IECs decreases during antibiotic treatment.³⁸ Therefore, we investigated whether intestinal PPAR- γ

was a mechanism connecting microbiota perturbations to dysregulated intestinal lipid metabolism during early-life exposure to an HF diet and LDP. LDP treatment did not impact the gene expression and protein abundance of PPAR- γ in SI IECs when mice were fed an LF diet (Figures S5A–S5D). However, PPAR- γ gene expression and protein abundance in SI IECs was significantly reduced in mice given an HF diet and LDP (relative to mice fed an HF diet alone) (Figures 5A–5D). No changes in the expression of *Pparg* or related target genes were observed in the adipose tissue or liver (Figures S5E and S5F). Our findings suggest that exposure to an HF diet and antibiotics specifically depleted PPAR- γ from SI IECs.

With the reduction of intestinal PPAR- γ observed in HF diet and LDP-treated mice, we next asked if genetic ablation of PPAR- γ specifically in IECs resulted in increased weight gain and metabolic dysfunction. Intestinal epithelial-specific PPAR- γ knockout mice (*Pparg* ^{Δ IEC}) and littermate control mice (*Pparg*^{fl/fl}) were fed an LF or HF diet for 10 weeks (Figures 5E and S5G). *Pparg* ^{Δ IEC} gained more weight and fat than their littermate controls when fed an HF diet (Figures 5E and S5H). Consistent with the effects of an HF diet and LDP, *Pparg* ^{Δ IEC} did not gain more weight or fat when given an LF diet (Figures S5G and S5H). To determine if the obesogenic effects of an HF diet and LDP act through depletion of intestinal PPAR- γ , we exposed *Pparg* ^{Δ IEC} and wild-type *Pparg*^{fl/fl} to an HF diet or an HF diet and LDP. Wild-type *Pparg*^{fl/fl} mice developed significantly more abdominal fat when exposed to an HF diet and LDP compared with when only fed an HF diet (Figure 5F). However, an HF diet and LDP treatment failed to increase the amount of adiposity accumulated by *Pparg* ^{Δ IEC} mice (Figure 5F), supporting our hypothesis that LDP promotes fat accumulation during exposure to an HF diet by blunting PPAR- γ signaling in the intestinal epithelium. To confirm that differences in adiposity were not attributable to changes in LDP activity, *Lactobacillus* abundance in the SI of *Pparg* ^{Δ IEC} and *Pparg*^{fl/fl} mice was measured. No difference in *Lactobacillus* abundance was observed between *Pparg* ^{Δ IEC} and *Pparg*^{fl/fl} mice fed an HF diet or between *Pparg* ^{Δ IEC} and *Pparg*^{fl/fl} mice exposed to an HF diet and LDP (Figure 5G).

Lactobacillus species can modulate lipid secretion in IECs through activation of the PPARs.^{22,23} As we observed a reduction in *Lactobacillus* spp. upon exposure to an HF diet and LDP (Figure 2A), we next investigated whether *Lactobacillus* spp. could activate PPAR- γ *in vitro*. As we observed that *L. murinus* protected young mice from HF diet-induced obesity (Figure 2I), we utilized this isolate to interrogate the relationship between *Lactobacillus* spp. and PPAR- γ . In intestinal epithelial (Caco-2) cells, we determined that *L. murinus* increased the transcriptional activity of PPAR- γ to a similar level as the known PPAR- γ agonist, butyrate³⁸ (Figure S6A). Furthermore, we observed that *L. murinus* significantly increased the nuclear localization of PPAR- γ (Figures S6B, S6C, and S6E). As *L. murinus* regulated both the abundance and activity of

(A, F, and G) Each dot represents one animal. Bars represent geometric mean. n = 5–7 mice/group. (B and M) Each dot represents one animal. Bars represent mean. n = 6–7 mice/group. Data representative from one independent cohort. (E) Dots represent mean \pm SEM (N = 8–9/ genotype). (H and I) Each dot represents one technical replicate (data shown from three biological replicates). Bars represent geometric mean. (J) Violin plot showing the distribution of PPAR- γ nuclear intensity (n = 90 cells). (A and G–J) *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001 using an unpaired two-tailed Student's t test. (B and M) *p < 0.05 using a two-tailed Mann-Whitney test. (E) *p < 0.0332 using a two-way ANOVA with Sidák's multiple comparisons test. (F) *p < 0.05 using a one-way ANOVA with Sidák's multiple comparisons test.

PPAR- γ , we then asked whether *L. murinus* affected lipid secretion in a PPAR- γ -dependent manner. In agreement with previous findings,²³ *L. murinus* decreased secretion of micelles containing a fluorescent fatty acid (BODIPY C₁₂) by Caco-2 cells (Figure 5H). However, adding a PPAR- γ inhibitor, GW9662, blocked the ability of *L. murinus* to inhibit lipid secretion (Figure 5H), suggesting that *L. murinus* increases PPAR- γ activity in IECs to reduce lipid secretion from the intestine.

To continue our investigation into the relationship between *L. murinus* and PPAR- γ , we transitioned to an SI cell line to recapitulate our mouse model more accurately. *Pparg* expression was increased in mouse small intestinal epithelial (MSIE) cells treated with *L. murinus* (Figure 5I). In contrast, treatment with *Escherichia coli* (*E. coli*) Mt1B1 or *Staphylococcus xylosum* failed to increase expression of *Pparg* in MSIE cells, suggesting that the increase in *Pparg* expression is unique to *L. murinus* and not a result of co-culture with bacteria (Figure 5I). Moreover, compared with mock-treated cells, *L. murinus* significantly increased the amount of PPAR- γ in the nucleus in MSIE cells (Figures 5J–5L) (consistent with our findings in Caco-2 cells; Figure S6). Further evidence of a specific interaction with *L. murinus* and intestinal PPAR- γ is seen in gnotobiotic mice colonized with a defined microbiota with or without *L. murinus* (Figure 2F). In agreement with our *in vitro* data, mice colonized with a microbiota containing *L. murinus* have significantly higher levels of PPAR- γ staining in their intestine compared with mice lacking *L. murinus* (Figures 5M–5O).

As PPAR- γ is known to be regulated by microbiota-derived metabolites,^{38,41} we next investigated if the cell-free supernatant from *L. murinus* cultures could increase PPAR- γ nuclear localization. Indeed, the supernatant from *L. murinus* cultures, but not *E. coli* Mt1B1, significantly increased the amount of PPAR- γ accumulating in the nucleus (Figures S6B–S6I). Together, our *in vitro* data reveal that *Lactobacillus*-derived metabolites activate PPAR- γ , leading us to hypothesize that the loss of PPAR- γ activity due to exposure to an HF diet and antibiotics occurred because of the depletion of *Lactobacillus*-derived metabolites.

PLA is depleted from the SI lumen and regulates intestinal PPAR- γ

To identify *Lactobacillus* metabolites responsible for activating intestinal PPAR- γ , untargeted metabolomics was performed on the SI contents from mice given an HF diet or an HF diet and LDP (Figure 6A). Multiple metabolites known to be produced by *Lactobacillus* species were significantly depleted in mice exposed to an HF diet and LDP, including PLA (Figure 6B). As previous work identified that *Lactobacillus*-derived PLA upregulates *PPARG* in adipocytes,⁴² we investigated whether PLA was the *L. murinus*-derived metabolite in supernatants that regulated PPAR- γ in IECs. First, we determined that intestinal PLA abundance is dependent on the gut microbiota as GF mice have significantly less PLA in their SI compared with conventional mice when fed an HF diet (Figure 6C). PLA was then measured in supernatants from cultures of the defined microbiota used to colonize GF mice (Figure 2F). PLA levels were significantly greater in supernatants of the defined microbiota containing *L. murinus* (Figure 6D), indicating that *L. murinus* is the primary producer of PLA compared with other microbes present in

our defined SI microbiota. Furthermore, we determined that *L. murinus*, alone, produced PLA in cell culture experiments (Figure 6E). Importantly, *L. murinus* Pen^R (Figures 2D and 2E), another strain of *L. murinus* (DSM 20452), different *Lactobacillus* spp., and *Bifidobacterium animalis* also all produce PLA *in vitro* (Figure 6E), highlighting that the ability to produce PLA is conserved between multiple bacterial species present in the gut microbiota. To investigate PLA production by *L. murinus* *in vivo*, we measured intestinal PLA levels in mice given *L. murinus* Pen^R during exposure to an HF diet and LDP (Figures 2D and 2E). Compared with mice treated with an HF diet and LDP alone, mice colonized with *L. murinus* Pen^R had significantly higher PLA levels in their SI content (Figure 6F), supporting our hypothesis that the *Lactobacillus* species is a significant contributor to the intestinal PLA abundance.

To determine if PLA activated PPAR- γ in IECs, we treated mouse enteroids with PLA and observed increased expression of *Angptl4* in mouse enteroids to a similar level as the PPAR- γ agonist, Rosiglitazone (Figure 6G). Furthermore, PLA treatment significantly increased PPAR- γ accumulation in the nucleus of IECs compared with mock-treated cells (Figures 6H–6J). PLA also reduced lipid micelle secretion by Caco-2 cells (Figure 6K), suggesting that this *Lactobacillus*-derived metabolite may play a role in protecting the host against metabolic dysfunction during consumption of an HF diet by regulating lipid metabolism in IECs through PPAR- γ .

PLA protects young mice from HF diet-induced metabolic dysfunction

Based upon our *in vitro* data, we asked whether supplementation with PLA increased PPAR- γ abundance in the SI of mice exposed to an HF diet and LDP. In agreement with the untargeted metabolomics data (Figure 6A), mice exposed to an HF diet and LDP had significantly less PLA in their SI compared with mice given only an HF diet, and supplementation with PLA restored PLA abundance in HF diet and LDP-treated mice to a similar level as mice given an HF diet alone (Figure 7A). Elevated intestinal PLA (Figure 7A) significantly increased the abundance of PPAR- γ in the intestinal epithelium when mice were exposed to an HF diet and LDP compared to mice given only a HF diet (Figures 7B and 7C). Consistent with an increase in intestinal PPAR- γ , HF diet and LDP-exposed mice treated with PLA developed less adiposity and gained significantly less weight (Figures 7D, 7E, and S7A) than mice exposed to an HF diet and antibiotics. PLA supplementation also significantly inhibited LDP-induced fat accumulation in hepatocytes (Figures 7F and 7G) and reduced fasting blood glucose (relative to mice exposed to an HF diet and LDP) (Figure S7B). Changes in physiology due to PLA supplementation occurred despite mice treated with PLA consuming more food and having similar reductions in *Lactobacillus* abundance in the SI (Figures S7C and S7D). We determined that PLA, a metabolite produced by the SI resident gut microbiota, protects against early-life metabolic dysfunction during exposure to an HF diet by regulating PPAR- γ function in IECs.

DISCUSSION

The worldwide prevalence of childhood obesity has risen dramatically over the last 30 years.⁴³ In the United States,




Figure 6. Exposure to an HF diet and LDP depletes phenylactic acid, an activator of intestinal PPAR- γ , from the small intestine

(A and B) Untargeted metabolomics was performed on ileum content from mice given an HF diet or an HF diet + LDP (N = 5/phenotype). (A) Heatmap clustering of the experimental sample groups and metabolites. Samples (columns) are clustered by group and relative abundance (rows) across different groups, ordered from low (blue) to high (red) abundance. (B) Normalized relative abundance, as determined by untargeted metabolomics, of phenylactic acid (PLA) in the small intestine content of mice fed an HF diet or an HF diet + LDP-treated.

(C) Concentration (pmol/mg of intestinal content) of PLA in the small intestine of germ-free (GF) and conventional mice fed an HF diet for 5 weeks.

(D) Concentration (μM) of PLA in supernatants from defined microbiota cultures lacking or containing *L. murinus*.

(E) Concentration (μM) of PLA in the supernatant of lactic acid bacteria grown in cell culture media.

(F) Concentration (pmol/mg of intestinal content) of PLA in the small intestine of mice. Dotted line indicates limit of detection = 0.025 pmol/mg.

(G) Mouse enteroids were treated with either 10 μM Rosiglitazone or 5 mM PLA or left untreated.

(H and I) Immortalized mouse small intestinal cells were treated with 5 mM PLA (H) or left untreated (I). Cells were subsequently stained with anti-PPAR- γ and phalloidin to identify F-actin.

(J) PPAR- γ nuclear intensity was quantified using ImageJ; 15 cells were selected from two images from three independent experiments.

(K) Amount of BODIPY C₁₂ secreted by treated Caco-2 cells was measured 6 h after addition of lipid micelles.

(B and C) Each dot represents one animal. Lines represent geometric mean. (F) Each dot represents one animal. Lines represent mean. (B, C, and F) Data represent one independent cohort. n = 5 or 6 mice/group. (D and E) Each dot represents a biological replicate. Lines show geometric mean. (G) Each dot represents one technical replicate. Data shown from three biological replicates. Lines show geometric mean. (J) Violin plot showing the distribution of PPAR- γ nuclear intensity (n = 90 cells). (K) Each dot represents one technical replicate. Data shown from three biological replicates. Lines show geometric mean. *p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001 using an unpaired two-tailed Student's t test.

approximately 20% of school-age children are considered obese, resulting in a significant public health concern.⁴⁴ The etiology of childhood obesity is complex, and both genetic and environmental factors contribute to the development of the dis-

ease.⁴⁵ Recently, early-life disruption to the gut microbiota by antibiotic treatment has been proposed as a risk factor for childhood obesity.^{10,11} However, epidemiological surveys have produced conflicting results regarding whether early-life antibiotics




Figure 7. Phenylactic acid upregulates intestinal PPAR- γ and inhibits HF + LDP-induced metabolic dysfunction

(A) Concentration (pmol/mg of intestinal content) of PLA in the small intestine of mice. Dotted line indicates limit of detection = 0.025 pmol/mg. (B) Sections from the ileum of HF diet mice given 10 mM PLA in their drinking water and/or LDP for 5 weeks were stained for PPAR- γ . Scale bar represents 200 μ m. (C) PPAR- γ abundance in the epithelium was quantified by scoring blinded sections of the ileum. (D and E) (D) Abdominal fat (% total body weight) and (E) weight gain of mice measured after 5 weeks on an HF diet \pm PLA \pm LDP. (F) Representative images of oil red O-stained sections of the liver. Scale bar represents 100 μ m. (G) Combined oil red scores from livers of mice after 5 weeks of treatment. (A and G) $n = 6$ mice/group. (C–E) $n = 12$ mice/group. (A, C–E, and G) Each dot represents one animal. Lines represent geometric mean. Data representative of two independent cohorts. (A, D, and E) * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$ using an unpaired two-tailed Student's t test. (C and G) * $p < 0.05$; ** $p < 0.01$ using Mann-Whitney test.

correlate with a higher body mass index, prompting further research into how the early-life microbiota could be protective against obesity.^{8,46} Importantly, these epidemiological surveys did not consider the children's diet and therefore may be missing relevant synergistic effects between antibiotics and other environmental factors.^{8,46} To answer this gap in knowledge, we developed a mouse model in which young mice were concurrently exposed to antibiotics and an obesogenic HF diet. Specifically, we investigated the effects of penicillin, which belongs to the most commonly prescribed class of antibiotics in the United States.⁴⁷ In this paper, we determined that early-life penicillin exposure accelerates the development of HF diet-induced obesity by disrupting interactions between the SI microbiota and IECs. We further demonstrated that depletion of *Lactobacillus* species in the SI of mice given antibiotics and an HF diet led to excess adiposity. Specifically, an HF diet and antibiotics

reduced the abundance of the *Lactobacillus*-derived metabolite, PLA, which regulates PPAR- γ in IECs and protects against HF diet-induced obesity.

We determined that early-life concurrent exposure to antibiotics and an HF diet perturbs lipid metabolism in IECs and provides further evidence for the SI's significant role in regulating whole-body lipid homeostasis.⁴⁸ In obese patients, SI lipid metabolism is significantly altered, suggesting that dysregulation of intestinal lipid metabolism contributes to metabolic dysfunction.⁴⁹ Interestingly, proteins increased in the SI of obese patients⁴⁹ (FABP1, FABP2, MTP) corresponded to genes upregulated in the SI of mice exposed to an HF diet and antibiotics. We determined that mice given an HF diet and antibiotics accumulate significantly more triglycerides in their IECs and serum but did not absorb more fat from their diet. Together, these data support the hypothesis that concurrent exposure to an HF

diet and antibiotics perturbs intestinal lipid metabolism and leads to greater lipoprotein secretion, consistent with what is observed in obese patients.⁴⁹ We propose that these changes in intestinal lipid metabolism, at least in part, are due to loss of PPAR- γ signaling. Previous research has demonstrated that disruption to the gut microbiota decreases PPAR- γ activity in the intestine,³⁸ consistent with our observation that an HF diet and LDP treatment reduces PPAR- γ in the SI. The role of PPAR- γ in lipid metabolism is cell-type-specific. For instance, in the adipose tissue, PPAR- γ promotes lipid storage and lipogenesis,⁵⁰ whereas in macrophages, PPAR- γ reduces free fatty acid and triglyceride accumulation.⁵¹ In the intestine, deletion of PPAR- γ resulted in elevated serum triglycerides,⁴⁰ agreeing with our model in which depletion of intestinal PPAR- γ coincides with significantly greater triglycerides in fasted serum. Therefore, intestinal PPAR- γ presents a promising target to regulate lipid metabolism and reduce metabolic dysfunction during exposure to an HF diet. Indeed, treatment with the PPAR- γ agonist, Rosiglitazone, represses the expression of intestinal *apoB* and reduces triglyceride secretion in a hamster model.⁵² Collectively, these findings highlight the role of intestinal lipid metabolism, specifically PPAR- γ , in protecting against metabolic dysfunction.

The depletion of *Lactobacillus* species from the SI in mice exposed to an HF diet and antibiotics poses a significant threat to early-life microbiota-host interaction. *Lactobacillus* is one of the earliest colonizers of the gut and is a predominant member of the infant microbiota.^{4,53} Despite a reduction in the relative abundance of *Lactobacillus* as a child matures,⁴ *Lactobacillus* species remain key constituents of the gut microbiota.⁵⁴ Importantly, antibiotic treatment, specifically with penicillin, leads to long-lasting reductions in *Lactobacillus* in children.⁹ Therefore, investigations into how loss of *Lactobacillus* impacts host health and development remain clinically relevant. Previous research has determined that specific *Lactobacillus* species, *L. paracasei* F19 and *L. reuteri* ATCC 4659, limit the effects of an obesogenic HF diet in adult mice,^{55,56} but the mechanism by which these strains exert their protection remains relatively unknown. Furthermore, the ability of *Lactobacillus* to prevent metabolic dysfunction is species-specific as some species seem to offer no protection or exacerbate the effect of an HF diet.^{55,57} Even more so, probiotic intervention studies have shown that *Lactobacillus* species largely fail to alter body composition in children,^{58,59} highlighting the critical need to understand how *Lactobacillus* species protect against HF diet-induced obesity. In our paper, we determine that *L. murinus*, a predominant *Lactobacillus* species in the human gut,⁵⁴ inhibits lipid secretion from IECs by upregulating intestinal PPAR- γ .

This paper did not explore the impact of other SI microbiota members on host adiposity during early-life exposure to an HF diet. However, in our 16S rRNA sequencing of the SI microbiota of mice exposed to an HF diet and LDP, we determined that the relative abundance of the Streptococcaceae family was increased. As the *Streptococcus* species is a dominant group in the human SI microbiota,⁶⁰ this increase in our model warrants further investigation. *Streptococcus* species are highly abundant in atherosclerotic plaques and correlate positively with HDL cholesterol level,⁶¹ suggesting that elevated Streptococcaceae in HF diet and LDP-treated mice could be adding to their meta-

bolic dysfunction. Interestingly, commensal *Streptococcus* species have been shown to downregulate PPAR- γ activity,⁶² which may contribute to the inhibition of intestinal PPAR- γ in mice given an HF diet and LDP. Therefore, future work will investigate how interactions between multiple SI microbiota members influence host metabolism through intestinal PPAR- γ .

Due to the transient colonization of probiotics,⁶³ direct administration of the beneficial *Lactobacillus*-derived metabolite may have better success in preventing metabolic dysfunction. Therefore, we identified a *Lactobacillus*-derived metabolite, PLA, that activates intestinal PPAR- γ and protects young mice from metabolic dysfunction due to concurrent exposure to an HF diet and antibiotics. Multiple bacterial species produce PLA, including species belonging to the Bifidobacteriaceae and Peptostreptococcaceae families.^{64–66} Interestingly, we see a depletion of Bifidobacteriaceae and Peptostreptococcaceae in mice fed an HF diet and mice exposed to an HF diet and antibiotics (Figure 2A), suggesting that multiple bacterial species may contribute to PLA production in the intestine. Significantly, PLA is present in the feces of infants and correlates with the abundance of *Bifidobacterium*,⁶⁴ indicating that this metabolite is a component of the early-life gut metabolome in humans. Previous work revealed that PLA upregulated an adipose tissue-specific isoform of PPAR- γ ,⁴² and we have now determined that PLA also increases intestinal PPAR- γ activity and abundance. We then extended these findings by showing that PLA inhibits lipid secretion in IECs and prevents antibiotic-induced excess adiposity in young mice during consumption of an obesogenic HF diet. Collectively, we identified a previously unknown mechanism by which the early-life microbiota communicates with the intestinal epithelium to decrease HF-induced obesity, providing new evidence linking early-life antibiotic exposure to childhood obesity.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- RESOURCE AVAILABILITY
 - Lead contact
 - Materials availability
 - Data and code availability
- EXPERIMENTAL MODEL DETAILS
 - Animal Experiments
 - Bacterial culture conditions
 - Mouse small intestinal epithelial cells
 - Caco-2 cell culture
 - Mouse enteroids
- METHOD DETAILS
 - Body composition
 - Oil Red O Staining
 - 16s rRNA sequencing and data analysis
 - *L. murinus*-specific qPCR
 - Colonization of HF and LDP-treated mice with *L. murinus* Pen^R
 - Bacterial culture with bile salts and penicillin
 - Gnotobiotic mouse experiments
 - Histopathology Analysis

- Enterocyte isolation
- RNA extraction and quantitative real-time PCR
- RNA-sequencing
- Lipid Measurements
- Non-invasive Measurement of Intestinal Fat Absorption
- Oral fat tolerance test
- PPAR- γ Immunostaining
- Lipid secretion assay in Caco-2 cells
- PPAR- γ fluorescent staining
- PPAR Reporter Assay
- Co-culture of bacterial strains and intestinal epithelial cells
- Untargeted metabolomics
- Preparation of defined microbiota supernatants
- Phenylactic Acid Measurements

● **QUANTIFICATION AND STATISTICAL ANALYSIS**

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.chom.2023.09.002>.

ACKNOWLEDGMENTS

We would like to acknowledge the Translational Pathology Shared Resource at Vanderbilt University Medical Center, which is supported by NCI/NIH Cancer Center support grant P30CA068485 and the Shared Instrumentation grant S10 OD023475-01A1 for the Leica Bond RX. Additionally, we would like to thank the University of California, San Diego (UCSD) Microbiome Core as this publication includes data generated at the UC San Diego IGM Genomics Center utilizing an Illumina NovaSeq 6000 that was purchased with funding from a National Institutes of Health SIG grant (S10 OD026929). This work was supported in part using the resources of the Center for Innovative Technology (CIT) at Vanderbilt University. The authors also wish to acknowledge the expert technical support of the Vanderbilt Technologies for Advanced Genomics Analysis and Research Design (VANGARD) core facilities. Immunofluorescent images were collected through the use of the Vanderbilt Cell Imaging Shared Resource (supported by NIH grants CA68485, DK20593, DK58404, DK59637, and EY08126) using a Zeiss LSM 880 Airyscan confocal microscope (supported by NIH grant 1 S10 OD021630 1). C.D.S. was supported by the NIH (T32AI112541 and 1F31AI161882-01A1), 2021-2022 Danone North America Gut Microbiome, Yogurt and Probiotics Fellowship, and VICTR (VR55193 and VR54476). N.G.S. was supported by NIH T32 training grant T32ES007028-46 and GT15104 from the Howard Hughes Medical Institute through the James H. Gilliam Fellowships for Advanced Study program. N.J.F. was supported by NIH T32 training grant T32DK007673. Work in N.O.M.'s lab was funded by Department of Veterans Affairs (BX005699). Worked in S.H.P.'s lab was funded by the Department of Veterans Affairs (11K2BX005401) and funds from the Department of Medicine at Vanderbilt University Medical Center. M.X.B. is an HHMI Freeman Hrabowski Scholar. Work in M.X.B.'s lab was funded by The V Foundation for Cancer Research V Scholar grant (V2020-013), the NIH (R01DK131104-01 and 1R01AI168302-01A1), The Pew Charitable Trusts (2022-A-19568), and the Vanderbilt Institute for Clinical and Translational Research (VICTR) (VR53102). This article is subject to HHMI's Open Access to Publications policy. HHMI lab heads have previously granted a nonexclusive CC BY 4.0 license to the public and a sublicensable license to HHMI in their research articles. Pursuant to those licenses, the author-accepted manuscript of this article can be made freely available under a CC BY 4.0 license immediately upon publication.

AUTHOR CONTRIBUTIONS

M.X.B. and C.D.S. designed and conceived the study. C.D.S., J.M., N.G.S., E.S., and J.T. performed all experiments. W.Y., G.N.F., T.P.T., N.J.F., and

J.K.Z. assisted with experimental design and data collection. M.W.C. performed phenylactic measurements. S.G.C., S.D.S., and J.A.M. executed the untargeted metabolomics sample preparation, data acquisition, and data analysis. T.T.H. and S.H.P. performed DEXA scanning. F.Y. assisted with the experimental design and data collection for all work done with the MSIE cells. N.O.M. provided technical expertise and reagents for culturing mouse enteroids. M.L. performed non-invasive measurements of intestinal fat absorption. All authors contributed to the data analysis and preparation of the manuscript. C.D.S. and M.X.B. wrote the manuscript and all authors reviewed it.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: April 6, 2023

Revised: August 7, 2023

Accepted: September 6, 2023

Published: October 3, 2023

REFERENCES

1. Gensollen, T., Iyer, S.S., Kasper, D.L., and Blumberg, R.S. (2016). How colonization by microbiota in early life shapes the immune system. *Science* 352, 539–544. <https://doi.org/10.1126/science.aad9378>.
2. Maynard, C.L., Elson, C.O., Hatton, R.D., and Weaver, C.T. (2012). Reciprocal interactions of the intestinal microbiota and immune system. *Nature* 489, 231–241. <https://doi.org/10.1038/nature11551>.
3. Derrien, M., Alvarez, A.S., and de Vos, W.M. (2019). The gut microbiota in the first decade of life. *Trends Microbiol.* 27, 997–1010. <https://doi.org/10.1016/j.tim.2019.08.001>.
4. Roswall, J., Olsson, L.M., Kovatcheva-Datchary, P., Nilsson, S., Tremaroli, V., Simon, M.C., Kiilerich, P., Akrami, R., Krämer, M., Uhlén, M., et al. (2021). Developmental trajectory of the healthy human gut microbiota during the first 5 years of life. *Cell Host Microbe* 29, 765–776.e3. e763. <https://doi.org/10.1016/j.chom.2021.02.021>.
5. Yatsunenko, T., Rey, F.E., Manary, M.J., Trehan, I., Dominguez-Bello, M.G., Contreras, M., Magris, M., Hidalgo, G., Baldassano, R.N., Anokhin, A.P., et al. (2012). Human gut microbiome viewed across age and geography. *Nature* 486, 222–227. <https://doi.org/10.1038/nature11053>.
6. Bokulich, N.A., Chung, J., Battaglia, T., Henderson, N., Jay, M., Li, H., D.L., A., Wu, F., Perez-Perez, G.I., Chen, Y., et al. (2016). Antibiotics, birth mode, and diet shape microbiome maturation during early life. *Sci. Transl. Med.* 8, 343ra382. <https://doi.org/10.1126/scitranslmed.aad7121>.
7. McNamara, M.P., Singleton, J.M., Cadney, M.D., Ruegger, P.M., Borneman, J., and Garland, T. (2021). Early-life effects of juvenile Western diet and exercise on adult gut microbiome composition in mice. *J. Exp. Biol.* 224, jeb239699. <https://doi.org/10.1242/jeb.239699>.
8. Chelimo, C., Camargo, C.A., Jr., Morton, S.M.B., and Grant, C.C. (2020). Association of repeated antibiotic exposure up to Age 4 years with body mass at age 4.5 years. *JAMA Netw. Open* 3, e1917577. <https://doi.org/10.1001/jamanetworkopen.2019.17577>.
9. Korpela, K., Salonen, A., Virta, L.J., Kekkonen, R.A., Forslund, K., Bork, P., and de Vos, W.M. (2016). Intestinal microbiome is related to lifetime antibiotic use in Finnish pre-school children. *Nat. Commun.* 7, 10410. <https://doi.org/10.1038/ncomms10410>.
10. Cho, I., Yamanishi, S., Cox, L., Methé, B.A., Zavadil, J., Li, K., Gao, Z., Mahana, D., Raju, K., Teitler, I., et al. (2012). Antibiotics in early life alter the murine colonic microbiome and adiposity. *Nature* 488, 621–626. <https://doi.org/10.1038/nature11400>.
11. Cox, L.M., Yamanishi, S., Sohn, J., Alekseyenko, A.V., Leung, J.M., Cho, I., Kim, S.G., Li, H., Gao, Z., Mahana, D., et al. (2014). Altering the intestinal microbiota during a critical developmental window has lasting metabolic consequences. *Cell* 158, 705–721. <https://doi.org/10.1016/j.cell.2014.05.052>.

12. Yoo, W., Zieba, J.K., Foegeeding, N.J., Torres, T.P., Shelton, C.D., Shealy, N.G., Byndloss, A.J., Cevallos, S.A., Gertz, E., Tiffany, C.R., et al. (2021). High-fat diet-induced colonocyte dysfunction escalates microbiota-derived trimethylamine N-oxide. *Science* 373, 813–818. <https://doi.org/10.1126/science.aba3683>.
13. Turnbaugh, P.J., Backhed, F., Fulton, L., and Gordon, J.I. (2008). Diet-induced obesity is linked to marked but reversible alterations in the mouse distal gut microbiome. *Cell Host Microbe* 3, 213–223. <https://doi.org/10.1016/j.chom.2008.02.015>.
14. Wang, Y., Guglielmo, D., and Welsh, J.A. (2018). Consumption of sugars, saturated fat, and sodium among US children from infancy through preschool age, NHANES 2009–2014. *Am. J. Clin. Nutr.* 108, 868–877. <https://doi.org/10.1093/ajcn/nqy168>.
15. Zetts, R.M., Stoesz, A., Smith, B.A., and Hyun, D.Y. (2018). Outpatient antibiotic use and the need for increased antibiotic stewardship efforts. *Pediatrics* 141, e20174124. <https://doi.org/10.1542/peds.2017-4124>.
16. Clara, R., Schumacher, M., Ramachandran, D., Fedele, S., Krieger, J.P., Langhans, W., and Mansouri, A. (2017). Metabolic adaptation of the small intestine to short- and medium-term high-fat diet exposure. *J. Cell. Physiol.* 232, 167–175. <https://doi.org/10.1002/jcp.25402>.
17. Zembroski, A.S., D'Aquila, T., and Buhman, K.K. (2021). Characterization of cytoplasmic lipid droplets in each region of the small intestine of lean and diet-induced obese mice in response to dietary fat. *Am. J. Physiol. Gastrointest. Liver Physiol.* 321, G75–G86. <https://doi.org/10.1152/ajpgi.00084.2021>.
18. Kondo, H., Minegishi, Y., Komine, Y., Mori, T., Matsumoto, I., Abe, K., Tokimitsu, I., Hase, T., and Murase, T. (2006). Differential regulation of intestinal lipid metabolism-related genes in obesity-resistant A/J vs. obesity-prone C57BL/6J mice. *Am. J. Physiol. Endocrinol. Metab.* 291, E1092–E1099. <https://doi.org/10.1152/ajpendo.00583.2005>.
19. Martinez-Guryn, K., Hubert, N., Frazier, K., Urlass, S., Musch, M.W., Ojeda, P., Pierre, J.F., Miyoshi, J., Sontag, T.J., Cham, C.M., et al. (2018). Small intestine microbiota regulate Host Digestive and Absorptive Adaptive Responses to Dietary Lipids. *Cell Host Microbe* 23, 458–469.e5. <https://doi.org/10.1016/j.chom.2018.03.011>.
20. Semova, I., Carten, J.D., Stombaugh, J., Mackey, L.C., Knight, R., Farber, S.A., and Rawls, J.F. (2012). Microbiota regulate intestinal absorption and metabolism of fatty acids in the zebrafish. *Cell Host Microbe* 12, 277–288. <https://doi.org/10.1016/j.chom.2012.08.003>.
21. El Aidy, S., Merrifield, C.A., Derrien, M., van Baarlen, P., Hooiveld, G., Levenez, F., Doré, J., Dekker, J., Holmes, E., Claus, S.P., et al. (2013). The gut microbiota elicits a profound metabolic reorientation in the mouse jejunal mucosa during conventionalisation. *Gut* 62, 1306–1314. <https://doi.org/10.1136/gutjnl-2011-301955>.
22. Tazi, A., Araujo, J.R., Mulet, C., Arena, E.T., Nigro, G., Pédrón, T., and Sansonetti, P.J. (2018). Disentangling Host-Microbiota Regulation of Lipid Secretion by Enterocytes: insights from Commensals *Lactobacillus paracasei* and *Escherichia coli*. *mBio* 9, e01493–e01418. <https://doi.org/10.1128/mBio.01493-18>.
23. Araújo, J.R., Tazi, A., Burlen-Defranoux, O., Vichier-Guerre, S., Nigro, G., Licandro, H., Demignot, S., and Sansonetti, P.J. (2020). Fermentation products of commensal bacteria alter enterocyte lipid metabolism. *Cell Host Microbe* 27, 358–375.e7. <https://doi.org/10.1016/j.chom.2020.01.028>.
24. Kappe, C., Zhang, Q., Nyström, T., and Sjöholm, A. (2014). Effects of high-fat diet and the anti-diabetic drug metformin on circulating GLP-1 and the relative number of intestinal L-cells. *Diabetol. Metab. Syndr.* 6, 70. <https://doi.org/10.1186/1758-5996-6-70>.
25. Poteres, E., Hubert, N., Poludasu, S., Brigando, G., Moore, J., Keeler, K., Isabelli, A., Ibay, I.C.V., Alt, L., Pytynia, M., et al. (2020). Selective regional alteration of the gut microbiota by diet and antibiotics. *Front. Physiol.* 11, 797. <https://doi.org/10.3389/fphys.2020.00797>.
26. Murakami, Y., Tanabe, S., and Suzuki, T. (2016). High-fat diet-induced intestinal hyperpermeability is associated with increased bile acids in the large intestine of mice. *J. Food Sci.* 81, H216–H222. <https://doi.org/10.1111/1750-3841.13166>.
27. Andersson, D.I., and Hughes, D. (2014). Microbiological effects of sublethal levels of antibiotics. *Nat. Rev. Microbiol.* 12, 465–478. <https://doi.org/10.1038/nrmicro3270>.
28. Chen, J., Ying, G.G., and Deng, W.J. (2019). Antibiotic residues in food: extraction, analysis, and human health concerns. *J. Agric. Food Chem.* 67, 7569–7586. <https://doi.org/10.1021/acs.jafc.9b01334>.
29. Everard, A., Geurts, L., Caesar, R., Van Hul, M., Matamoros, S., Duparc, T., Denis, R.G., Cochez, P., Pierard, F., Castel, J., et al. (2014). Intestinal epithelial MyD88 is a sensor switching host metabolism towards obesity according to nutritional status. *Nat. Commun.* 5, 5648. <https://doi.org/10.1038/ncomms6648>.
30. Wang, Y., Kuang, Z., Yu, X., Ruhn, K.A., Kubo, M., and Hooper, L.V. (2017). The intestinal microbiota regulates body composition through NFIL3 and the circadian clock. *Science* 357, 912–916. <https://doi.org/10.1126/science.aan0677>.
31. Ko, C.W., Qu, J., Black, D.D., and Tso, P. (2020). Regulation of intestinal lipid metabolism: current concepts and relevance to disease. *Nat. Rev. Gastroenterol. Hepatol.* 17, 169–183. <https://doi.org/10.1038/s41575-019-0250-7>.
32. Ntambi, J.M., Miyazaki, M., Stoehr, J.P., Lan, H., Kendziorski, C.M., Yandell, B.S., Song, Y., Cohen, P., Friedman, J.M., and Attie, A.D. (2002). Loss of stearoyl-CoA desaturase-1 function protects mice against adiposity. *Proc. Natl. Acad. Sci. USA* 99, 11482–11486. <https://doi.org/10.1073/pnas.132384699>.
33. Hussain, M.M., Kancha, R.K., Zhou, Z., Luchoomun, J., Zu, H., and Bakillah, A. (1996). Chylomicron assembly and catabolism: role of apolipoproteins and receptors. *Biochim. Biophys. Acta* 1300, 151–170. [https://doi.org/10.1016/0005-2760\(96\)00041-0](https://doi.org/10.1016/0005-2760(96)00041-0).
34. Mattijssen, F., Alex, S., Swarts, H.J., Groen, A.K., van Schothorst, E.M., and Kersten, S. (2014). Angptl4 serves as an endogenous inhibitor of intestinal lipid digestion. *Mol. Metab.* 3, 135–144. <https://doi.org/10.1016/j.molmet.2013.11.004>.
35. Lynes, M., Narisawa, S., Millán, J.L., and Widmaier, E.P. (2011). Interactions between CD36 and global intestinal alkaline phosphatase in mouse small intestine and effects of high-fat diet. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 301, R1738–R1747. <https://doi.org/10.1152/ajpregu.00235.2011>.
36. Lagakos, W.S., Gajda, A.M., Agellon, L., Binas, B., Choi, V., Mandap, B., Russnak, T., Zhou, Y.X., and Storch, J. (2011). Different functions of intestinal and liver-type fatty acid-binding proteins in intestine and in whole body energy homeostasis. *Am. J. Physiol. Gastrointest. Liver Physiol.* 300, G803–G814. <https://doi.org/10.1152/ajpgi.00229.2010>.
37. Peng, H., Chiu, T.Y., Liang, Y.J., Lee, C.J., Liu, C.S., Suen, C.S., Yen, J.J., Chen, H.T., Hwang, M.J., Hussain, M.M., et al. (2021). PRAP1 is a novel lipid-binding protein that promotes lipid absorption by facilitating MTP-mediated lipid transport. *J. Biol. Chem.* 296, 100052. <https://doi.org/10.1074/jbc.RA120.015002>.
38. Byndloss, M.X., Olsan, E.E., Rivera-Chávez, F., Tiffany, C.R., Cevallos, S.A., Lokken, K.L., Torres, T.P., Byndloss, A.J., Faber, F., Gao, Y., et al. (2017). Microbiota-activated PPAR-gamma signaling inhibits dysbiotic Enterobacteriaceae expansion. *Science* 357, 570–575. <https://doi.org/10.1126/science.aam9949>.
39. Alex, S., Lange, K., Amolo, T., Grinstead, J.S., Haakonsson, A.K., Szalowska, E., Koppen, A., Mudde, K., Haenen, D., Al-Lahham, S., et al. (2013). Short-chain fatty acids stimulate angiopoietin-like 4 synthesis in human colon adenocarcinoma cells by activating peroxisome proliferator-activated receptor gamma. *Mol. Cell. Biol.* 33, 1303–1316. <https://doi.org/10.1128/MCB.00858-12>.
40. Duszka, K., Oresic, M., Le May, C., König, J., and Wahli, W. (2017). PPARgamma modulates long chain fatty acid processing in the intestinal epithelium. *Int. J. Mol. Sci.* 18, 2559. <https://doi.org/10.3390/ijms18122559>.
41. Nepelska, M., de Wouters, T., Jacouton, E., Béguet-Crespel, F., Lapaque, N., Doré, J., Arulampalam, V., and Blottière, H.M. (2017). Commensal gut

- bacteria modulate phosphorylation-dependent PPARgamma transcriptional activity in human intestinal epithelial cells. *Sci. Rep.* 7, 43199. <https://doi.org/10.1038/srep43199>.
42. Ilavenil, S., Kim, D.H., Valan Arasu, M., Srigopalram, S., Sivanesan, R., and Choi, K.C. (2015). Phenylactic acid from *Lactobacillus plantarum* promotes adipogenic activity in 3T3-L1 Adipocyte via up-regulation of PPAR-gamma2. *Molecules* 20, 15359–15373. <https://doi.org/10.3390/molecules200815359>.
 43. Di Cesare, M., Sorić, M., Bovet, P., Miranda, J.J., Bhutta, Z., Stevens, G.A., Laxmaiah, A., Kengne, A.P., and Bentham, J. (2019). The epidemiological burden of obesity in childhood: a worldwide epidemic requiring urgent action. *BMC Med.* 17, 212. <https://doi.org/10.1186/s12916-019-1449-8>.
 44. National Center for Health (2021). National health and nutrition examination survey 2017–March 2020 prepandemic data files development of files and prevalence estimates for selected health outcomes. National Health Statistics Reports. <https://doi.org/10.15620/cdc:106273>.
 45. Kumar, S., and Kelly, A.S. (2017). Review of childhood obesity: from epidemiology, etiology, and comorbidities to clinical assessment and treatment. *Mayo Clin. Proc.* 92, 251–265. <https://doi.org/10.1016/j.mayocp.2016.09.017>.
 46. Leong, K.S.W., McLay, J., Derrai, J.G.B., Gibb, S., Shackleton, N., Taylor, R.W., Glover, M., Audas, R., Taylor, B., Milne, B.J., and Cutfield, W.S. (2020). Associations of prenatal and childhood antibiotic exposure with obesity at Age 4 years. *JAMA Netw. Open* 3, e1919681. <https://doi.org/10.1001/jamanetworkopen.2019.19681>.
 47. Centers for Disease Control and Prevention (2021). Outpatient antibiotic prescriptions—United States. <https://www.cdc.gov/antibiotic-use/data/report-2021.html>.
 48. Duca, F.A., Waise, T.M.Z., Peppler, W.T., and Lam, T.K.T. (2021). The metabolic impact of small intestinal nutrient sensing. *Nat. Commun.* 12, 903. <https://doi.org/10.1038/s41467-021-21235-y>.
 49. Veilleux, A., Grenier, E., Marceau, P., Carpentier, A.C., Richard, D., and Levy, E. (2014). Intestinal lipid handling: evidence and implication of insulin signaling abnormalities in human obese subjects. *Arterioscler. Thromb. Vasc. Biol.* 34, 644–653. <https://doi.org/10.1161/ATVBAHA.113.302993>.
 50. He, W. (2013). Adipose tissue-specific PPARgamma gene targeting. *Methods Mol. Biol.* 952, 117–135. https://doi.org/10.1007/978-1-62703-155-4_8.
 51. Ye, G., Gao, H., Wang, Z., Lin, Y., Liao, X., Zhang, H., Chi, Y., Zhu, H., and Dong, S. (2019). PPARalpha and PPARgamma activation attenuates total free fatty acid and triglyceride accumulation in macrophages via the inhibition of Fatp1 expression. *Cell Death Dis.* 10, 39. <https://doi.org/10.1038/s41419-018-1135-3>.
 52. Lewis, G.F., Uffelman, K., Naples, M., Szeto, L., Haidari, M., and Adeli, K. (2005). Intestinal lipoprotein overproduction, a newly recognized component of insulin resistance, is ameliorated by the insulin sensitizer rosiglitazone: studies in the fructose-fed Syrian golden hamster. *Endocrinology* 146, 247–255. <https://doi.org/10.1210/en.2004-1143>.
 53. Dominguez-Bello, M.G., Costello, E.K., Contreras, M., Magris, M., Hidalgo, G., Fierer, N., and Knight, R. (2010). Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc. Natl. Acad. Sci. USA* 107, 11971–11975. <https://doi.org/10.1073/pnas.1002601107>.
 54. Rossi, M., Martínez-Martínez, D., Amaretti, A., Ulrici, A., Raimondi, S., and Moya, A. (2016). Mining metagenomic whole genome sequences revealed subdominant but constant *Lactobacillus* population in the human gut microbiota. *Environ. Microbiol. Rep.* 8, 399–406. <https://doi.org/10.1111/1758-2229.12405>.
 55. Fåk, F., and Bäckhed, F. (2012). *Lactobacillus reuteri* prevents diet-induced obesity, but not atherosclerosis, in a strain dependent fashion in ApoE^{-/-} mice. *PLoS One* 7, e46837. <https://doi.org/10.1371/journal.pone.0046837>.
 56. Aronsson, L., Huang, Y., Parini, P., Korach-André, M., Håkansson, J., Gustafsson, J.Å., Pettersson, S., Arulampalam, V., and Rafter, J. (2010). Decreased fat storage by *Lactobacillus paracasei* is associated with increased levels of angiopoietin-like 4 protein (ANGPTL4). *PLoS One* 5, e13087. <https://doi.org/10.1371/journal.pone.0013087>.
 57. Zhong, W., Wang, H., Yang, Y., Zhang, Y., Lai, H., Cheng, Y., Yu, H., Feng, N., Huang, R., Liu, S., et al. (2022). High-protein diet prevents fat mass increase after dieting by counteracting *Lactobacillus*-enhanced lipid absorption. *Nat. Metab.* 4, 1713–1731. <https://doi.org/10.1038/s42255-022-00687-6>.
 58. Karlsson Videhult, F., Öhlund, I., Stenlund, H., Hernell, O., and West, C.E. (2015). Probiotics during weaning: a follow-up study on effects on body composition and metabolic markers at school age. *Eur. J. Nutr.* 54, 355–363. <https://doi.org/10.1007/s00394-014-0715-y>.
 59. Luoto, R., Kalliomäki, M., Laitinen, K., and Isolauri, E. (2010). The impact of perinatal probiotic intervention on the development of overweight and obesity: follow-up study from birth to 10 years. *Int. J. Obes. (Lond)* 34, 1531–1537. <https://doi.org/10.1038/ijo.2010.50>.
 60. Zoetendal, E.G., Raes, J., van den Bogert, B., Arumugam, M., Booijink, C.C., Troost, F.J., Bork, P., Wels, M., de Vos, W.M., and Kleerebezem, M. (2012). The human small intestinal microbiota is driven by rapid uptake and conversion of simple carbohydrates. *ISME J.* 6, 1415–1426. <https://doi.org/10.1038/ismej.2011.212>.
 61. Koren, O., Spor, A., Felin, J., Fåk, F., Stombaugh, J., Tremaroli, V., Behre, C.J., Knight, R., Fagerberg, B., Ley, R.E., and Backhed, F. (2011). Human oral, gut, and plaque microbiota in patients with atherosclerosis. *Proc. Natl. Acad. Sci. USA* 108, 4592–4598. <https://doi.org/10.1073/pnas.1011383107>.
 62. Couvigny, B., de Wouters, T., Kaci, G., Jacouton, E., Delorme, C., Doré, J., Renault, P., Blottière, H.M., Guédon, E., and Lapaque, N. (2015). Commensal *Streptococcus salivarius* modulates PPARgamma transcriptional activity in human intestinal epithelial cells. *PLoS One* 10, e0125371. <https://doi.org/10.1371/journal.pone.0125371>.
 63. Maldonado-Gómez, M.X., Martínez, I., Bottacini, F., O'Callaghan, A., Ventura, M., van Sinderen, D., Hillmann, B., Vangay, P., Knights, D., Hutkins, R.W., and Walter, J. (2016). Stable engraftment of *Bifidobacterium longum* AH1206 in the human gut depends on individualized features of the resident microbiome. *Cell Host Microbe* 20, 515–526. <https://doi.org/10.1016/j.chom.2016.09.001>.
 64. Laursen, M.F., Sakanaka, M., von Burg, N., Mörbe, U., Andersen, D., Moll, J.M., Pekmez, C.T., Rivollier, A., Michaelsen, K.F., Molgaard, C., et al. (2021). *Bifidobacterium* species associated with breastfeeding produce aromatic lactic acids in the infant gut. *Nat. Microbiol.* 6, 1367–1382. <https://doi.org/10.1038/s41564-021-00970-4>.
 65. Valerio, F., Lavermicocca, P., Pascale, M., and Visconti, A. (2004). Production of phenylactic acid by lactic acid bacteria: an approach to the selection of strains contributing to food quality and preservation. *FEMS Microbiol. Lett.* 233, 289–295. <https://doi.org/10.1016/j.femsle.2004.02.020>.
 66. Yu, S., Zhou, C., Zhang, T., Jiang, B., and Mu, W. (2015). Short communication: 3-phenylactic acid production in milk by *Pediococcus pentosaceus* SK25 during laboratory fermentation process. *J. Dairy Sci.* 98, 813–817. <https://doi.org/10.3168/jds.2014-8645>.
 67. Whitehead, R.H., VanEeden, P.E., Noble, M.D., Ataliotis, P., and Jat, P.S. (1993). Establishment of conditionally immortalized epithelial cell lines from both colon and small intestine of adult H-2Kb-tsA58 transgenic mice. *Proc. Natl. Acad. Sci. USA* 90, 587–591. <https://doi.org/10.1073/pnas.90.2.587>.
 68. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21. <https://doi.org/10.1093/bioinformatics/bts635>.
 69. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30, 923–930. <https://doi.org/10.1093/bioinformatics/btt656>.

70. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550. <https://doi.org/10.1186/s13059-014-0550-8>.
71. Murphy, R., Stewart, A.W., Braithwaite, I., Beasley, R., Hancox, R.J., and Mitchell, E.A.; ISAAC Phase Three Study Group (2014). Antibiotic treatment during infancy and increased body mass index in boys: an international cross-sectional study. *Int. J. Obes. (Lond)* **38**, 1115–1119. <https://doi.org/10.1038/ijo.2013.218>.
72. Salinero, A.E., Anderson, B.M., and Zuloaga, K.L. (2018). Sex differences in the metabolic effects of diet-induced obesity vary by age of onset. *Int. J. Obes. (Lond)* **42**, 1088–1091. <https://doi.org/10.1038/s41366-018-0023-3>.
73. Nobel, Y.R., Cox, L.M., Kirigin, F.F., Bokulich, N.A., Yamanishi, S., Teitler, I., Chung, J., Sohn, J., Barber, C.M., Goldfarb, D.S., et al. (2015). Metabolic and metagenomic outcomes from early-life pulsed antibiotic treatment. *Nat. Commun.* **6**, 7486. <https://doi.org/10.1038/ncomms8486>.
74. Eberl, C., Ring, D., Münch, P.C., Beutler, M., Basic, M., Slack, E.C., Schwarzer, M., Srutkova, D., Lange, A., Frick, J.S., et al. (2019). Reproducible colonization of Germ-free mice with the oligo-mouse-microbiota in different animal facilities. *Front. Microbiol.* **10**, 2999. <https://doi.org/10.3389/fmicb.2019.02999>.
75. Bacic, M.K., and Smith, C.J. (2008). Laboratory maintenance and cultivation of bacteroides species. *Curr. Protoc. Microbiol. Chapter 13*. Unit 13C.1. <https://doi.org/10.1002/9780471729259.mc13c01s9>.
76. Yildiz, S., Pereira Bonifacio Lopes, J.P., Bergé, M., González-Ruiz, V., Baud, D., Kloehn, J., Boal-Carvalho, I., Schaeren, O.P., Schotsaert, M., Hathaway, L.J., et al. (2020). Respiratory tissue-associated commensal bacteria offer therapeutic potential against pneumococcal colonization. *eLife* **9**, e53581. <https://doi.org/10.7554/eLife.53581>.
77. Barman, M., Unold, D., Shifley, K., Amir, E., Hung, K., Bos, N., and Salzman, N. (2008). Enteric salmonellosis disrupts the microbial ecology of the murine gastrointestinal tract. *Infect. Immun.* **76**, 907–915. <https://doi.org/10.1128/IAI.01432-07>.
78. Jandacek, R.J., Heubi, J.E., and Tso, P. (2004). A novel, noninvasive method for the measurement of intestinal fat absorption. *Gastroenterology* **127**, 139–144. <https://doi.org/10.1053/j.gastro.2004.04.007>.
79. Pauquai, T., Bouchoux, J., Chateau, D., Vidal, R., Rousset, M., Chambaz, J., and Demignot, S. (2006). Adaptation of enterocytic Caco-2 cells to glucose modulates triacylglycerol-rich lipoprotein secretion through triacylglycerol targeting into the endoplasmic reticulum lumen. *Biochem. J.* **395**, 393–403. <https://doi.org/10.1042/BJ20051359>.
80. Kim, J.B., Wright, H.M., Wright, M., and Spiegelman, B.M. (1998). ADD1/SREBP1 activates PPARgamma through the production of endogenous ligand. *Proc. Natl. Acad. Sci. USA* **95**, 4333–4337. <https://doi.org/10.1073/pnas.95.8.4333>.
81. Popay, T.M., Wang, J., Adams, C.M., Howard, G.C., Codreanu, S.G., Sherrod, S.D., McLean, J.A., Thomas, L.R., Lorey, S.L., Machida, Y.J., et al. (2021). MYC regulates ribosome biogenesis and mitochondrial gene expression programs through its interaction with host cell factor-1. *eLife* **10**, e60191. <https://doi.org/10.7554/eLife.60191>.
82. Wishart, D.S., Jewison, T., Guo, A.C., Wilson, M., Knox, C., Liu, Y., Djombou, Y., Mandal, R., Aziat, F., Dong, E., et al. (2013). HMDB 3.0—the human metabolome database in 2013. *Nucleic Acids Res.* **41**, D801–D807. <https://doi.org/10.1093/nar/gks1065>.
83. Smith, C.A., O'Maille, G., Want, E.J., Qin, C., Trauger, S.A., Brandon, T.R., Custodio, D.E., Abagyan, R., and Siuzdak, G. (2005). Metlin: a metabolite mass spectral database. *Ther. Drug Monit.* **27**, 747–751. <https://doi.org/10.1097/01.ftd.0000179845.53213.39>.
84. Jablonski, F., Salvat, F., Powell, C.J., and Lee, A.Y. (2016). NIST standard reference database number 64. <https://doi.org/10.6028/NIST.NSRDS.64>.
85. Schrimpe-Rutledge, A.C., Codreanu, S.G., Sherrod, S.D., and McLean, J.A. (2016). Untargeted metabolomics strategies-challenges and emerging directions. *J. Am. Soc. Mass Spectrom.* **27**, 1897–1905. <https://doi.org/10.1007/s13361-016-1469-y>.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal anti-PPAR- γ	Santa Cruz Biotechnology	Cat #: sc-7273, RRID:AB_628115
Goat anti-mouse IgG Alexa Fluor 488	Thermo Fisher Scientific	Cat #: A-11001, RRID:AB_2534069
Bacterial and virus strains		
<i>Ligilactobacillus murinus</i>	This study	CS010
<i>Muribaculum intestinale</i>	DSMZ	YL27
<i>Clostridium clostridioforme</i>	DSMZ	YL32
<i>Clostridium sporogenes</i>	DSMZ	B-CC-163-3B
<i>Streptococcus daniellae</i> ERD01G	DSMZ	ERD01G
<i>Staphylococcus xyloso</i> 33-ERD13C	DSMZ	33-ERD13C
<i>Escherichia coli</i> Mt1B1	DSMZ	Mt1B1
<i>Ligilactobacillus murinus</i> Pen ^R	This study	CS052
<i>Ligilactobacillus murinus</i>	DSMZ	M-6244-3B
<i>Lactocaseibacillus rhamnosus</i>	ATCC	Cat #: 53103
<i>Lactobacillus acidophilus</i>	ATCC	Scav; Cat #: 4356
<i>Bifidobacterium longum subsp animalis</i>	DSMZ	YL2
Chemicals, peptides, and recombinant proteins		
Difco Lactobacilli MRS agar	Becton Dickinson	Cat #: 288210
Difco Lactobacilli MRS broth	Becton Dickinson	Cat #: 288130
Penicillin V potassium salt	Sigma	Cat #: P1382
D-(+)-3-phenyllactic acid	Sigma	Cat #: 376906
Difco LB broth	Becton Dickinson	Cat #: 244620
Difco Brain Heart Infusion agar	Becton Dickinson	Cat #: 241830
Bacto Yeast Extract	Becton Dickinson	Cat #: 212750
Bacto Tryptic soy broth	Becton Dickinson	Cat #: 211825
Dipotassium hydrogen phosphate	Fisher Chemicals	Cat #: P288
Hemin	Sigma	Cat #: 51280
D(+)-Glucose	Acros Organics	Cat #: 41095
Sodium carbonate	Fisher Chemicals	Cat #: 5263
(L) – cysteine	Sigma	Cat #: 168149
Menadione	Sigma	Cat #: 47775
Heat-Inactivated Fetal Bovine Serum	Gibco	Cat #: 16140-071
RPMI 1640 Medium	Gibco	Cat #: 11875093
Recombinant Mouse IFN-gamma Protein	R&D Systems	Cat #: 485-MI-100
Penicillin-Streptomycin	Gibco	Cat #: 15140122
ITS Liquid Media Supplement	Sigma	Cat #: I3146
Minimal Essential Media	Gibco	Cat #: 11095-080
Glutamax	Gibco	Cat #: 35050-061
Nonessential amino acids	Gibco	Cat #: 11140-050
Sodium pyruvate	Gibco	Cat #: 11360070
Trypsin-EDTA (0.05%)	Gibco	Cat #: 25300054
Primocin	InvivoGen	Cat #: ant-pm-1
Dithiothreitol	Acros Organics	Cat #: 16568
EDTA	Sigma	Cat #: E5134
EGTA	Sigma	Cat #: E3889

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Matrigel Membrane matrix	Corning	Cat #: 354234
Intesticult Organoid Growth Medium (Mouse)	STEMCELL Technologies	Cat #: 06005
Y-27632 dihydrochloride	TOCRIS	Cat #: 1254
Rosiglitazone	Sigma	Cat #: R2408
O.C.T. Compound	Fisher Chemicals	Cat #: 4585
iQ SYBR Green Supermix	BioRad	Cat #: 1708880
Bile salts	Sigma	Cat #: B8756
Lipid mixture	Sigma	Cat #: L0288
TRIzol Reagent	Invitrogen	Cat #: 15596026
QIAzol Lysis Reagent	Qiagen	Cat #: 79306
BioRad Protein Reagent	BioRad	Cat #: 5000006
Tyloxapol	Sigma	Cat #: 1704003
GW9662	Sigma	Cat #: M6191
Oleic acid	Sigma	Cat #: O1008
Sodium taurocholate	Sigma	Cat #: T4009
2-mono-oleoylglycerol	Sigma	Cat #: M7765
Cholesterol	Sigma	Cat #: C8667
L- α -lysophosphatidylcholine	Sigma	Cat #: L4129
BODIPY 558/568 C ₁₂	Invitrogen	Cat #: D3835
HEPES	Gibco	Cat #: 15630-080
Paraformaldehyde	Electron Microscopy Sciences	Cat #: 15710-5
Triton X-100	Acros Organics	Cat #: 327372500
Alexa Fluor 647 Phalloidin	Invitrogen	Cat #: A22287
ProLong Diamond Antifade Mountant with DAPI	Invitrogen	Cat #: P36971
Butyric acid, sodium salt	Acros Organics	Cat #: 263191000
DL-3-phenyllactic Acid-d3	Toronto Research Chemicals	Cat #: P335567
DL-3-phenyllactic acid	Toronto Research Chemicals	Cat #: H953713

Critical commercial assays

Total RNA Purification Kits	Norgen Biotek Corp.	Cat #: 37500
Oil Red O, Propylene Glycol staining kit	Newcomer Supply	Cat #: 12772
DNeasy PowerSoil Kit	Qiagen	Cat #: 47014
PureLink Genomic DNA Mini Kit	Invitrogen	Cat #: K182001
iScript gDNA Clear cDNA synthesis kit	BioRad	Cat #: 1725034
NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina	New England BioLabs Inc.	Cat #: E7760
Lipid Quantification Kit	Cell BioLabs, Inc.	Cat #: STA-617
Serum Triglyceride Quantification kit	Cell BioLabs, Inc.	Cat #: STA-396
Free Fatty Acid Assay Kit	Cell BioLabs, Inc.	Cat #: STA-618
Total Cholesterol Assay Kit	Cell BioLabs, Inc.	Cat #: STA-384
Bond Refine Polymer detection system	Leica Biosystems	Cat #: DS9800
Luciferase Assay System	Promega	Cat #: E1500

Deposited data

16S rRNA sequencing of distal small intestine contents	European Nucleotide Archive	ENA: PRJEB52997
RNA sequencing of distal small intestine epithelial cells	European Nucleotide Archive	ENA: PRJEB53157
Untargeted metabolomics of distal small intestine contents	Metabolomics Workbench	Metabolomics Workbench: ST002186

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental models: Cell lines		
Mouse: small intestinal cells from <i>H-2Kb-tsA58</i> transgenic mice	Whitehead et al. ⁶⁷	MSIE
Human: Caco-2 cells	ATCC	HTB-37
Experimental models: Organisms/strains		
Mouse: B6.Cg-Tg(Vil1-cre)997Gum/J	The Jackson Laboratory	JAX: 004586
Mouse: B6.129- <i>Pparg</i> ^{tm2Rev/J}	The Jackson Laboratory	JAX: 004584
Mouse: <i>Pparg</i> ^{fl/fl} <i>Villin</i> ^{cre/-}	Byndloss et al. ³⁸	N/A
Mouse: C57BL/6NCrl	Charles River Laboratory	CRL: 027
Mouse: Germ-free C57Bl/6N	In-house colony	N/A
Oligonucleotides		
Information regarding oligonucleotides used in this study is listed in Table S2		
Recombinant DNA		
PPRE x3-TK-Luc	Addgene	Plasmid #: 1015
Software and algorithms		
Excel	Microsoft	N/A
Prism 9 (Version 9.2.0)	Graphpad	https://www.graphpad.com/features
QIIME2 (version 2022.2)	N/A	https://qiime2.org/
STAR (v2.7.3a)	Dobin et al. ⁶⁸	https://github.com/alexdobin/STAR
featureCounts (v2.0.0)	Liao et al. ⁶⁹	https://subread.sourceforge.net/
DESeq2 (v.1.24.0)	Love et al. ⁷⁰	http://bioconductor.org/packages/release/bioc/html/DESeq2.html
WebGestaltR (v0.4.4)	N/A	https://www.webgestalt.org/
Progenesis QI v. 3.0	Non-linear Dynamics, Water Corporation	https://www.nonlinear.com/progenesis/qi/v3.0/download/
Metaboanalyst 5.0	N/A	https://www.metaboanalyst.ca/MetaboAnalyst/home.xhtml

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Mariana Byndloss (mariana.x.byndloss@vumc.org).

Materials availability

All unique/stable reagents generated in this study are available from the [lead contact](#) without restriction.

Data and code availability

16S rRNA sequencing data has been deposited in the European Nucleotide Archive under the accession number: ENA: PRJEB52997. RNA sequencing data has been deposited in the European Nucleotide Archive under the accession number: ENA: PRJEB53157. The mass spectrometry data are available in the Metabolomics Workbench public repository, with the data set identifier as Metabolomics Workbench: ST002186. All data sets are publicly available. This paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available from the [lead contact](#) upon request.

EXPERIMENTAL MODEL DETAILS

Animal Experiments

All experiments in this study were approved by the Institutional Animal Care and Use Committee (IACUC) at Vanderbilt University Medical Center.

Early-life antibiotic treatment using clinical doses has been linked to increased childhood body mass index in males.⁷¹ Additionally, the effects of diet-induced obesity are significantly greater in young male mice compared to females.⁷² As we focused our study on the synergistic effects of antibiotics and a high fat (HF) diet, male mice were used for this work. Male C57BL/6N mice (#027),

3-weeks-old, were obtained from Charles River Laboratory. Prior to experiments, fecal samples were collected from mice and initial *Lactobacillus* levels determined by homogenizing fecal content in 1 mL of sterile PBS, followed by serial dilutions of the samples which were plated on De Man, Rogosa, and Sharpe (MRS) agar (BD Difco #288210). After overnight incubation under anaerobic conditions, colony forming units (CFUs) were calculated to obtain *Lactobacillus* colonization levels. The initial weight of mice was also collected. Mice were then assigned to groups so that no significant differences in weight or *Lactobacillus* abundance between groups occurred before beginning treatments.

Female and male C57BL/6J *Pparg^{fl/fl}Villin^{cre/-}* and littermate *Pparg^{fl/fl}Villin^{-/-}* (control) mice were generated at Vanderbilt University Medical Center by mating *Pparg^{fl/fl}* mice (The Jackson Laboratory, #004584) with *Pparg^{fl/fl}Villin^{cre/-}* mice.³⁸ Genotype of offspring determined by PCR using protocol primers described by The Jackson Laboratory. Mice were weaned at 3 weeks of age and assigned to one of the treatment groups described below.

Animals were fed either a 60% fat diet (HF) (Research Diets Inc., #D12492) or a 10% fat control diet (LF) (Research Diets Inc., #D12450J) for 5 weeks. Groups of LF or HF diet mice were also given low doses of penicillin (Sigma Aldrich #P1382)(LDP) (6.67 mg/L¹⁰) in their drinking water throughout the experiment or clinical doses of penicillin (0.167 g/L⁷³) in their drinking water from days 0 – 5, 15 – 20, and 30 – 35. For long-term experiments, mice were fed either a LF or HF diet or a LF or HF diet with LDP for 5 weeks before being switched to a HF or LF diet alone for an additional 5 weeks. *Pparg^{fl/fl}Villin^{cre/-}* and littermate *Pparg^{fl/fl}Villin^{-/-}* mice were maintained on a HF diet or a HF diet and LDP treatment for 6 weeks as they required additional time to reach a similar weight as C57BL/6N mice on a HF diet (data not shown). In some experiments, drinking water was supplemented with 10 mM D-(+)-3-phenyllactic acid (Sigma-Aldrich #376906) for 5 weeks.

C57BL/6N germ-free mice (Taconic, B6NTac – GF) were bred in-house and moved into a positive pressure cage system at 3-weeks-old.

Over the course of the experiments, mice were weighed weekly. Food consumption of groups of mice was measured twice per week and normalized to the group or cage weight. In addition, fecal samples were collected weekly or bi-weekly, depending on the experiment, and plated onto MRS agar to measure *Lactobacillus* abundance. After 5, 8, or 10 weeks, mice were fasted for approximately 6 hours before fasting blood glucose was measured using a glucometer.

At the end of the experiment (5-weeks or 10-weeks after starting diet and antibiotic treatments), mice were humanely euthanized by CO₂ administration. Afterwards, serum, ileum (distal small intestine) content, ileum tissue (for epithelial isolation as described below), and samples for histopathology, IHC, and oil-red staining were collected. Liver and adipose tissue was flash frozen for RNA extraction. In addition, the epididymal (abdominal) fat from mice was removed and weighed.

Bacterial culture conditions

The bacterial strains used in this study are listed in Table S1. *Lactobacillus* strains were routinely grown anaerobically at 37 °C in MRS broth (BD Difco # 288130) or on MRS agar plates. *E. coli* Mt1B1 was grown aerobically in LB broth (BD Difco #244620) at 37 °C. *Muribaculum intestinale* was grown anaerobically at 37 °C in Anaerobic Akkermanisa Medium.⁷⁴ *Clostridium clostridioforme* and *Clostridium sporogenes* were grown anaerobically at 37 °C on Brain Heart Infusion (BHI) agar (BD Difco #21 1059) supplemented with 5% (v/v) defibrinated sheep blood (Thermo Scientific #R54016) and single colonies used to inoculate Supplemented Brain Heart Infusion (BHIS) medium.⁷⁵ *Streptococcus danielae* ERD01G was routinely grown anaerobically at 37 °C in Tryptic Soy Broth (TSB) (BD Difco #211825). *Staphylococcus xylosum* 33-ERD13C was routinely grown aerobically at 37 °C in LB broth (BD Difco #244620).

Mouse small intestinal epithelial cells

MSIE cells were generously provided by Dr. Fan Yang and were previously generated from the small intestine of 6- to 8-week-old C57BL/6 mice harboring a thermolabile mutation (tsA58) under the control of an IFN- γ -inducible H-2Kb promoter and a temperature-sensitive simian virus 40 (SV40) large T antigen.⁶⁷ Expression of the SV40 large T antigen was induced by culturing cells at 33°C with IFN- γ which allowed for cell proliferation. MSIE cells were routinely cultured in RPMI 1640 medium (Gibco #11875093) supplemented with 5% Fetal Bovine Serum (Gibco #16140071), 5 U/ml of mouse IFN- γ (R&D Systems #485-MI), 100 U/ml penicillin and streptomycin (Gibco #15140122), 5 μ g/ml insulin, 5 μ g/ml transferrin, and 5 ng/ml selenous acid (Sigma-Aldrich #I3146) at 33°C (permissive temperature) under 5% CO₂. Cells were passaged every 2 to 3 days. MSIE cells were seeded into 12-well plates (1 x 10⁵ cells/ well) or 6-well plates (6 x 10⁵ cells/well), incubated until confluent (usually 2 to 3 days), and then treated.

Caco-2 cell culture

Caco-2 cells (ATCC #HTB-37) were grown in Caco-2 cell culture media (Minimal Essential Media (MEM) (Gibco #11090081), 10% Fetal Bovine Serum (Gibco #16140071), 1% Glutamax (Gibco #35030061), 1% MEM Nonessential Amino Acids (Corning #11140050), and 1% Sodium Pyruvate (Gibco #11360070)) at 37°C and 5% CO₂. Cells were passaged every 3-4 days in T75 flasks according to ATCC protocols. For lipid secretion assays, 70 – 80% confluent T75 flasks were diluted, and approximately 1 x 10⁵ cells seeded onto 12-well transwell plates (Corning Costar #3493) containing collagen coated 0.4 μ m pore size inserts. Cells were differentiated for 14 days on the transwell inserts with the media being replaced every two days. For immunofluorescence experiments, 70% - 80% confluent T75 flasks were diluted, and then approximately 6 x 10⁵ cells seeded onto glass coverslips in 6-well plates.

Mouse enteroids

Enteroids were generated from 10–12-week-old female C57BL/6J mice. Mice were humanely euthanized, and the ileum removed, rinsed with ice-cold PBS, and then opened. Tissue was incubated in 0.5% bleach in PBS for 5 minutes before rinsing with 0.1 mg/mL Primocin (InvivoGen #ant-pm-05) in PBS (PBS/P). Crypts were then isolated from the tissue by gently rocking the ileum in chelation buffer (1M dithiothreitol and 20 mM EDTA/EGTA in PBS) at 4°C for 30 minutes. The supernatant was then discarded, and additional PBS/P was added before shaking the tissue vigorously. The supernatant was filtered through a 70 µm filter and then spun at 500 x g for 5 minutes. The cell pellet was resuspended in PBS/P and Matrigel (Corning #356234) (final concentration of Matrigel was approximately 80%). 30 µL of the mixture was aliquoted into a 6-well non-tissue culture treated plate (6 domes/ well) and incubated at 37°C until solidified. Media (Intesticult (Stemcell Technologies #06005), 10 µM Y-27632 (bio-technie | TOCRIS #1254), and Primocin) was then added and enteroids incubated for 2 days at 37°C, 5% CO₂. Y-27632 was removed after two days. Media was replaced every 2–3 days and enteroids were passaged every 7–10 days. Enteroids were passaged by dissociating organoids from Matrigel and adding TrypLE Express (Gibco #12604013).

Differentiated enteroids (10 – 14 days after passaging) were treated with either 10 µM Rosiglitazone (Sigma-Aldrich #R2408) or 5 mM PLA and incubated for 24 hours. Media was replenished during the incubation at 8 hours. Organoids were then dissociated out of the Matrigel, and RNA extracted from the cell pellets (Norgen BioTek Corp. #17200).

METHOD DETAILS

Body composition

Mice were fed a HF diet or exposed to a HF diet and LDP for 5 weeks. Then, dual-energy X-ray absorptiometry (DEXA) was performed before their sacrifice on anesthetized mice by using a Faxitron UltraFocus (Hologic). The instrument was calibrated for offset images, dark images, and flat-field images before the measurement by a method provided by the manufacturer.

Oil Red O Staining

Portions of liver were frozen in optimal cutting temperature (O.C.T.) compound (Fisher Scientific #23730571) and 5 µm thick sections were cut and mounted to slides. Slides were stored at -80 °C until staining. To stain lipids, slides were brought to room temperature and then placed in 10% neutral-buffered formalin solution for 10 minutes. The Newcomer Supply Tech Oil Red O, Propylene Glycol staining kit (Newcomer Supply #12772B) was used for visualization. Slides were then coverslipped with aqueous mounting medium. Scoring of blinded tissue sections were performed by a veterinary pathologist based on the criteria listed in [Table S4](#). Representative images were taken using a Leica DM750 microscope and a Leica ICC50W camera.

16s rRNA sequencing and data analysis

Content from the distal small intestine was collected at the end of the 5-week experiment and stored on ice. DNA was then extracted from the samples using the DNeasy PowerSoil Kit (Qiagen #47016) according to the manufacturer's instructions. Sequencing libraries were prepared by the University of California, San Diego (UCSD) Microbiome Core using protocols and primers published on the Earth Microbiome Project website and sequencing performed on Illumina NovaSeq 6000 according to the manufacturer's instructions. Other samples were sequenced by Seqmatic on Illumina MiSeq. Single end sequences were imported into QIIME2 (version 2022.2). Reads were trimmed to exclude regions that fell below a quality score of 25. Features tables for each data set were constructed and then merged following denoising. Data was rarefied based on minimum read quantity. Phylogenetic trees were then constructed, and taxonomy was assigned based on the Silva genes classifier. Pseudocounts added and ANCOM was performed to identify alterations in gut microbiota composition based on diet or administration of low-dose penicillin. Feature tables constructed for taxonomic levels 5(genera), 6(family), and 2(phylum). Pseudocounts added to each of the collapsed feature tables and ANCOMs performed at for each experimental variable as previously performed. Distance matrices (jaccard, bray-curtis, weighted, and unweighted) were exported from QIIME2 for further processing in R-studio. Alpha diversity metrics (Shannon, Faith_pd, and Observed features) were extracted from QIIME2 for downstream applications. PERMANOVAs performed for each distance matrix, as well as the creation of non-metric multidimensional scaling (nmds) plots based on each distance matrix.

L. murinus-specific qPCR

To measure the abundance of *L. murinus* in the small intestine, a protocol was adapted from Yildiz et al.⁷⁶ Briefly, a DNA (ng) and CFU correlation was generated using *L. murinus* (isolate collected from male C57BL/6N mice (#027), 3-weeks-old; species identified by 16S rRNA sequencing of the colony by Genewiz). *L. murinus* was cultured anaerobically in MRS broth overnight and then 1 mL of culture was serially diluted and plated on MRS agar for determination of CFU. A second 1 mL aliquot of *L. murinus* culture was lysed and genomic DNA extracted using the PureLink Genomic DNA Mini Kit (Invitrogen #K1820-01). CFU/ng of *L. murinus* was calculated in triplicate using the results from above.

DNA from content in the distal small intestine was extracted as described above using the DNeasy PowerSoil Kit. Primers specific for either the 16S rRNA gene of *Lactobacillus* species⁷⁷ or *L. murinus* ([Table S2](#)) were combined with iQ SYBR Green Supermix (Bio-Rad #1708882) and 20 ng of DNA from either the small intestine content or a standard curve of *L. murinus* gDNA. Quantitative PCR (qPCR) was performed using the following thermal cycling program: initial denaturation step at 95°C for 5 min, followed by 40 cycles of denaturation at 95°C for 30 s and annealing/extension at 60 °C for 60 s. Cq values obtained from the qPCR reactions were

compared to a *L. murinus* gDNA standard curve to determine the amount of *L. murinus* gDNA present in each sample. Then, CFU/ng of small intestine content DNA was calculated using the correlation calculated as described above.

Colonization of HF and LDP-treated mice with *L. murinus* Pen^R

To create a strain of *L. murinus* resistant to low doses of penicillin used in the mouse experiments described above, a single colony of *L. murinus* was inoculated into MRS broth containing 50 µg/L penicillin (Sigma Aldrich #P1382). After 24 hours, the culture was diluted 1:100 into MRS broth containing 100 µg/L penicillin. The culture was passaged as described into increasing concentrations of penicillin until *L. murinus* grew in MRS broth containing 6.67 mg/L penicillin. To isolate single colonies, passaged *L. murinus* was plated on MRS agar containing 6.67 mg/L penicillin. Single colonies were inoculated into fresh MRS broth and cultures used to make glycerol stocks.

To colonize mice with a *L. murinus* Pen^R (strain generation described below), male C57BL/6N mice (#027), 3-weeks-old, from Charles River Laboratory were inoculated by intragastric gavage (1×10^9 CFU in 0.1 mL MRS broth) with *L. murinus* Pen^R. At 48 hours post colonization, mice were switched onto a HF diet and given LDP in their drinking water. Mice were given *L. murinus* Pen^R by intragastric gavage (1×10^9 in 0.1 mL MRS broth) weekly for the 5 weeks of exposure to a HF diet and LDP.

Bacterial culture with bile salts and penicillin

For measuring the growth of bacteria upon exposure to penicillin, bile salts, lipids or a combination, bacterial species were cultured as follows: *L. murinus* was streaked onto MRS agar and single colonies used to start cultures in MRS broth. Overnight cultures were harvested by centrifugation at 4,000 x g for 5 minutes and then pellets were resuspended in 1 mL fresh media. Optical density at 600 nm (OD₆₀₀) was measured and cultures adjusted to an OD₆₀₀ = 0.1. Cultures were then diluted again to an OD₆₀₀ = 0.001. Stocks of penicillin (Sigma Aldrich #P1382) were prepared at a concentration of 250 mg/L in dH₂O and then batches at 10x the concentration tested in the assay were made in MRS broth (pH = 6.4). Bile salts (Sigma Aldrich #B8756) were dissolved in dH₂O at a concentration of 250 mM and then diluted in MRS broth (pH = 6.4) to a concentration of 0.9375 mM. Lipid mixture (Sigma Aldrich #L0288) was diluted to a concentration of 12.5% (v/v) in MRS broth. For measuring the growth of untreated bacterial species, 10 µl of adjusted bacterial culture was added to 90 µl of media in a 96 well plate. To measure the impact of penicillin on bacterial growth, 10 µl of 10x penicillin stocks were added to wells containing 10 µl of adjusted bacterial culture and 80 µl of media. The effect of bile salts on bacterial growth was determined by adding 80 µl of media containing 0.9375 mM bile salts to 10 µl of adjusted culture and 10 µl of plain media. The impact of lipids on bacterial growth was determined by adding 80 µl of media containing 12.5% lipids to 10 µl of adjusted culture and 10 µl of plain media. To measure the effect of the combination of penicillin and bile salts or penicillin and lipids on bacterial growth, 10 µl of adjusted culture, 10 µl of 10x penicillin stocks, and 80 µl of media containing 0.9375 mM bile salts, or 12.5% bile salts were combined. Plates were incubated anaerobically at 37 °C. OD₆₀₀ was measured every hour for 24 hours using the Epoch 2 plate reader (BioTek). *In vitro* growth assays were performed in triplicate with different colonies.

Gnotobiotic mouse experiments

C57BL/6N germ-free mice (from our in-house colony) were colonized with a defined microbiota community representative of a small intestine microbiota at 3 weeks of age. The defined microbiota included 1×10^9 CFU of the following bacteria: *Muribaculum intestinale*, *Clostridium clostridioforme*, *Clostridium sporogenes*, and *Streptococcus daniellae* ERD01G. *Ligilactobacillus murinus* (isolation and characterization described below) was added to the defined microbiota (1×10^{10} CFU) for a subset of mice. Germ-free mice were intragastrically inoculated with 0.1 mL of the defined microbiota and maintained on a chow diet for 5 days. Colonized mice were then given 1×10^8 CFU of *Escherichia coli* Mt1b1 and 48-hours later were switch onto a HF diet (Research Diets, Inc., D12492j). Mice were maintained on a HF diet for 5 weeks. Groups of mice colonized with *L. murinus* continued to be gavaged weekly with 1×10^{10} CFU of the bacterium and *L. murinus* was also supplied in their drinking water (1×10^8 CFU/mL).

Histopathology Analysis

Tissue from the distal small intestine was fixed in 10% neutral buffered formalin, paraffin-embedded and 5 µm thick sections were cut and mounted to slides and stained with hematoxylin and eosin. Scoring of blinded tissue sections were performed by a veterinary pathologist based on the criteria listed in Table S3. Representative images were taken using a Leica DM750 microscope and a Leica ICC50W camera.

Enterocyte isolation

The ileum (distal small intestine) was removed from mice and opened lengthwise before placing in cold phosphate buffered saline (PBS) (Gibco #14190144). Tubes were gently inverted four times. Tissue was then placed in a 15 mL conical centrifuge tube containing dissociation reagent #1 (30 mM EDTA, 1.5 mM DTT, pH 8 in PBS) and incubated for 20 minutes buried in ice. Then, tissue was transferred to a new 15 mL conical centrifuge tube containing pre-warmed (37°C) dissociation reagent #2 (30 mM EDTA, pH 8 in PBS) and samples were incubated at 37°C for 10 min. After incubation, tubes containing the samples were shaken vigorously for 30 seconds to release epithelium from the basement membrane. Cell suspension was transferred to a new 15 mL conical centrifuge tube, leaving any remnant tissue, and cells were pelleted by centrifugation at 800 x g for 5 minutes at 4°C. Supernatant was removed and the cell pellet was flash frozen in liquid nitrogen for subsequent RNA extraction or triglyceride measurements.

RNA extraction and quantitative real-time PCR

Liver tissue and enterocyte cell pellets were homogenized using a FastPrep-24 and RNA isolated by the TRIzol method (Invitrogen #15596018) following the manufacturer's protocol. Adipose tissue was homogenized using a Fastprep-24 and RNA isolated using QIAzol Lysis Reagent (Qiagen #79306) according to the manufacturer's protocol. RNA was reverse transcribed using iScript gDNA Clear cDNA Synthesis Kit (Bio-Rad #1725035). Quantitative real-time PCR was performed using iQ SYBR Green Supermix (Bio-Rad #1708882) and the appropriate primer sets (Table S2). Relative gene expression was calculated using *Act2b* as the house-keeping gene.

RNA-sequencing

RNA was extracted from enterocyte cell pellets as described above and samples sequenced by the Vanderbilt Technologies for Advanced Genomics (VANTAGE). VANTAGE performed an mRNA enrichment and then prepared a cDNA library using the stranded mRNA (polyA-selected) library preparation kit (New England BioLabs #E7760). Sequencing was then performed at paired-end 150 bp on the Illumina NovaSeq 6000 targeting an average of 25 M reads per sample. Results were then analyzed by Vanderbilt Technologies for Advanced Genomics Analysis and Research Design (VANGARD) as follows. Adapters were trimmed by Cutadapt (v2.10). After trimming, reads were mapped to the mouse reference genome GRCm38.p6 using STAR (v2.7.3a)⁶⁸ and quantified by featureCounts (v2.0.0).⁶⁹ DESeq2 (v.1.24.0)⁷⁰ was used to detect differential expression between two groups. WebGestaltR (v0.4.4) was used to perform functional enrichment analysis against Gene Ontology and KEGG. Gene Ontology analysis was performed using the online tool David Bioinformatics Resources (<https://david.ncifcrf.gov/>).

Lipid Measurements

After 5 weeks of exposure to a HF diet or a HF diet and LDP, mice were fasted overnight and then humanely euthanized by CO₂. Plasma was collected and then the amount of neutral lipids (Cell Biolabs STA-617), triglycerides (Cell Biolabs STA-396), free fatty acids (Cell Biolabs STA-618), or cholesterol (Cell Biolabs STA-390) were measured using the appropriate assay kit. Enterocytes from fasted mice were collected as described above and triglycerides measured in the cell lysate. The concentration of triglycerides was normalized to protein abundance in the cell lysate as determined by a standard Bradford assay (Bio-Rad #5000006).

Non-invasive Measurement of Intestinal Fat Absorption

Measurements of dietary fat absorption were based off the protocol by Jandacek et al.⁷⁸ C57BL/6N were given a HF diet or a HF diet and LDP for 5 weeks as described above. After the 5 weeks of treatment, mice were placed in a clean cage and switched to a diet containing sucrose polybehenate (Research Diets, Inc., #D12492 with 5% of the fat replaced with sucrose polybehenate). The mice were moved again to a clean cage after 24 hours and maintained on the diet containing sucrose polybehenate. On the third day of feeding with sucrose polybehenate, fecal pellets were collected from the mice before placing animals in a clean cage. After 4 days on the diet, fecal pellets were collected again.

Fecal samples were saponified with methanolic sodium hydroxide to obtain free fatty acids which were then methylated using 14% BF₃ methanol to produce fatty acid methyl esters. Hexane was then used to extract the fatty acid methyl esters. The fatty acids were detected using gas chromatography and compared to a set of standards and an internal standard. The percentage of fat absorption was determined based on the ratio of total fatty acids to behenic acid in the diet and in the feces.

Oral fat tolerance test

Mice were given a HF diet or a HF diet and LDP for 5 weeks before being fasted for 6 hours. Then, mice were injected intraperitoneally with 0.5 g/kg tyloxapol (Sigma Aldrich #1704003; dissolved in sterile PBS) 30 minutes before they were given 200 μ l of olive oil via intragastric gavage. Blood samples were collected from the tail 2 hours after olive oil gavage. Samples were then spun at 1000 x g for 20 minutes at 4 °C and the supernatant collected and stored at -80 °C. Triglycerides were measured in the serum as described below.

PPAR- γ Immunostaining

Pieces of the distal small intestine were fixed in 10% neutral-buffered formalin, paraffin-embedded and 5 μ m thick sections were cut and mounted to slides. Slides were placed on the Leica Bond-Rx IHC Stainer. All steps besides dehydration, clearing and coverslipping were performed on the Bond-Rx. Slides were deparaffinized. Heat induced antigen retrieval was performed on the Bond Max using their Epitope Retrieval 2 solution (Leica Biosystems #AR9640) for 20 minutes. Slides were placed in a protein block (Agilent DAKO # x0909) for 10 minutes. Slides were then placed in a PowerVision Block (Leica Biosystems #PV6122) for 60 minutes. The sections were incubated with anti-PPAR γ (Santa Cruz Biotechnology #sc-7273), diluted 1:100, for 30 minutes. The Bond Refine Polymer detection system (Leica Biosystems #DS9800) was used for visualization. Slides were the dehydrated, cleared and coverslipped. Scoring of blinded tissue sections were performed by a veterinary pathologist based on the criteria listed in Table S5. Representative images were taken using a Leica DM750 microscope and a Leica ICC50W camera.

Lipid secretion assay in Caco-2 cells

Differentiated Caco-2 wells were treated with either 3 μ M GW9662 (Sigma-Aldrich #M6191), live *L. murinus* (prepared as described above), both 3 μ M GW9662 (Sigma-Aldrich #M6191) and live *L. murinus*, or 5 mM D-(+)-3-phenyllactic acid (Sigma-Aldrich #376906)

for 16 hours. Lipid micelles were prepared as previously described^{22,23,79} in serum-free Caco-2 cell culture media. Micelles, containing 0.6 mM oleic acid (Sigma-Aldrich #O1008), 2 mM sodium taurocholate (Sigma-Aldrich #86339), 0.2 mM 2-mono-oleoylglycerol (Sigma-Aldrich #M7765), 0.05 mM cholesterol (Sigma-Aldrich #C8667), 0.2 mM L- α -lysophosphatidylcholine (Sigma-Aldrich #L4129) and 0.02 mM BODIPY C₁₂ fatty acid (Invitrogen #D3835), were added to the apical compartment of cells. After a 10-minute incubation, media containing lipid micelles was removed and replaced with serum-free Caco-2 cell culture media. Four and six hours after addition of lipid micelles, 150 μ L from the basal compartment was removed and 50 μ L added in triplicate to wells of black 96-well plate. Fluorescence was measured (excitation at 548 nm and emission at 578 nm) (BioTek).

PPAR- γ fluorescent staining

MSIE or Caco-2 cells were seeded onto coverslips as described above and incubated for 48 hours. Then, cells were mock-treated or treated with live *L. murinus* (prepared as described above), *L. murinus* or *E. coli* Mt1B1 supernatant (prepared as described above), or 5 mM D-(+)-3-phenyllactic acid (Sigma-Aldrich #376906) for 16 hours. All conditions were buffered with 20 mM HEPES (Gibco #15630080). After treatment, media was removed, and cells were fixed with 4% paraformaldehyde (Electron Microscopy Sciences #15710-S) for 15 minutes at 37 °C. Cells were then washed 3 times for 5 minutes in PBS (Gibco #14190144). Subsequently, cells were treated with 0.1% Triton X-100 (Acros Organics #AC327371000) for 15 minutes and then washed again 3 times for 5 minutes with PBS (Gibco #14190144). Cells were blocked in 5% BSA (Alfa Aesar #J64655) for one hour before being incubated with anti-PPAR- γ (Santa Cruz Biotechnology #sc-7273) (Diluted 1:50) for one hour at room temperature. After incubation with primary antibody, cells were washed 3 times for 5 minutes in PBS (Gibco #14190144) and then goat anti-mouse IgG Alexa Fluor 488 (Invitrogen #A-11001) (diluted 1:1000) was added for 1 hour at room temperature. Coverslips were washed in PBS (Gibco #14190144) and then Alexa Fluor 546-Phalloidin (Thermo Fisher #A22283) or Alexa Fluor 647-Phalloidin (Thermo Fisher #A22287) was added for 20 minutes. Afterwards, coverslips were washed in PBS (Gibco #14190144) and then mounted on glass slides in ProLong Diamond Antifade Mountant with DAPI (Molecular Probes #P36962). Imaging was performed on a Zeiss LSM880 Airyscan Confocal Microscope. Representative images shown in paper were collected using a Nikon A1 Microscope equipped with 405, 488, 561, and 645 nm LASERS and a Plan Fluor 40X/ 1.30 NA immersion objective. For imaging in all microscope modalities, imaging acquisition parameters were matched between samples during image acquisition. Images were contrast enhanced and cropped using FIJI/ImageJ software (NIH). Abundance of nuclear PPAR- γ was calculated using FIJI/ImageJ measurement tools and dimensions of DAPI stained nuclei.

PPAR Reporter Assay

Eighty to ninety percent confluent Caco-2 cells were seeded into 96 well plates at a concentration of approximately 1×10^4 cells/well. Forty-eight hours post seeding, wells containing Caco-2 cells were transfected with 150 ng of PPRE x3-TK-Luc⁸⁰ using Lipofectamine 3000 (Thermo Fisher L3000008) according to manufacturer's instructions. After overnight incubation, transfected cells were mock-treated or treated with 8 mM sodium butyrate (Acros Organics #263191000), or live *L. murinus* for 5 hours. Media was then removed, and cells washed with 100 μ L PBS. Cells were then lysed and luciferase assay substrate (Promega #E1500) added according to manufacturer's instructions. Luciferase was then measured using plate reader (BioTek).

Co-culture of bacterial strains and intestinal epithelial cells

For *in vitro* experiments, strains were grown overnight as follows: *L. murinus* was grown anaerobically in MRS broth and *E. coli* Mt1B1 and *S. xylosum* 33-ERD13C were grown aerobically in LB broth. Overnight cultures were harvested at 4000 x g for 5 minutes. *L. murinus* was then adjusted to an OD = 1 in Caco-2 cell culture media, *E. coli* Mt1B1 was adjusted to an OD = 0.025, and *S. xylosum* 33-ERD13C was adjusted to an OD = 0.05. Strains were adjusted to different starting concentrations due to the ability of *E. coli* and *S. xylosum* 33-ERD13C to grow in the cell culture media. Bacterial strains were added to Caco-2 or MSIE cells for co-culture experiments. Similar bacterial burdens between strains were observed after 16 hours of co-culture. To prepare *Lactobacillus* and *E. coli* Mt1B1 supernatants, Caco-2 media was inoculated with a single colony of *L. murinus* and grown aerobically (37 °C, 5% CO₂) for 24 hours. Culture was then harvested at 4000 x g for 5 minutes and the supernatant filter sterilized (0.2 μ m pore size). The volume of the supernatant used was normalized by the OD of the bacterial culture when two different strains were used. Both live cultures and supernatants were buffered using 20 mM HEPES. MSIE cells were incubated with live bacterial cultures for 16 hours and then either RNA was extracted from the cells (Norgen BioTek Corp. #17200), or the cells were stained for PPAR- γ (protocol described below). Caco-2 cells were incubated with either live bacteria or their supernatant for 16 hours before cells were stained for PPAR- γ (protocol described below).

Untargeted metabolomics

Optima grade LC-MS solvents for the mass spectrometry analyses were purchased from Thermo Fisher Scientific, USA (Thermo Fisher Scientific, Waltham, MA). Individual isotopically labeled standards were purchased from Cambridge Isotope Laboratories (Tewksbury, MA) and CDN Isotopes (Pointe-Claire, Quebec, Canada).

Metabolomics sample preparation

Frozen mouse intestinal content samples (n=20, 5 biological replicates for each sample group) were lysed in 500 μ L ice-cold lysis buffer (1:1:2, v:v:v, acetonitrile: methanol: ammonium bicarbonate 0.1M - pH 8.0) and sonicated individually using a probe tip sonicator at 50% power (10 pulses). The lysis buffer contained isotopically labeled standards (n=2) to determine sample process

variability (see Table S6). Homogenized samples were normalized by weight to the smallest amount of tissue sample such that each sample contained an equal amount of tissue. Proteins were precipitated from individual samples by addition of 800 μL of ice-cold methanol followed by overnight incubation at -80°C . Precipitated proteins were pelleted by centrifugation (15,000 rpm, 15 min) and metabolite extracts were dried down *in vacuo* and stored at -80°C . Individual samples were reconstituted in 100 μL of reverse phase liquid chromatography reconstitution buffer (acetonitrile/water with 0.1% formic acid, 3:97, v/v) containing isotopically labeled standards ($n=2$) to assess instrument variability (see Table S6). A pooled quality control (QC) sample was prepared by pooling equal volumes (10 μL) from each individual sample following reconstitution.

Metabolomics based reverse phase liquid chromatography high resolution tandem mass spectrometry analysis

Prepared samples were analyzed by RPLC-HRMS/MS in the Vanderbilt Center for Innovative Technology (CIT) using a modified version of a reversed phase chromatography negative ionization method.⁸¹ Metabolites were separated on a Thermo Fisher Scientific (Waltham, MA) Hypersil Gold C18 column (100 x 2.1 mm, 1.9 μm particle size) using water/acetonitrile gradient with formic acid (0.1%) added to both mobile phases. Samples were analyzed on a high-resolution Thermo Fisher Scientific Q-Exactive HF hybrid quadrupole-Orbitrap mass spectrometer (Bremen, Germany) equipped with a Thermo Fisher Scientific Vanquish UHPLC binary system and autosampler controlled by Xcalibur 4.4 software (Waltham, MA). Liquid chromatography was performed at 250 $\mu\text{L min}^{-1}$ using solvent A (0.1% formic acid in water) and solvent B (0.1% formic acid in 80:20 acetonitrile:water) with a gradient length of 30 min as previously described in Popay et al.⁸¹ Solvents and column were held at 45°C .

Mass spectrometry (full MS) analyses (5 μL injection volume) were acquired in negative ion mode at maximum resolution of 120,000 and scanned from m/z 70 to m/z 1050 with a scan rate of 3.5 Hz, automatic gain control (AGC) target of 1×10^6 , and maximum ion injection time of 100 ms. MS/MS spectra were collected at 15,000 resolution, AGC target of 2×10^5 ions, and maximum ion injection time of 100 ms. Pooled QC samples were used for column conditioning (10x pooled QC sample prior to individual sample analysis and 2x pooled QC sample was injected every five samples), retention time alignment and to assess mass spectrometry instrument reproducibility (intra-batch) throughout the sample set. A total of 20 samples (randomized injection 1x) were analyzed by RPLC-HRMS/MS analysis after conditioning the analytical platform with pooled QC samples. Quality assurance (QA) practices were applied to assess the analytical method performance. Further, a system suitability sample (LCMS QC reference material, Waters Corporation, Milford, MA) was used to verify the stability of retention times, peak shapes and areas before and after sample batch analysis.

Metabolomics data processing and statistical analysis

Spectral features (retention time, m/z pairs) were extracted from the RPLC-HRMS/MS data using Progenesis Q1 v. 3.0 (Non-linear Dynamics, Waters Corporation, Milford, MA). Briefly, the data from five ($n=5$) biological replicates from each sample type were imported, processed, normalized (integration) and reviewed. All MS and MS/MS sample runs were aligned against a pooled QC sample reference run. Spectral features (retention time, m/z pairs) were further de-adducted and de-isotoped to generated compounds or metabolites. Data was further curated by applying QA practices to the data. Specifically, compounds or metabolites with spectral features with $>25\%$ coefficient of variation (CV) in the pooled QC samples were removed (total: 2270 metabolites). Data were normalized to all compounds using Progenesis Q1.

Principal component Analysis (PCA) was used to identify and remove outliers in the data set (see Figure S8A). Heatmap analysis was performed in Metaboanalyst 5.0, data was pareto scaled and log transformed using Pearson distance and average clustering (see Figure S8B) to illustrate differences based on metabolite abundances across the experimental groups. Sample process and instrument variability were also assessed (see Table S6) to determine sample and batch acceptance. Briefly, QA metrics for sample process variability and instrument variability are $\leq 25\%$ CV and $\leq 10\%$ CV, respectively. In these studies, no samples were identified as outliers. Statistical analyses were performed in Progenesis Q1 using variance stabilized measurements achieved through log normalization to calculate p-values by one-way analysis of variance (ANOVA) test. Significantly changed metabolites were chosen with the criteria $p\text{-value} < 0.05$ and $|\text{fold change}| > 2$.

Tentative (Level 3, L3) and putative annotations (Level 2, L2) were determined within Progenesis Q1 software using accurate mass measurements (<5 ppm error), isotope distribution similarity, retention time matching and fragmentation spectrum matching using database searches against Human Metabolome Database (HMDB),⁸² METLIN,⁸³ the National Institute of Standards and Technology database⁸⁴ and an in-house library. Annotations (Level 1-3)⁸⁵ were determined for significant metabolites by searching the above libraries or databases. The metabolite annotations of significantly different compounds (pairwise comparisons between HF diet and HF diet + LDP-treated mice) are represented in Table S7. The MS data are available in the metabolomics Workbench public repository, with the data set identifier as ST002186.

Preparation of defined microbiota supernatants

To measure PLA in supernatants from defined microbiota cultures, fecal samples were collected from gnotobiotic mice colonized with the defined microbiota described above (including or lacking *L. murinus*) prior to colonization with *E. coli* Mt1B1. Fecal samples were resuspended at a concentration of 1 g feces to 25 mL PBS and combined in a 1:1 ratio with 50% sterile glycerol. Samples were then stored at -80 . 1 mL of the defined microbiota fecal sample (either with or without *L. murinus*) was used to inoculate 10 mL of Anaerobic Akkermanisa Medium. Cultures were grown anaerobically for 48 hours. After 48 hours, cultures were harvested at $4000 \times g$ for 10 minutes. The resulting pellets were resuspended in MRS broth and separated into 3 5 mL aliquots of Caco-2 media (previously incubated anaerobically for 48 hours). Cultures were then incubated for 16 hours before being harvested at $4000 \times g$ for 10 minutes. The supernatant was filter sterilized and PLA measured as described below.

Phenylactic Acid Measurements

Extraction and normalization

Intestinal contents were weighed, diluted to a final density of 40 mg/mL in water/acetonitrile (3:2) buffered with 250 mM sodium phosphate pH 2.5 and homogenized/extracted using ceramic beads (2.8 mm dia.) and a *Bead Mill 4* homogenizer (Thermo Fisher; Waltham, MA). Insoluble debris was removed by centrifugation (10,000 \times g, 30 min, 5 °C); the supernatants were transferred to clean Eppendorf tubes and stored at -20 °C until the day of analysis.

LC-HRMS Analysis

3-Phenylactic acid (PLA) and a deuterium-labeled internal standard (PLA- d_3) were derivatized with the reagent dansyl hydrazine and the carboxyl activating agent 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) to their corresponding dansyl hydrazone derivatives. Due to the limited stability of EDC under acidic conditions, stock solutions of EDC were made up fresh in water and used immediately. Briefly, 25 μ L aliquots of thawed extracts were spiked with 2 nmol PLA- d_3 , diluted 1:3 with homogenization buffer, and derivatized at room temperature with dansyl hydrazine (25 μ L \times 50 mg/mL) and EDC (25 μ L \times 150 mg/mL). After one hour at room temperature, 25 μ L methanol/water (1:1) containing 500 mM glucose was added to quench the excess reagents. Following centrifugation (10,000 \times g, 30 min, 5 °C), quenched reaction mixtures were transferred to 2-mL autosampler vials equipped with low-volume polypropylene inserts and Teflon-lined rubber septa. The sample injection volume was 10 μ L. PLA calibration standards were prepared in water and derivatized in the same manner. LC-HRMS analysis was performed using a Thermo Q Exactive HF hybrid quadrupole/orbitrap high resolution mass spectrometer interfaced to a Vanquish Horizon HPLC system (Thermo Fisher). The mass spectrometer was operated in positive ion mode, with targeted selected ion monitoring (t-SIM) detection of specified precursor ions at a resolving power of 30,000, an isolation window of 2.0 m/z , and the following HESI source parameters: spray voltage 4 kV; capillary temperature 300 °C; HESI temperature 100 °C; S-lens RF level 50; N₂ sheath gas 40; N₂ auxiliary gas 10; sweep gas 2.0; in-source CID off. Extracted ion chromatograms were constructed for PLA and PLA- d_3 with the following exact masses and a mass tolerance of \pm 5 ppm: PLA: [M+H]⁺ 414.1482; PLA- d_3 : [M+H]⁺ 417.1670. Data acquisition and quantitative spectral analysis were done using Thermo Xcalibur version 4.1.31.9 and Thermo LCQuan version 2.7, respectively. Calibration curves were constructed by plotting peak areas against analyte concentrations for a series of eleven calibration standards, ranging from 0.010 to 1000 μ M. A weighting factor of $1/C^2$ was applied in the linear least-squares regression analysis to maintain homogeneity of variance across the concentration range. An Acquity BEH C18 reverse phase analytical column (2.1 \times 100 mm, 1.7 μ m, Waters, Milford, MA) was used for all chromatographic separations. Mobile phases were made up of 0.2 % HCOOH in (A) water and in (B) CH₃CN. Gradient conditions were as follows: 0–1.0 min, B = 5 %; 1–6 min, B = 5–95 %; 6–8 min, B = 95 %; 8–8.5 min, B = 95–5 %; 8.5–12 min, B = 5 %. The flow rate was maintained at 300 μ L/min, and the total chromatographic run time was 12 min. A software-controlled divert valve was used to transfer the LC eluent from 0 to 4.5 min and from 6.5 to 12 min of each chromatographic cycle to waste.

QUANTIFICATION AND STATISTICAL ANALYSIS

Fold changes of ratios (mRNA relative expression, PPRE activity, BODIPY C₁₂ secretion) as well as percent abdominal fat, fasting glucose, PLA concentration (relative abundance and absolute concentration), and bacterial abundance were transformed logarithmically prior to statistical analysis. For quantification of PPAR- γ nuclear intensity, the integrated density of PPAR- γ fluorescent signal within the DAPI stained nucleus was measured. The integrated density was then transformed logarithmically prior to statistical analysis. An unpaired Student's *t* test was used on the transformed data to determine whether differences between groups were statistically significant ($p < 0.05$). Comparisons between more than 3 groups were determined using a one-way ANOVA followed by Tukey's or Šidák's multiple comparison test. When comparisons between multiple groups and containing two independent variables were assessed, statistically significant differences between groups were determined by two-way ANOVA followed by Tukey's or Šidák's multiple comparison test. Significance of differences in histopathology, PPAR- γ immunohistochemistry, and oil red O staining were determined by a two-tailed non-parametric test (Mann-Whitney).