



Consensus

The Youth Fitness International Test (YFIT) battery for monitoring and surveillance among children and adolescents: A modified Delphi consensus project with 169 experts from 50 countries and territories

Francisco B. Ortega^{a,b,*}, Kai Zhang^{c,d,1}, Cristina Cadenas-Sanchez^{a,e,f}, Mark S. Tremblay^{c,g}, Gregor Jurak^h, Grant R. Tomkinsonⁱ, Jonatan R. Ruiz^{a,j}, Katja Keller^k, Christine Delisle Nyström^l, Jennifer Sacheck^m, Russell Pateⁿ, Kathryn L. Weston^o, Tetsuhiro Kidokoro^{i,p}, Eric Poon^q, Lucy-Joy M. Wachira^r, Ronald Ssenyonga^s, Thayse Natacha Q.F. Gomes^{t,u}, Carlos Cristi-Montero^v, Brooklyn J. Fraser^{i,w}, Claudia Niessner^k, Vincent O. Onywere^x, Yang Liu^y, Li-Lin Liang^z, Stephanie A. Prince^{aa,bb}, David R. Lubans^{b,cc}, Justin J. Lang^{c,i,aa,bb,*},
the Delphi Fitness Expert Group²

^a Department of Physical Education and Sports, Faculty of Sport Sciences, Sport and Health University Research Institute (iMUDS), University of Granada; and CIBEROBN Physiopathology of Obesity and Nutrition; Granada ES18071, Spain

^b Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä FI40014, Finland

^c Healthy Activity Living and Obesity (HALO) research group, Children's Hospital of Eastern Ontario Research Institute, Ottawa, ON K1H 8L1, Canada

^d School of Human Kinetics, Faculty of Health Sciences, University of Ottawa, Ottawa, ON K1N 6N5, Canada

^e Department of Cardiology, Stanford University, Stanford, CA 94305, USA

^f Veterans Affairs Palo Alto Health Care System, Palo Alto, CA 94304, USA

^g Department of Pediatrics, University of Ottawa, Ottawa, ON K1H 8L1, Canada

^h University of Ljubljana, Faculty of Sport, 1000 Ljubljana, Slovenia

ⁱ Alliance for Research in Exercise, Nutrition and Activity (ARENA), Allied Health and Human Performance, University of South Australia, Adelaide, SA 5001, Australia

^j Instituto de Investigación Biosanitaria, ibs.Granada, Granada ES18012, Spain

^k Institute for Sport and Sport Science, Karlsruhe Institute of Technology, Karlsruhe 71631, Germany

^l Department of Medicine Huddinge, Karolinska Institute 141 83, Huddinge, Sweden

^m Department of Exercise and Nutrition Sciences, Milken Institute School of Public Health, The George Washington University, Washington, DC 20052, USA

ⁿ Department of Exercise Science, Arnold School of Public Health, University of South Carolina, Columbia, SC 29208, USA

^o Department of Psychological Sciences and Health, University of Strathclyde, Glasgow G1 1QE, UK

^p Nippon Sport Science University, Faculty of Sport Science, Tokyo 158-8508, Japan

^q Department of Sports Science and Physical Education, The Chinese University of Hong Kong, Hong Kong, China

^r Department of Physical Education, Exercise and Sports Science, School of Health Sciences, Kenyatta University, P.O Box 43844-00100, Nairobi, Kenya

^s Department of Epidemiology and Biostatistics, College of Health Sciences, Makerere University, P. O Box 7072, Kampala, Uganda

^t Department of Physical Education, Federal University of Sergipe, São Cristóvão 49107-230, Brazil

^u Department of Physical Education and Sport Sciences, Health Research Institute, Physical Activity for Health Research cluster, University of Limerick, Limerick V94 T9PX, Ireland

^v IRyS group, Physical Education School, Pontificia Universidad Católica de Valparaíso, Valparaíso 2374631, Chile

^w Menzies Institute for Medical Research, University of Tasmania, Hobart, Tasmania 7000, Australia

^x KCA University, Division of Research, Innovation and Outreach, P. O. Box 56808 - 00200, Nairobi, Kenya

^y School of Physical Education, Shanghai University of Sport, Shanghai 200438, China

^z Institute of Public Health, College of Medicine, National Yang Ming Chiao Tung University

Peer review under responsibility of Shanghai University of Sport.

* Corresponding author. Department of Physical Education and Sports, Faculty of Sport Sciences, Sport and Health University Research Institute (iMUDS), University of Granada; and CIBEROBN Physiopathology of Obesity and Nutrition; Granada, ES18071, Spain.

E-mail addresses: ortegaf@ugr.es (F.B. Ortega), justin.lang@phac-aspc.gc.ca (J.J. Lang).

¹ Both authors equally contributed to this study.

² The list of all fitness experts that completed the Delphi survey are provided as Appendix at the end of the main text.

<https://doi.org/10.1016/j.jshs.2024.101012>

^{aa} Centre for Surveillance and Applied Research, Public Health Agency of Canada, Ottawa, ON K1A 0K9, Canada
^{bb} School of Epidemiology and Public Health, Faculty of Medicine, University of Ottawa, Ottawa, ON K1H 8M5, Canada
^{cc} Centre for Active Living and Learning, University of Newcastle, Newcastle, NSW 2308, Australia

Received 3 August 2024; revised 11 September 2024; accepted 26 September 2024

Available online xxx

2095-2546/© 2024 Published by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract

Background: Physical fitness in childhood and adolescence is associated with a variety of health outcomes and is a powerful marker of current and future health. However, inconsistencies in tests and protocols limit international monitoring and surveillance. The objective of the study was to seek international consensus on a proposed, evidence-informed, Youth Fitness International Test (YFIT) battery and protocols for health monitoring and surveillance in children and adolescents aged 6–18 years.

Methods: We conducted an international modified Delphi study to evaluate the level of agreement with a proposed, evidence-based, international fitness battery (YFIT) of core health-related fitness tests and protocols to be used worldwide in 6- to 18-year-olds. This proposal was based on previous European and North American projects that systematically reviewed the existing evidence to identify the most valid, reliable, health-related, safe, and feasible fitness tests to be used in children and adolescents aged 6–18 years. We designed a single-panel modified Delphi study and invited 216 experts from all around the world to answer this Delphi survey, of whom one-third are from low-to-middle income countries and one-third are women. Four experts were involved in the piloting of the survey and did not participate in the main Delphi study to avoid bias. We pre-defined an agreement of $\geq 80\%$ among the expert participants to achieve consensus.

Results: We obtained a high response rate (78 %) with a total of 169 fitness experts from 50 countries and territories, including 63 women and 61 experts from low- or middle-income countries/territories. Consensus ($>85\%$ agreement) was achieved for all proposed tests and protocols, supporting the YFIT battery, which includes weight and height (to compute body mass index as a proxy of body size/composition), the 20-m shuttle run (cardiorespiratory fitness), handgrip strength, and standing long jump (muscular fitness).

Conclusion: This study contributes to standardizing fitness tests and protocols used for research, monitoring, and surveillance across the world, which will allow for future data pooling and the development of international and regional sex- and age-specific reference values, health-related cut-points, and a global picture of fitness among children and adolescents.

Keywords: Fitness; Experts; Delphi; Protocols; Youth Fitness International Test

1. Introduction

Since the early 2000s, a wealth of studies have provided clear and consistent evidence supporting physical fitness (a set of attributes that people have or achieve that relates to their ability to perform physical activity) in childhood and adolescence as a powerful marker of current and future health.^{1–4} The physical fitness components that have demonstrated the strongest health-related benefits are generally cardiorespiratory and muscular fitness. For instance, poor cardiorespiratory and/or muscular fitness in young people predicts future psychiatric diseases and suicide,^{5,6} cardiovascular disease risk factors,^{1–3} cardiovascular disease morbidity and mortality,^{5,7–13} and all-cause disability and mortality.^{5,7,9,10,13} Although mostly based on cross-sectional evidence, it has also been shown that better cardiorespiratory and muscular fitness in childhood and adolescence is associated with better mental health, cognitive and academic performance, higher neuroelectric activity, and larger gray matter and total brain volumes.^{6,14–24}

It is therefore clear that assessing fitness in childhood and adolescence can provide valuable information about the current and future health status of the population. The added benefit of including fitness in health surveillance and monitoring is that it is non-invasive, cost-effective, and relatively simple (requires modest tester training and expertise). In fact, there are examples of national fitness monitoring and

surveillance systems for children and adolescents in Europe proving its feasibility, as recently described by the European Union (EU) funded FitBack project,²⁵ and in other regions of the world.²⁶ Examples of these countries and territories include Finland; France; Hungary; Portugal; Serbia; Slovenia; Scotland, UK; Japan; China; Republic of Korea; and Brazil. Some of them include optional or compulsory fitness assessment as part of school curriculum, and others conduct representative sampling in the form of national surveys (e.g., Canadian Health Measures Survey (CHMS)). An example of the usefulness of these fitness surveillance systems is the fact that they have been able to evaluate the impact of the coronavirus disease 2019 (COVID-19) pandemic on health-related fitness.^{27–29} These systems also have the capacity to identify regions, municipalities or neighborhoods, as well as sub-populations (e.g., by income status, immigrant status, cultural/racial background) at increased risk of poor health as indicated by their low relative fitness levels, which is information of clear importance for targeted public health policy. Moreover, some fitness monitoring systems, such as those in Finland and Slovenia, have linked fitness data to national health information systems. These systems have found that the most feasible way of conducting fitness testing among children and adolescents is through schools, during physical education sessions.²⁵

Despite the great potential of fitness assessment for monitoring and surveillance, the major limitation has been the

inconsistency in fitness tests and protocols used across studies and national testing systems, which limits comparability. These inconsistencies hamper the interpretation of the fitness test results when comparing to representative normative values or sex- and age-specific health-related cut-points. Furthermore, for each fitness test there are a number of measurement protocols available, which also negatively impacts comparability, data pooling, and interpretation. Consequently, there is an important need to identify a core set of fitness tests for health monitoring and surveillance internationally, and to standardize the protocols for each test.

Therefore, the objective of the current study was to seek international consensus on a proposed, evidence-informed,^{30,31} Youth Fitness International Test (YFIT) battery and protocols for health monitoring and surveillance in children and adolescents aged 6–18 years. To achieve this objective, we conducted an international Delphi study with a large and diverse expert group to investigate the level of agreement with the proposed tests and protocols. The Delphi approach is a systematic expert consensus procedure for gathering the most reliable opinions from a group (ideally large and diverse) of independent experts who cannot meet in real-time for logistic or economic reasons and with the ultimate goal of attaining consensus.³² This Delphi study is directly related to 3 of the 10 international priorities (i.e., international surveys using common measures; develop universal health-related fitness cut-points; develop international field-based fitness test) for physical fitness research and surveillance among children and adolescents, as recently identified by international fitness experts.³³

2. Methods

2.1. The evidence supporting the proposed fitness tests

This Delphi study builds on 2 major evidence-based sources, the EU-funded Assessing Levels of PHysical Activity and fitness (ALPHA-FIT) project³¹ and the Institute of Medicine (IOM, currently named the National Academy of Medicine) report from the USA.³⁰ Briefly, the ALPHA project aimed to identify a set of valid, reliable, feasible, and safe field-based fitness tests to assess health-related fitness in school-aged children and adolescents (6–18 years old) to support standardized public health monitoring within the EU. The focus was to select tests that would be easy and feasible for use in school settings. The evidence used to support decisions was based on 4 separate reviews (3 of which employed systematic review methodology): (a) cross-sectional associations between physical fitness and health outcomes;⁴ (b) validity of fitness tests to predict future health;³ (c) criterion validity of fitness tests;³⁴ and (d) test-retest reliability of fitness tests.³⁵ Moreover, a number of methodological studies were conducted to address the knowledge gaps identified,³¹ with the feasibility and safety of the selected tests subsequently studied.³⁶ As an example, skinfold thicknesses were part of the evidence-based test battery but were not included in the high-priority test battery because they showed limited feasibility (due to equipment, expertise, sensitivity issues, and time

needed). Therefore, the high-priority ALPHA fitness test battery proposed the following measures: (a) weight and height (to compute body mass index (BMI)) and (b) waist circumference to assess anthropometry and body composition; (c) the 20-m shuttle run to assess cardiorespiratory fitness; and (d) handgrip strength and (e) standing long jump to assess musculoskeletal fitness.³¹ The ALPHA project provided an operations manual with specific protocols to conduct the tests and supporting videos that are available online at the FitBack website (<https://www.fitbackeurope.eu/en-us/fitness-report/about-testing>).

The IOM report on *Fitness Measures and Health Outcomes in Youth* aimed to recommend the best health-related fitness measures to include in a national fitness survey of children and adolescents, and also to recommend fitness test items that would be feasible to administer in a school environment (Section *Fitness Measures for Schools and Other Education Settings*).³⁰ The evidence used to inform decisions was based on a systematic review of the literature that focused on longitudinal and experimental studies measuring both fitness and health outcomes in children and adolescents aged 5–18 years. The review included evidence on field-based measures of fitness published between the years 2000 and 2010. The full IOM report can be freely accessed at <https://www.ncbi.nlm.nih.gov/books/NBK241315/>. The IOM report proposed the following measures: (a) weight and height (to compute BMI) to assess anthropometry and body composition; (b) the 20-m shuttle run to assess cardiorespiratory fitness; and (c) handgrip strength and (d) standing long jump to assess musculoskeletal fitness.³⁰ The report also recommended measures of skinfold thicknesses and waist circumference for the U.S. National Survey, but not for school-based testing due to challenges with the time and expertise required as well as potential privacy issues when conducting the tests. The IOM report did not recommend specific testing protocols.

2.2. The proposed YFIT battery: Core tests and protocols

We propose the YFIT battery that is aligned with those recommended by the ALPHA project and the IOM report, and as such, the proposed YFIT battery is considered evidence-based, valid, reliable, health-related, feasible, and safe. The proposed core fitness tests for monitoring and surveillance in children and adolescents include (a) weight and height (to compute BMI) to assess body composition; (b) the 20-m shuttle run to assess cardiorespiratory fitness; and (c) handgrip strength and (d) standing long jump to assess muscular fitness. Waist circumference was not included in this proposal, in line with the IOM report, due to potential privacy issues with exposing the abdomen for measurement and possible cultural or ethical issues that may arise in some cultures and countries.

For the YFIT battery, we proposed using the protocol instructions and accompanying videos developed for the ALPHA project. Additional details can be found in the ALPHA Test Manual (<https://www.ugr.es/~cts262/ES/documents/ALPHA-FitnessTestManualforChildren-Adolescents.pdf>). The original ALPHA protocols, after language

333 editing and revision, were used to develop the Delphi survey
 334 materials used in this study (see the original protocols with
 335 track changes made as results of the Delphi survey, [Supple-](#)
 336 [mentary Table 1](#)).

337 2.3. Designing and piloting the Delphi survey

339 We conducted a single-panel modified Delphi study with a
 340 group of international fitness experts, and approval by the
 341 ethics committee was not required. This Delphi study was
 342 developed and reported in accordance with the Conducting
 343 and Reporting Delphi Studies (CREDES) guidelines (See
 344 checklist as [Supplementary Table 2](#)).³⁷ We developed a stan-
 345 dardized survey using Microsoft Forms (Microsoft Corp.,
 346 Redmond, WA, USA). The survey outlined the evidence for
 347 each of the proposed tests, including details on the proposed
 348 test protocols. For each test, we asked participants whether
 349 they agreed with the proposed test for international surveil-
 350 lance and monitoring and whether they agreed with the corre-
 351 sponding protocol. For each question, participants were able to
 352 respond “yes” or “no”. When a participant selected “no”, they
 353 were asked to explain why they disagreed, using an open-
 354 ended response. The survey was piloted with a small group of
 355 experts (Grant R. Tomkinson, Jonatan R. Ruiz, Katja Keller,
 356 and Christine Delisle Nyström, 50 % women) to assess the
 357 clarity of the survey content. The 4 experts involved in the
 358 piloting of the survey did not participate in the main Delphi
 359 study to avoid bias. The Delphi procedure allowed the expert
 360 participants to provide their opinions and to systematically
 361 refine, if necessary, the content to attain consensus.³² We
 362 aimed for agreement of ≥ 80 % among the expert participants
 363 for consensus.^{37–42}

364 2.4. The process of selecting fitness experts to be invited to 365 participate

366 Sampling of expert participants took place in 4 phases.
 367 First, we invited participants who took part in the Global
 368 Youth Fitness Forum on September 7th, 2023 that was orga-
 369 nized by the Public Health Agency of Canada. Most of these
 370 participants co-authored a previous Delphi study.³³ Second,
 371 we ran a SciVal (www.scival.com) search on November 14th,
 372 2023. SciVal is a bibliometric repository that categorizes
 373 Scopus publications into different topic clusters, each repre-
 374 senting a distinct field of research. These topics are identified
 375 through direct citation analysis and named based on key terms
 376 from the aggregated publications with a unique classification
 377 number. The topic identified as “Cardiorespiratory Fitness;
 378 Skinfold Thickness; Body Mass (T.7814)” covers comprehen-
 379 sive studies on physical fitness components and their tests in
 380 youth. Experts who had been a first or senior (i.e., last/corre-
 381 sponding) author on relevant publications in this field since
 382 January 2020 with an h-index of ≥ 5 were invited.³³ Notably,
 383 SciVal updated a new generation of topics on May 21, 2024.
 384 The new topic most relevant to the “youth fitness test” was
 385 “Adolescents; Muscle Strength; Fitness (T.6179),” which
 386 covered 77.6 % of the studies previously categorized under
 387 T.7814. Third, we ran an additional search of [www.](http://www.

 388 <a href=)

389 [expertscape.com](http://www.expertscape.com) on November 9th, 2023 to identify top
 390 researchers who have published in the area of pediatric phys-
 391 ical fitness. Last, we searched our personal networks to iden-
 392 tify additional people with expertise in fitness testing in
 393 children and/or adolescents, while prioritizing those from low-
 394 and middle-income countries and women to ensure that we
 395 obtained insights from a gender and internationally diverse
 396 expert group. It is important to note that to reduce risk of bias
 397 only two of the experts invited to participate in this Delphi
 398 survey participated in the ALPHA project and only two partic-
 399 ipated in the IOM report.

400 2.5. Delphi study methods, data management, and analysis

401 In total, we invited 216 participants with expertise on
 402 fitness in children and adolescents to complete the Delphi
 403 survey. The survey was first circulated by email on December
 404 15th, 2023 and data collection closed on March 8th, 2024. We
 405 provided participants with up to 3 reminder emails between
 406 January and February 2024. All analyses were conducted in
 407 Microsoft Excel 2016 (Microsoft Corp.) or SAS Enterprise
 408 Guide 7.1 (SAS, Cary, NC, USA). Analyses were carried out
 409 as frequencies or means stratified by gender and geographic
 410 region.

411 3. Results

412 3.1. Characteristics of the expert group responding to the 413 Delphi survey

414 The characteristics of the Delphi expert panel members are
 415 presented in [Table 1](#). Of the 216 fitness experts invited to
 416 participate, 169 from 50 countries and territories (including
 417 special administrative regions such as Hong Kong, China)
 418 responded to our survey, resulting in a 78 % response rate
 419 ([Fig. 1](#)). The χ^2 test revealed that the response rates across
 420 both genders and various world regions did not statistically
 421 differ from those not responding, with the exception of Asia,
 422 where the response rate was lower (52.4 %) probably due to
 423 their winter vacation period and the Spring Festival in China.
 424 [Fig. 1](#) visually shows how the percentage of experts
 425 responding from different world region and gender groups is
 426 similar to those invited, indicating overall representativeness
 427 in the participation/response. A detailed gender- and country-
 428 level description of the respondents is provided as [Table 2](#). In
 429 short, respondents were on average 46 years of age, mostly
 430 scientists/researchers (88 %), and had more than 10 years of
 431 postgraduate experience (65 %). A total of 63 women
 432 completed the survey, representing 37 % of respondents.
 433 Sixty-one respondents were from low- or middle-income coun-
 434 tries (self-reported by the respondents), representing 36 % of
 435 respondents. Most respondents were from Europe (54 %),
 436 followed by the Americas (25 %), with a balanced distribution
 437 between North and South America (13 % and 12 %, respec-
 438 tively), followed by Africa (8 %), Asia (7 %), and Oceania
 439 (5 %).

Table 1
Descriptive statistics of the panel of experts ($n = 169$).

	Total	Men	Women
Age (year)	46.3 ± 10.8	47.3 ± 11.4	44.7 ± 9.8
Gender			
Man	105 (62.1)	–	–
Woman	63 (37.3)	–	–
Non-binary	1 (0.6)	–	–
Occupation			
Scientist/researcher (e.g., professor, scientist, post-doctoral fellow)	149 (88.2)	92 (87.6)	56 (88.9)
Research assistant/research manager	4 (2.4)	2 (1.9)	2 (3.2)
Student (e.g., PhD student)	8 (4.7)	5 (4.8)	3 (4.8)
Other	8 (4.7)	6 (5.7)	2 (3.2)
Career stage (years of experience post-graduation)			
Current student	12 (7.1)	6 (5.7)	6 (9.5)
0–5 years	19 (11.2)	10 (9.5)	9 (14.3)
6–10 years	29 (17.2)	19 (18.1)	9 (14.3)
11–20 years	53 (31.4)	31 (29.5)	22 (34.9)
21+ years	56 (33.1)	39 (37.1)	17 (27)
Primary region of occupation			
Africa	14 (8.3)	6 (5.7)	8 (12.7)
Asia	11 (6.5)	8 (7.6)	3 (4.8)
Oceania	9 (5.3)	5 (4.8)	4 (6.3)
Europe	92 (54.4)	61 (58.1)	31 (49.2)
North America	22 (13.0)	13 (12.4)	8 (12.7)
South America	21 (12.4)	12 (11.4)	9 (14.3)
Primary country GDP (self-reported)			
High-income	108 (63.9)	70 (66.7)	37 (58.7)
Middle-income	45 (26.6)	26 (24.8)	19 (30.2)
Low-income	16 (9.5)	9 (8.6)	7 (11.1)

Notes: Data are presented as n (%) or mean ± SD unless otherwise stated. Percentages may not add up to 100 % due to rounding.

Abbreviation: GDP = gross domestic product.

3.2. Main results of the Delphi study

The main results of the Delphi study are presented in Fig. 2 for the full sample, by gender and by world region. The percentage of agreement was above 80 % (targeted consensus) for the 4 fitness measures proposed, precluding the need for a second Delphi survey round. The agreement for using BMI was the lowest at 87 %, with higher agreements obtained for the other tests (i.e., 92 % for handgrip strength, 93 % for 20-m shuttle run, and 98 % for standing long jump). The level of agreement for all proposed tests was above 80 % for both men (85 %–98 %) and women (92 %–97 %) and across all the world regions (85 %–100 %). Comments and suggestions related to the tests proposed, as well as authors' overall answers are presented in Supplementary Tables 3–6.

We also attained consensus for all proposed test protocols, >80 % in the full sample and by gender (Fig. 2). This level of consensus precluded the need for a second Delphi survey round. Levels of agreement for the test protocols were also >80 % for most world regions (Fig. 2). There were, however, a few test protocols in certain regions (4 of 24 bars represented in Fig. 2) where the percentage was lower (i.e., 67 %–68 %). For example, only 68 % of North American participants agreed with the protocol for BMI, and we did not attain the 80 % threshold for the 20-m shuttle run (78 %), handgrip strength (78 %), or standing long jump (67 %) protocols in

Australia and Oceania. It is important to note that choosing “No” in the survey does not necessarily mean disagreement with the proposed protocol, but it was the only way in which to provide opinions/observations. In such cases, participants were able to suggest slight modifications to the protocols or considerations, which are presented in Supplementary Tables 3–6. The original protocols, with changes highlighted, are presented in Supplementary Table 1, and the final protocols (clean of track changes), which were refined based on the expert panel's comments, are presented in Table 3. As seen in the version of the protocols with changes tracked (Supplementary Table 1), the main content of the protocols was not largely modified and, therefore, a second Delphi survey round was not conducted. Importantly, some of the refinements made to the protocols based on comments from the international and diverse expert group included important cultural and religious considerations regarding the clothing recommended for the measurements, which makes YFIT a more inclusive test battery. We also provide additional guidance for the sequence of fitness tests and for handgrip strength testing (i.e., optimal hand size grip span converter table) since it has been shown that maximal handgrip strength is associated with an optimal grip span that is dependent on the hand size of the child⁴³ and adolescent⁴⁴ (Supplementary Tables 7–8). An illustration summarizing the project and findings is presented as Fig. 3.




Fig. 1. The gender and geographic distribution of the (A) Delphi survey invitees ($n = 216$) and responders ($n = 169$) and (B) authors ($n = 26$). Data shown are n (%). Note that 80 (woman) + 135 (man) = 215, which plus 1 person reporting non-binary gender totals 216 survey invitees. The same applies to the 169 Responders.

4. Discussion

4.1. Summary of main findings

In the present study, we proposed the YFIT battery, an evidence-based and international consensus-based fitness test battery for monitoring and surveillance among children and adolescents that is valid, reliable, health-related, feasible, and safe. To attain international consensus on the YFIT battery, we completed a Delphi survey using a large, gender, and geographically diverse expert panel. We obtained a high response rate (78 %) and high agreement (≥ 85 %) for all proposed tests across genders and geographical regions, indicating overall consensus (≥ 80 %) for the core set of fitness tests included in the YFIT battery. The core measures included body weight, body height, and BMI as markers of body size/composition; 20-m shuttle run as a marker of cardiorespiratory fitness; and handgrip strength and standing long jump as markers of upper and lower body muscular fitness, respectively. We also attained an overall consensus (≥ 80 %) for the corresponding test protocols. A number of minor modifications were proposed by respondents, which were incorporated into the final test battery protocol provided in this study. In

addition, there are guidance tools available to help avoid student’s frustration or bad experiences during fitness testing at school that can be accessed on the FitBack website (See: (a) Before fitness testing; and (b) Provide appropriate fitness testing environment).

4.2. Interpretation and relevant comments provided by the respondents

It is important to consider the YFIT battery as the core or minimum number of fitness tests and protocols recommended for international monitoring and surveillance. However, these measures should not dissuade countries/regions from including additional fitness components or other fitness tests depending on the purpose of the evaluations. For example, if a certain country/region has historically collected fitness data using tests other than those proposed here, it is reasonable to continue using some or all of those tests for tracking temporal trends. In this case, the inclusion of the YFIT core tests could be considered as complementary to existing tests. Likewise, if a country/region plans to start a new fitness monitoring/surveillance system, then the YFIT core fitness tests are

Table 2
Detailed gender- and country-level information of the Delphi responders, representing 50 countries and territories.

Countries and territories	<i>n</i>
Africa (N = 8)	14 (8 women)
Botswana	1
Ghana	1
Kenya	4
Malawi	1
Nigeria	1
South Africa	2
Uganda	3
Zimbabwe	1
Asia (N = 4)	11 (3 women)
China	6
Hong Kong, China	3
Japan	1
Saudi Arabia	1
Australia and Oceania (N = 1)	9 (4 women)
Australia	9
Europe (N = 26)	92 (31 women)
Austria	2
Belgium	1
Croatia	3
Czech Republic	2
Denmark	1
Estonia	5
Finland	6
France	3
Germany	3
Greece	1
Hungary	2
Iceland	1
Ireland	1
Italy	6
Lithuania	3
Montenegro	1
North Macedonia	2
Poland	1
Portugal	7
Serbia	2
Slovenia	4
Spain	19
Sweden	3
Switzerland	2
The Netherlands	2
UK	9
North America (N = 3)	22 (8 women)
Canada	11
Mexico	2
USA	9
South America (N = 8)	21 (9 women)
Argentina	3
Brazil	6
Chile	7
Colombia	1
Ecuador	1
Paraguay	1
Peru	1
Uruguay	1
Total (N = 50)	169 (63 women)

Note: The word “territories” includes internationally recognized “special administrative regions”, for example, Hong Kong, China.

recommended at a minimum, plus other potentially relevant tests (if any). Although this study has focused on monitoring/surveillance, the same principle could apply to research projects that include these core tests plus any other tests relevant for their analytical goals. In the future, standardized international fitness measurement will result in a large amount of comparable data across countries/regions, thus opening many possibilities for research while informing policy interventions seeking improvements and equity.

Furthermore, when a test is finally selected, it is of utmost importance to follow the same protocols to increase standardization, data comparability, generation of future reference values, and health-related cut-points. In this context, our study obtained opinions on fitness test protocols from world experts, resulting in some minor protocol refinements. We are not aware of other fitness test protocols that have followed a similar expert consensus process. Therefore, strict adherence to these protocols is highly recommended for standardization. The protocols also include demonstration videos, key information about the equipment needed, as well as assessor instructions and specific instructions for the person being evaluated, which are all important for measurement standardization. Moreover, the protocols suggest a testing sequence and important safety considerations.

The overall consensus on the use of the 20-m shuttle run, handgrip strength, and standing long jump tests was very high (>90 % agreement). Across all participants, the level of agreement for the standing long jump was the highest at 98 %, suggesting it is the preferred fitness test for monitoring and surveillance. Although the agreement for using BMI as a measure of body composition was still high, it was the lowest among the proposed measures (87 %). After reading the comments provided by the expert panel, we believe that this lower agreement was mainly due to BMI being framed as a measure of body composition, which was not considered ideal by some experts because BMI is neither able to distinguish between fat and lean mass nor central/abdominal adiposity. We fully appreciate this concern, and it may be more appropriate for the terminology to change from “body composition” to “body size/weight status”, which appears to better reflect the underlying construct. Regardless of this terminology, we believe that weight, height, and BMI are the minimum anthropometric indices needed in this core fitness battery for several reasons. First, BMI is currently the only globally accepted measure to define overweight and obesity across the lifespan for surveillance purposes. Second, BMI is simple, feasible, and robust—unlike body fat percentage, which is known to be costlier and varies largely by method, even between gold standard methods (e.g., there are large differences between air-displacement plethysmography and dual-energy X-ray absorptiometry⁴⁵). Third, regardless of its validity as a marker of adiposity, BMI is an excellent marker of future disease risk.^{46,47} Fourth, as indicated above, it is important to stress that other field-based measures such as skinfolds or waist




Fig. 2. The percent agreement for BMI, 20mSRT, HGS, and SLJ by total sample, gender, and region of occupation ($n = 169$). Agreement was assessed by asking participants whether they agree with recommending each measure as the best and most feasible to use in international health surveillance and monitoring. Agreement with the protocol was assessed by asking participants whether they agree with the recommended protocol identified for international health surveillance and monitoring. Consensus was considered successful if reaching 80 % agreement or higher. The response options for both questions were “Yes” or “No”. 20mSRT = 20-m shuttle run test; BMI = body-mass index; HGS = handgrip strength; SLJ = standing long jump.

circumference raise privacy and feasibility issues for use at a large scale and especially in certain settings/cultures, and the YFIT battery was developed to be international and inclusive. Fifth, it is also important to note that the separate measures of weight, height, and BMI can be useful for scaling and interpreting the other fitness tests. Sixth, other indices, such as the tri-ponderal mass index (body mass divided by height cubed) or different sex- and age-specific z-score measures, can be computed at any time based on collected weight and height data. Nevertheless, if there are no time, resource, or privacy concerns, additional anthropometric measurements will provide valuable information about body composition and fat distribution. In those cases, our recommendation is to follow the ALPHA and IOM evidence-based proposals,^{30,31} which include skinfold thicknesses (to estimate body fat percentage) and waist circumference (which also allows for abdominal adiposity and body shape (i.e., waist-to-height ratio) to be estimated).

It is important to note that the systematic reviews that informed the ALPHA project and IOM report were conducted roughly a decade ago. However, more recent systematic reviews also support the same key fitness tests proposed. For the cardiorespiratory fitness component, there is strong evidence suggesting the 20-m shuttle run is the best field-based measure given it is reliable,⁴⁸ valid,⁴⁹ health-related,⁵⁰ health-discriminant,⁵¹ and scalable for use in large-scale surveys.⁵² For the muscular fitness component, García-Hermoso and colleagues² concluded that the handgrip strength and standing long jump were the most studied tests in relation to health outcomes in children and adolescents, and Fraser and colleagues⁵³ concluded that handgrip strength and standing long jump were the 2 muscular fitness tests with the highest health-related discriminatory ability. The usefulness of BMI as an internationally agreed metric to define overweight and obesity for surveillance purposes in children and adolescents and its association with health outcomes is well supported by

Table 3

The final protocol descriptions for each of the identified fitness tests after incorporating modifications based on suggestions from the Delphi participants.

Fitness test	Content	Description
Body composition	Demonstration	See the video link (https://www.youtube.com/watch?v=BXhqQZcEaLk).
	Equipment	An electronic scale and a telescopic height-measuring instrument.
	Examiner instructions	(a) Body mass: The child must stand on the platform of the scale without support. The child stands still over the center of the platform, with the body weight evenly distributed between both feet. Light clothing (as culturally accepted) is recommended for this measurement, excluding shoes, heavy long pants and sweaters. (b) Body height: Hair ornaments (e.g., hats) must be removed, braids undone, and ponytails must be positioned at the back, not on top. The child stands on the stadiometer without shoes and with feet placed slightly apart and the back of the head, shoulders, buttocks, calves, and heels touching the vertical board. Legs must be kept straight and the feet flat. The tester must position the child's head so the ear canal and the lower edge of the eye socket are parallel to the base-board (i.e., the Frankfort plane positions horizontally). The headboard must be pulled down to rest firmly on top of the head while flattening hair. The measurement is registered after the child is asked to take a deep breath in. If hair/head ornaments cannot be removed due to cultural reasons, the tester should estimate the height in centimeters (cm) after subtracting the height of the hair/head ornament.
20-m shuttle run test	Scoring	Two measurements of both body weight and body height are performed, and the mean of each is retained. Weight is recorded in kilograms to 0.1 kg, and height is recorded in centimetres to 0.1 cm. <i>Example:</i> 58.4 kg and 157.3 cm.
	Demonstration	See the video link (https://www.youtube.com/watch?v=Fg7Suqa46hU).
	Equipment	Select a test site, preferably a 22- to 25-m-long gym. An outdoor court or a grass field is also suitable, provided it has enough space and is even in surface. Allow for a space of at least 1 m at either end of the track. A wide area is recommended to test more children simultaneously, allowing 1 m between each child to improve safety. The running surface should be as flat as possible and not slippery. The 2 ends of the 20-m track should be clearly marked. Additional equipment includes 4 cones to mark the 20-m distance, a tape to measure the 20-m distance, a speaker or device to play the audio cues, and the pre-recorded audio cues. The recommended audio version, ideal for recording half stages, is available here: (https://onedrive.live.com/?authkey=%21AKuxDwIL7LEpBG0&id=8F0D6BC76AAB9FAE%2163832&cid=8F0D6BC76AAB9FAE&parId=root&parQt=sharedby&o=OneUp). Note 1: We acknowledge the support of Prof. Léger from the University of Montreal, Canada for donating the original audio file of the 20-m shuttle run test under the framework of the FitBack platform, where the audio file is hosted. Note 2: If the minimum space (i.e., 22 m) required to conduct this test is not available within the school facilities (either indoors or outdoors), consider the possibility of conducting the test in the close surroundings of the school (e.g., a park or sport facility) as long as the safety considerations are maintained (flat and non-slippery surfaces, no cars or objects that can be harmful, etc.).
Handgrip strength	Examiner instructions	Children are required to run between 2 lines 20-m apart in pace with the audio signals. The initial running speed is 8.5 km/h with the speed increasing by 0.5 km/h at each consecutive min (1 min is equal to 1 stage in the test). The examiner terminates the test when the child fails to reach the line for 2 consecutive times. Otherwise, the test ends when the child stops due to fatigue. Since pacing for this test can be difficult for children, the examiner can allow some flexibility when arriving to the line slightly before or after the audio signal for the first couple laps of a stage. "The shuttle run test gives an indication of your aerobic capacity, i.e., your endurance, and involves running there and back along a 20-m track. Speed will be controlled by an audio track emitting a beeping sound at regular intervals. Pace yourselves to be at one end of the 20-m track or the other when you hear a sound. Touch the line at the end of the track with your foot, then turn and run in the opposite direction. At first, the speed is slow, but it will increase slowly and steadily every minute. Your aim in the test is to follow the set rhythm for as long as you can. You should stop when you can no longer keep up with the set rhythm. Remember the number announced by the recording when you stop, that is your score. The length of the test varies according to the individual: the fitter you are, the longer the test lasts. To sum up, the test is maximal and progressive; in other words, easy at the beginning and hard towards the end. Good Luck!"
	Participant instructions	
	Scoring	After the child has stopped the test, the last completed (i.e., audio announced) half stage is recorded. Only 1 test trial is needed. <i>Example:</i> a score of 6.5 stages. If higher precision is required (e.g., intervention studies aiming to detect small changes), the final time spent in the test can be expressed in seconds (s) in addition to the number of shuttles/laps.

(continued on next page)

Table 3 (Continued)

Fitness test	Content	Description
	Examiner instructions	Place the top of the dynamometer so it lies across the middle of the palm. Hand size should be measured as the distance separating the distal extremes of the first and 5th fingers of the right hand. The result of the hand size should be rounded to the nearest whole cm. Alternatively, a hand size ruler can be used (see Pages 18–19: https://www.ugr.es/~cts262/ES/documents/ALPHA-FitnessTestManualforChildren-Adolescents.pdf). Using the hand size, adjust the dynamometer grip span. Ask the participant which is their dominant hand to record on the result sheet. The tester can ask the participant which hand is used for throwing if the participant isn't confident in which hand is dominant. The dominant hand should be tested first. During the test, the arm and hand holding the dynamometer should not touch the body. The dynamometer is held in line with the forearm and hangs down at the side with the hand held in a neutral position with the thumb forward. While standing (if possible) the child squeezes gradually and continuously for at least 2 seconds (s), performing the test twice (alternating hands) with the optimal grip span (previously calculated, according to the hand size, see Supplementary Tables 7–8) and allowing a short 30-s rest between testing rounds. The indicator must be returned to 0 after each attempt, depending on the dynamometer.
	Participant instructions	"Take the dynamometer with one hand, the dominant hand first. Squeeze it as forcefully as you can while holding the dynamometer away from your body. Don't let it touch you during the test. Squeeze gradually and continuously for at least 2 s. Do the test twice per hand: the best result scores."
	Scoring	Both hands are to be tested twice (dominant and non-dominant, 30-s break, dominant and non-dominant), and the best result (of each hand) is scored. The participants should indicate their dominant hand (the one used for throwing) with that information being recorded (e.g., dominant hand = right or left or ambidextrous). The result is expressed in kg to the closest 0.1 kg. <i>Example:</i> a result of 24 kg scores 24.0 kg. It is very important to report the values for the right and left hand separately (indicating the dominant), which will allow in the future any type of analyses (including asymmetry analyses). As a general recommendation and in addition to reporting the value of both hands separately, reporting the average of both hands is informative from a health point of view since it indicates the overall handgrip strength level of the person that is informative of both hands.
Standing long jump	Demonstration	See the video link (https://www.youtube.com/watch?v=jHJ5OMZE_MA).
	Equipment	A non-slippery hard surface, a stick to identify the landing location, a tape to measure the jumping distance, an adhesive tape to identify the start line, and cones to mark off the testing location.
	Examiner instructions	Jumping for distance from a standing start and with both feet at the same time. One horizontal line is drawn or marked using tape to identify the start line. Place a tape measure at a right angle to and off to the side of the start line. The jumping distance is measured from the jump line to the point where the back of the heel nearest to the start line lands on the ground. The child is asked to place their toes behind the start line. While bending their knees and swinging their arms, they are asked to jump as far as possible while balanced on both feet. If the child falls backward or touches the floor with their hands or any body part behind their feet, the attempt needs to be repeated. If the child falls forward but keeps the back foot planted after landing, the test is considered valid.
	Participant instructions	"Stand with your feet shoulder-width apart and toes just behind the line. Bend your knees with your arms in front of you, parallel to the ground. As you swing both arms, push off vigorously and jump as far as possible, taking off and landing with both feet at the same time. Try to land with your feet together and stay upright. If you fall forward after landing, try to keep your back foot planted so that the measurement can be done, otherwise the jump needs to be repeated. If you fall backward after landing, you need to repeat the jump."
	Scoring	Two trials are carried out and the best result is recorded. The result is recorded to the nearest full cm. <i>Example:</i> a jump of 1 m 56 cm scores 156 cm.
	Recommended sequence for testing and safety considerations (Note: This section with recommended sequence and safety considerations was not included in the Delphi survey and comes from the original ALPHA manual of operations, but the authors consider it worth including together with the protocols information.)	<ol style="list-style-type: none"> Whenever possible, it is better to measure weight and height first before participants start sweating and losing body fluids (i.e., before warming up). A short warm-up (5 min) including jogging in a line while focusing on changing directions and speed. Dynamic stretching exercises are encouraged during the warm-up. Practice some long jumps with both feet at the same time to become familiar with the later testing. Conduct the handgrip strength and standing long jump in random order after the warm-up. Finally, the 20-m shuttle run should be done last since it is a maximal test and children are highly fatigued afterward. <p>*During all testing the examiners should use verbal encouragement to motivate participants.</p> <p>Safety considerations.</p> <p>A non-slippery surface is necessary for the standing long jump and the 20-m shuttle run tests. For the 20-m shuttle run test, a minimum of 1 extra m at each side is required (i.e., a total of 22 m is the minimum needed), but if more space is available, a 25-m-length space is better for a safe administration of the 20-m shuttle run test. Avoid fitness testing in very hot environments.</p> <p>Parents/legal guardians and/or children and adolescents should inform the evaluators of any condition that might be considered a contraindication for vigorous exercise. As a general rule, any child who participates without restrictions in physical education classes are able to participate in these fitness tests. If there is any doubt about whether a person can do the physical testing, the Physical Activity Readiness Questionnaire for Everyone (PAR-Q+) should be completed by the participant. If the participant answers YES to 1 or more questions, the person needs to check with their doctor before participating in the fitness evaluations. In any case, it is important for the evaluator to be alert to any negative symptoms during testing, such as skin paleness, dizziness, or any other adverse symptoms. The tests should be immediately stopped if there is any sign of problems during the testing.</p>

Abbreviation: ALPHA = Assessing Levels of Physical Activity.




Fig. 3. Illustration summarizing the whole project and findings.

extensive evidence and has been discussed above. Finally, an additional advantage of the YFIT core fitness tests is that they are not new; historically, they have been considered acceptable for use in many countries. As an example, temporal trends in fitness using the proposed tests have been published in different parts of the world.^{54–62} Likewise, data have been pooled using the proposed tests to generate norm-referenced values based on 8-million test results for European children and adolescents aged 6–18 years⁶³ as well as to develop the FitBack free-access, automated, multilingual web-platform that facilitates interpretation of fitness assessment. Moreover, we previously published international normative values for the 20-m shuttle run test⁶⁴ and are currently working to update these norms and develop new international normative values for handgrip strength and standing long jump for those aged 6–18 years. These will be made freely available when complete.

If the proposed fitness tests and protocols are widely adopted, initiatives like FitBack as well as the MOratorREsearch (MO|RE) data open access database in Europe,⁶⁵

can be extended to other world regions, leading to the development of a global fitness observatory.

4.3. Limitations and strengths

A lower proportion of women and researchers from Africa and Asia (and none from Central America) participated in the Delphi survey, which further reflects the under-representation of these groups in research in general^{66–68} and in sport/exercise sciences in particular.^{69,70} Nevertheless, we specifically targeted these under-represented groups and managed to include >60 women and >60 researchers from low- and middle-income countries with the requisite topical expertise, which represented one-third of all participants. Although we have gathered a large and diverse sample of participants, the results of this and any other Delphi survey reflect the opinions only of those included in the study, and including different participants could potentially have resulted in different outcomes. Likewise, there is also risk of respondent bias, since the participants were not blinded to the identity of the authors leading the study.

No fitness test is perfect, and all come with certain limitations. For example, the 20-m shuttle run, despite being one of the most widely used tests in the world, has the main limitation of requiring physical space, namely a flat and non-slippery surface that is at least 22 m long. If this minimum space is not available within the school facilities (either indoors or outdoors), assessors can consider the possibility of conducting the test in the close surroundings of the school (e.g., a park or sport facility) as long as safety is maintained (flat and non-slippery surfaces, no cars or objects that can be harmful, *etc.*). Likewise, handgrip strength testing has the limitation of needing handgrip dynamometers, which may not be available or affordable in certain cases. However, a recent study has shown that low-cost handgrip dynamometers (USD\$45) are highly valid and reliable and produce similar results as other dynamometers that are 10 times more expensive, such as Jamar or TKK dynamometers.^{71,72} If any of the YFIT core tests are not feasible to conduct for any reason, it is better to measure as many of the core tests as possible, as they would still provide valuable health-related fitness information. Finally, it is important to acknowledge that the YFIT battery is meant for the general population and is not adapted for children and adolescents with special needs. However, some fitness monitoring systems, such as those in Hungary and Finland, have already adapted fitness testing; guidance on this is provided on the FitBack platform (https://www.fitbackeu.rope.eu/Portals/0/Adapted_FitBack_final.pdf?ver=2022-08-10-124500-620) and future research directions have recently been discussed.^{73,74} When more evidence is accumulated in the future, similar methods to those presented here might lead to consensus on fitness testing adapted for children and adolescents with special needs.

5. Conclusion

This study provides an evidence-based, gender and geographically diverse, international consensus on the fitness tests and protocols for monitoring and surveillance of children and adolescents worldwide. The YFIT battery resulting from this consensus study includes: weight, height, and BMI as markers of body size (weight status) and composition; 20-m shuttle run as a marker of cardiorespiratory fitness; and handgrip strength and standing long jump as markers of muscular fitness. This core set of tests can be considered evidence-based, valid, reliable, health-related, feasible, safe, and inclusive. This consensus is based on high agreement rates (87 %–98 %) and is consistent among diverse experts across the world. Importantly, we made specific protocol refinements based on expert feedback. When these tests and protocols are applied consistently worldwide, they will improve standardization and opportunities for international comparison, data pooling, health-related cut-point development, and informed policy changes. Given the strong evidence suggesting fitness as a powerful, non-invasive, and feasible public health marker among children and adolescents, international efforts using the proposed tests and protocols will provide highly valuable information for monitoring and surveillance purposes across the globe.

Authors' contributions

FBO conceived the study, and participated in its design and coordination and wrote the first draft of the manuscript; FBO, KZ and JJJ designed the Delphi survey; KZ and JJJ created the survey in Microsoft Forms, handled the data and created the tables/figures; GT, JRR, KK and CDN participated in the piloting of the Delphi survey. All the authors contributed intellectually to the content of the manuscript. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

FBO's research activity is supported by the Grant [PID2020-120249RB-I00](#) and [PID2023-148404OB-I00](#) funded by [MCIN/AEI/10.13039/501100011033](#) and by the Andalusian Government (Junta de Andalucía, Plan Andaluz de Investigación, ref. P20_00124), and by the Erasmus+ Sport Programme of the European Union within the project FitBack4Literacy (No [101089829](#)). Additional support is provided by the University of Granada, Plan Propio de Investigación, Units of Excellence; Unit of Excellence on Exercise, Nutrition and Health (UCEENS), by the CIBERobn Physiopathology of Obesity and Nutrition, and by the Spanish Network in Exercise and Health, EXERNET Network ([RED2022-134800-T](#) and [EXP_99828](#)).

Supplementary materials

Supplementary material associated with this article can be found in the online version at [doi:10.1016/j.jshs.2024.101012](https://doi.org/10.1016/j.jshs.2024.101012).

Appendix: The names and affiliations of the Fitness Expert Group that responded to the Delphi survey (*n* = 169)

Adewale L. Oyeyemi, University of Maiduguri, Nigeria; Adilson Marques, University of Lisbon, Portugal; Aidan Gribbon, Statistics Canada, Canada; Anelise Reis Reis Gaya, Federal University of Pelotas, Brazil; Antonio García-Hermoso, Public University of Navarre, Spain; Arja Sääkslahti, University of Jyväskylä, Finland; Avery Faigenbaum, The College of New Jersey, USA; Bernadette Nakabazzi, Makerere University, Uganda; Bojan Leskošek, University of Ljubljana, Slovenia; Boris Popovic, University of Novi Sad, Serbia; Brendan O'Keeffe, University of Limerick, Ireland; Brian W. Timmons, McMaster University, Canada; Brigita Mieziene, Lithuanian Sports University, Lithuania; Brooklyn Fraser, University of Tasmania, Australia; Caroline Brand, Pontifical Catholic University of Valparaíso, Chile; Cecilia Anza-Ramirez, Universidad Peruana Cayetano Heredia, Peru; Celia Alvarez-Bueno, University of Castilla-La Mancha, Spain; Cesar A. Agostinis-Sobrinho, Klaipeda University, Lithuania; Charles H. Hillman, Northeastern University, USA; Claudia Niessner, Karlsruhe Institute of

- 1359 Technology, Germany; Clemens Drenowatz, University of
1360 Education Upper Austria, Austria; Costan G. Magnussen,
1361 Baker Heart and Diabetes Institute, Australia; Cristina
1362 Cadenas-Sanchez, University of Granada, Spain; Daniel
1363 Berglind, Karolinska Institutet, Sweden; Danilo Rodrigues
1364 Pereira da Silva, Federal University of Sergipe, Brazil; Dario
1365 Colella, University of Salento, Italy; David Lubans, University
1366 of Newcastle, Australia; David Matelot, Université Bretagne-
1367 Sud, France; Dawn M. Tladi, University of Botswana,
1368 Botswana; Diego Augusto Santos Silva, Federal University of
1369 Santa Catarina, Brazil; Diego Moliner-Urdiales, Universitat
1370 Jaume I, Spain; Dot Dumuid, University of South Australia,
1371 Australia; Dylan Blain, University of Wales Trinity Saint
1372 David, UK; Edtna Jáuregui Ulloa, University of Guadalajara,
1373 Mexico; Emeljanovas Arunas, Lithuanian Sports University,
1374 Lithuania; Enrique Pintos-Toledo, University of the Republic,
1375 Uruguay; Eric Tsz Chun Poon, Chinese University of Hong
1376 Kong, Hong Kong, China; Eun-Young Lee, Queen's Univer-
1377 sity, Canada; Eva Maria Riso, University of Tartu, Estonia;
1378 Evelin Mäestu, University of Tartu, Estonia; Farid Bardid,
1379 University of Strathclyde, UK; Felicia Cañete, Ministry of
1380 Public Health and Social Welfare of Paraguay, Paraguay;
1381 Fernando Rodríguez Rodríguez, Pontificia Universidad
1382 Católica de Valparaíso, Chile; Gabriela De Roia, Universidad
1383 de Flores, Argentina; Garden Tabacchi, University of Palermo,
1384 Italy; Gareth Stratton, Swansea University, UK; Gavin R.
1385 Sandercock, University of Essex, UK; Georgi Georgiev, Ss.
1386 Cyril and Methodius University, North Macedonia; Germán
1387 Vicente-Rodríguez, University of Zaragoza, Spain; Gerson
1388 Ferrari, University of Santiago de Chile, Chile; Gil Rosa,
1389 Universidade de Lisboa, Portugal; Greg Welk, Iowa State
1390 University, USA; Gregor Jurak, University of Ljubljana, Slo-
1391 venia; Gregor Starc, University of Ljubljana, Slovenia; Idoia
1392 Labayen, Public University of Navarre, Spain; Igor Cigarroa,
1393 Universidad Santo Tomás, Chile; Irene Esteban-Cornejo,
1394 University of Granada, Spain; Iván Clavel San Emeterio,
1395 University of A Coruña, Spain; Ivana Milanović, University of
1396 Belgrade, Serbia; Janine Clarke, Statistics Canada, Canada;
1397 Janusz Dobosz, Józef Piłsudski University of Physical Educa-
1398 tion in Warsaw, Poland; Jarek Mäestu, University of Tartu,
1399 Estonia; Javier Brazo-Sayavera, Universidad Pablo de
1400 Olavide, Spain; Jennifer Sacheck, George Washington Univer-
1401 sity, USA; Jennifer Servais, Statistics Canada, Canada;
1402 Jeremías David Secchi, Universidad Adventista del Plata,
1403 Argentina; Jérémy Vanhelst, Université Sorbonne Paris Nord,
1404 France; Jesús Viciano Ramírez, University of Granada, Spain;
1405 João Magalhães, University of Lisbon, Portugal; Johana Soto-
1406 Sánchez, Universidad Mayor, Chile; Johannes Jaunig, Univer-
1407 sity of Graz, Austria; John J. Reilly, University of Strathclyde,
1408 UK; Jordan Smith, University of Newcastle, Australia; Jorge
1409 Mota, University of Porto, Portugal; Jose Castro-Piñero,
1410 University of Cádiz, Spain; José Francisco López-Gil,
1411 Currently at Universidad de Las Américas, Ecuador (formerly
1412 at Universidad Loyola Andalucía, Spain, during the Delphi
1413 survey period); Julia Wambui Githang'a, Kenyatta Univer-
1414 sity, Kenya; Júlio Brugnara Mello, Pontificia Universidad
1415 Católica de Valparaíso, Chile; Kathleen Janz, University of
Iowa, USA; Kathryn L. Weston, University of Strathclyde,
UK; Kelly Laurson, Illinois State University, USA; Kevin Till,
Leeds Beckett University, UK; Khanyile Dlamini, National
University of Science and Technology, Zimbabwe; Konstan-
tinos D. Tambalis, National and Kapodistrian University of
Athens, Greece; Laura Basterfield, Newcastle University, UK;
Laura Joensuu, University of Jyväskylä, Finland; Laurent
Béghin, Lille University Hospital, France; Leandro dos
Santos, Londrina State University, Brazil; Lillian Mugisha,
Makerere University, Uganda; Liye Zou, Shenzhen University,
China; Juan Ricardo López Y Taylor, University of Guadala-
ajara, Mexico; Lovro Štefan, Masaryk University, Czechia; Luc
Léger, University of Montreal, Canada; Lucy-Joy Wachira,
Kenyatta University, Kenya; Luís B. Sardinha, University of
Lisbon, Portugal; Lukas Rubin, Technical University of
Liberec, Czechia; Mablina Thuany, States University of Pará,
Brazil; Magdalena Cuenca-García, University of Cádiz, Spain;
Mai Chin A Paw, Amsterdam University Medical Centre, the
Netherlands; Mairena Sanchez-Lopez, University of Castilla-
La Mancha, Spain; Maret Pihu, University of Tartu, Estonia;
Maria Jose Noriega, University of Cantabria, Spain; María
Medrano-Echeverría, Public University of Navarre, Spain;
Maria Reyes Beltran-Valls, Jaume I University, Spain; Marie
Löf, Karolinska Institute, Sweden; Marjeta Misigoj-Dura-
kovic, University of Zagreb, Croatia; Mark Tremblay, Child-
ren's Hospital of Eastern Ontario, Canada; Markus Gerber,
University of Basel, Switzerland; Maroje Sorić, University of
Zagreb, Croatia; Matteo Giuriato, University of Pavia, Italy;
Matteo Vandoni, University of Pavia, Italy; Matthieu Lenoir,
Ghent University, Belgium; Mauro D. Santander, Observatory
of sport Physical activity and culture of Neuquén, Argentina;
Miguel Peralta, University of Lisbon, Portugal; Mikko Huhti-
niemi, University of Jyväskylä, Finland; Mohamed Ahmed
Said, King Faisal University, Saudi Arabia; Mónika Kaj,
Hungarian School Sport Federation, Hungary; Naomi Burn,
University of South Australia, Australia; Nicola D. Ridgers,
University of South Australia, Australia; Nicola Lovecchio,
University of Bergamo, Italy; Nicolas Aguilar-Farias, Univer-
sidad de La Frontera, Chile; Niels Wedderkopp, University of
Southern Denmark, Denmark; Oscar L. Veiga, Autonomous
University of Madrid, Spain; Pedro Saint-Maurice, Champali-
maud Foundation, Portugal; Peter Katzmarzyk, Pennington
Biomedical Research Center, USA; Pontus Henriksson, Link-
öping University, Sweden; Rachel Colley, Statistics Canada,
Canada; Reginald T-A. Ocansey, University of Ghana, Ghana;
Reinhold Kliegl, University of Potsdam, Germany; Robinson
Ramírez-Vélez, Currently at the Public University of Navarre,
Spain (yet originally from Colombia with large experience in
fitness testing in Colombia); Ronald Ssenyonga, Makerere
University, Uganda; Rowena Naidoo, University of KwaZulu-
Natal, South Africa; Russell R. Pate, University of South Caro-
lina, USA; Ryan McGrath, North Dakota State University,
USA; Saima Kuu, Tallinn University, Estonia; Sanja Salaj,
University of Zagreb, Croatia; Seryozha Gontarev, Ss. Cyril
and Methodius University, North Macedonia; Shawnda A.
Morrison, Currently at National University of Singapore,
Singapore (formerly at University of Ljubljana, Slovenia,

during the Delphi survey period); Siphesihle Nqweniso, Nelson Mandela University, South Africa; Sitong Chen, Victoria University, Australia; Stanley Kagunda, Kenyatta University, Kenya; Stephanie Prince Ware, Public Health Agency of Canada, Canada; Stephen H. Wong, Chinese University of Hong Kong, Hong Kong, China; Stevo R. Popovic, University of Montenegro, Montenegro; Stuart J. Fairclough, Edge Hill University, UK; Susana Andrade, University of Cuenca, Ecuador; Susi Kriemler, University of Zurich, Switzerland; Tamás Csányi, Hungarian University of Sports Science, Hungary; Taru Manyanga, University of Northern British Columbia, Canada; Tawonga W. Mwase-Vuma, University of Malawi, Malawi; Tetsu Kidokoro, Nippon Sport Science University, Japan; Thayse Natacha Gomes, Federal University of Sergipe, Brazil; Thordis Gisladottir, University of Iceland, Iceland; Tim Takken, University Medical Center Utrecht, the Netherlands; Timo Jaakkola, University of Jyväskylä, Finland; Timo Lakka, University of Eastern Finland, Finland; Timothy Olds, University of South Australia, Australia; Tuija Tammelin, Jamk University of Applied Sciences, Finland; Urs Granacher, University of Freiburg, Germany; Valerie Carson, University of Alberta, Canada; Vincent Martinez-Vizcaino, University of Castilla-La Mancha, Spain; Vincent Onywera, KCA University, Kenya; Vittoria Carnevale Pellino, University of Pavia, Italy; Wendy Y. Huang, Hong Kong Baptist University, Hong Kong, China; Xiaojian Yin, Shanghai Institute of Technology, China; Yang Liu, Shanghai University of Sport, China; Yi Song, Peking University, China; Yi Sun, East China Normal University, China; Yuan Liu, Shanghai University, China.

References

- García-Hermoso A, Ramírez-Vélez R, García-Alonso Y, Alonso-Martínez AM, Izquierdo M. Association of cardiorespiratory fitness levels during youth with health risk later in life: A systematic review and meta-analysis. *JAMA Pediatr* 2020;**174**:952–60.
- García-Hermoso A, Ramírez-Campillo R, Izquierdo M. Is muscular fitness associated with future health benefits in children and adolescents? A systematic review and meta-analysis of longitudinal studies. *Sport Med* 2019;**49**:1079–94.
- Ruiz JR, Castro-Piñero J, Artero EG, et al. Predictive validity of health-related fitness in youth: A systematic review. *Br J Sports Med* 2009;**43**:909–23.
- Ortega FB, Ruiz JR, Castillo MJ, Sjörström M. Physical fitness in childhood and adolescence: A powerful marker of health. *Int J Obes (Lond)* 2008;**32**:1–11.
- Ortega FB, Silventoinen K, Tynelius P, Rasmussen F. Muscular strength in male adolescents and premature death: Cohort study of one million participants. *BMJ* 2012;**345**:e7279. doi:10.1136/bmj.e7279.
- Cadenas-Sanchez C, Mena-Molina A, Torres-Lopez LV, et al. Healthier minds in fitter bodies: A systematic review and meta-analysis of the association between physical fitness and mental health in youth. *Sports Med* 2021;**51**:2571–605.
- Rabiee R, Agardh E, Kjellberg K, Falkstedt D. Low cardiorespiratory fitness in young adulthood and future risk of disability pension: A follow-up study until 59 years of age in Swedish men. *J Epidemiol Community Health* 2015;**69**:266–71.
- Henriksson H, Henriksson P, Tynelius P, et al. Cardiorespiratory fitness, muscular strength, and obesity in adolescence and later chronic disability due to cardiovascular disease: A cohort study of 1 million men. *Eur Heart J* 2020;**41**:1503–10.
- Henriksson H, Henriksson P, Tynelius P, Ortega FB. Muscular weakness in adolescence is associated with disability 30 years later: A population-based cohort study of 1.2 million men. *Br J Sports Med* 2018;**53**:1221–30.
- Henriksson P, Henriksson H, Tynelius P, et al. Fitness and body mass index during adolescence and disability later in life: A cohort study. *Ann Intern Med* 2019;**170**:230–9.
- Henriksson P, Shiroma EJ, Henriksson H, et al. Fit for life? Low cardiorespiratory fitness in adolescence is associated with a higher burden of future disability. *Br J Sports Med* 2021;**55**:128–9.
- Laakso PTT, Ortega FB, Huotari P, Tolvanen AJ, Kujala UM, Jaakkola TT. The association of adolescent fitness with cardiometabolic diseases in late adulthood: A 45-year longitudinal study. *Scand J Med Sci Sport* 2024;**34**:e14529. doi:10.1111/sms.14529.
- Raghuveer G, Hartz J, Lubans DR, et al. Cardiorespiratory fitness in youth: An important marker of health: A scientific statement from the American Heart Association. *Circulation* 2020;**142**:e101–18.
- Mora-Gonzalez J, Esteban-Cornejo I, Cadenas-Sanchez C, et al. Physical fitness, physical activity, and the executive function in children with overweight and obesity. *J Pediatr* 2019;**208**:50–56.e1. doi:10.1016/j.jpeds.2018.12.028.
- Cadenas-Sanchez C, Migueles JH, Esteban-Cornejo I, et al. Fitness, physical activity and academic achievement in overweight/obese children. *J Sports Sci* 2020;**38**:731–40.
- Mora-Gonzalez J, Esteban-Cornejo I, Cadenas-Sanchez, et al., et al. Fitness, physical activity, working memory, and neuroelectric activity in children with overweight/obesity. *Scand J Med Sci Sport* 2019;**29**:1352–63.
- Mora-Gonzalez J, Esteban-Cornejo I, Migueles JH, et al. Physical fitness and brain source localization during a working memory task in children with overweight/obesity: The ActiveBrains project. *Dev Sci* 2021;**24**:e13048. doi:10.1111/desc.13048.
- Mora-Gonzalez J, Esteban-Cornejo I, Solis-Urra P, et al. Fitness, physical activity, sedentary time, inhibitory control, and neuroelectric activity in children with overweight or obesity: The ActiveBrains project. *Psychophysiology* 2020;**57**:e13579. doi:10.1111/psyp.13579.
- Rodriguez-Ayllon M, Esteban-Cornejo I, Verdejo-Román J, et al. Physical fitness and white matter microstructure in children with overweight or obesity: The ActiveBrains project. *Sci Rep* 2020;**10**:12469. doi:10.1038/s41598-020-67996-2.
- Esteban-Cornejo I, Rodriguez-Ayllon M, Verdejo-Roman J, et al. Physical fitness, white matter volume and academic performance in children: Findings from the ActiveBrains and FITKids2 projects. *Front Psychol* 2019;**10**:208. doi:10.3389/fpsyg.2019.00208.
- Ortega FB, Campos D, Cadenas-Sanchez C, et al. Physical fitness and shapes of subcortical brain structures in children. *Br J Nutr* 2019;**122** (Suppl. 1):S49–58.
- Esteban-Cornejo I, Cadenas-Sanchez C, Contreras-Rodriguez O, et al. A whole brain volumetric approach in overweight/obese children: Examining the association with different physical fitness components and academic performance. The ActiveBrains project. *Neuroimage* 2017;**159**:346–54.
- Esteban-Cornejo I, Mora-Gonzalez J, Cadenas-Sanchez C, et al. Fitness, cortical thickness and surface area in overweight/obese children: The mediating role of body composition and relationship with intelligence. *Neuroimage* 2019;**186**:771–81.
- Cadenas-Sanchez C, Migueles JH, Erickson KI, Esteban-Cornejo I, Catena A, Ortega FB. Do fitter kids have bigger brains? *Scand J Med Sci Sports* 2020;**30**:2498–502.
- Joensuu L, Csányi T, Huhtiniemi M, et al. How to design and establish a national school-based physical fitness monitoring and surveillance system for children and adolescents: A 10-step approach recommended by the FitBack network. *Scand J Med Sci Sports* 2024;**34**:e14593. doi:10.1111/sms.14593.
- Brazo-Sayavera J, Silva DR, Lang JJ, et al. Physical fitness surveillance and monitoring systems inventory for children and adolescents: A scoping review with a global perspective. *Sports Med* 2024;**54**:1755–69.
- Martinko A, Sorić M, Jurak G, Starc G. Physical fitness among children with diverse weight status during and after the COVID-19 pandemic: A

- population-wide, cohort study based on the Slovenian physical fitness surveillance system (SLOfit). *Lancet Reg Health Eur* 2023;**34**:100748. doi:10.1016/j.lanepe.2023.100748.
28. Jurak G, Morrison SA, Kovač M, et al. A COVID-19 crisis in child physical fitness: Creating a barometric tool of public health engagement for the Republic of Slovenia. *Front Public Health* 2021;**9**:644235. doi:10.3389/fpubh.2021.644235.
 29. Kidokoro T, Tomkinson GR, Lang JJ, Suzuki K. Physical fitness before and during the COVID-19 pandemic: Results of annual national physical fitness surveillance among 16,647,699 Japanese children and adolescents between 2013 and 2021. *J Sport Health Sci* 2023;**12**:246–54.
 30. Institute of Medicine (IOM). *Fitness measures and health outcomes in youth*. Washington, DC: The National Academies Press; 2012.
 31. Ruiz JR, Castro-Piñero J, España-Romero V, et al. Field-based fitness assessment in young people: The ALPHA health-related fitness test battery for children and adolescents. *Br J Sports Med* 2011;**45**:518–24.
 32. Dalkey N, Helmer O. An experimental application of the Delphi method to the use of experts. *Manage Sci* 1963;**9**:458–67.
 33. Lang JJ, Zhang K, Agostinis-Sobrinho C, et al. Top 10 international priorities for physical fitness research and surveillance among children and adolescents: A Twin-Panel Delphi Study. *Sports Med* 2023;**53**:549–64.
 34. Castro-Piñero J, Artero EG, España-Romero V, et al. Criterion-related validity of field-based fitness tests in youth: A systematic review. *Br J Sports Med* 2010;**44**:934–43.
 35. Artero EG, España-Romero V, Castro-Piñero J, et al. Reliability of field-based fitness tests in youth. *Int J Sports Med* 2011;**32**:159–69.
 36. España-Romero V, Artero EG, Jimenez-Pavón D, et al. Assessing health-related fitness tests in the school setting: Reliability, feasibility and safety; the ALPHA Study. *Int J Sports Med* 2010;**31**:490–7.
 37. Jünger S, Payne SA, Brine J, Radbruch L, Brearley SG. Guidance on Conducting and REporting DELphi Studies (CREDES) in palliative care: Recommendations based on a methodological systematic review. *Palliat Med* 2017;**31**:684–706.
 38. Williamson C, Kelly P, Tomasone JR, et al. A modified Delphi study to enhance and gain international consensus on the Physical Activity Messaging Framework (PAMF) and Checklist (PAMC). *Int J Behav Nutr Phys Act* 2021;**18**:108. doi:10.1186/s12966-021-01182-z.
 39. Gracia-Sánchez A, López-Pineda A, Lázaro-Martínez JL, et al. Consensus-based recommendations on physical activity and exercise in patients with diabetes at risk of foot ulcerations: A Delphi study. *Braz J Phys Ther* 2023;**27**:100500. doi:10.1016/j.bjpt.2023.100500.
 40. Virgara R, Phillips A, Lewis L, Richardson M, Maher C. Development of Australian physical activity and screen time guidelines for outside school hours care: An international Delphi study. *Int J Behav Nutr Phys Act* 2021;**18**:3. doi:10.1186/s12966-020-01061-z.
 41. Diamond IR, Grant RC, Feldman BM, et al. Defining consensus: A systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014;**67**:401–9.
 42. Barrios M, Guilera G, Nuño L, Gómez-Benito J. Consensus in the delphi method: What makes a decision change? *Technol Forecast Soc Change* 2021;**163**:120484. doi:10.1016/j.techfore.2020.120484.
 43. España-Romero V, Artero EG, Santaliestra-Pasias AM, Gutierrez A, Castillo MJ, Ruiz JR. Hand span influences optimal grip span in boys and girls aged 6 to 12 years. *J Hand Surgery Am* 2008;**33**:378–84.
 44. Ruiz JR, España-Romero V, Ortega FB, Sjöström M, Castillo MJ, Gutierrez A. Hand span influences optimal grip span in male and female teenagers. *J Hand Surgery Am* 2006;**31**:1367–72.
 45. Vicente-Rodríguez G, Rey-López JP, Mesana MI, et al. Reliability and inter-method agreement for body fat assessment among two field and two laboratory methods in adolescents. *Obesity (Silver Spring)* 2012;**20**:221–8.
 46. Wells JCK. Commentary: The paradox of body mass index in obesity assessment: Not a good index of adiposity, but not a bad index of cardiometabolic risk. *Int J Epidemiol* 2014;**43**:672–4.
 47. Ortega FB, Sui X, Lavie CJ, Blair SN. Body mass index, the most widely used but also widely criticized index: Would a criterion standard measure of total body fat be a better predictor of cardiovascular disease mortality? *Mayo Clin Proc* 2016;**91**:443–55.
 48. Bianco A, Jemni M, Thomas E, et al. A systematic review to determine reliability and usefulness of the field-based test batteries for the assessment of physical fitness in adolescents—The ASSO Project. *Int J Occup Med Environ Health* 2015;**28**:445–78.
 49. Batista MB, Romanzini CLP, Castro-Piñero J, Ronque ERV. Validity of field tests to estimate cardiorespiratory fitness in children and adolescents: A systematic review. *Rev Paul Pediatr* 2017;**35**:222–33.
 50. Lang JJ, Belanger K, Poitras V, Janssen I, Tomkinson GR, Tremblay MS. Systematic review of the relationship between 20m shuttle run performance and health indicators among children and youth. *J Sci Med Sport* 2018;**21**:383–97.
 51. Rollo S, Fraser BJ, Seguin N, et al. Health-related criterion-referenced cut-points for cardiorespiratory fitness among youth: A systematic review. *Sports Med* 2022;**52**:101–22.
 52. Domone S, Mann S, Sandercock G, Wade M, Beedie C. A method by which to assess the scalability of field-based fitness tests of cardiorespiratory fitness among schoolchildren. *Sports Med* 2016;**46**:1819–31.
 53. Fraser BJ, Rollo S, Sampson M, et al. Health-related criterion-referenced cut-points for musculoskeletal fitness among youth: A systematic review. *Sports Med* 2021;**51**:2629–46.
 54. Vanhelst J, et al. Cardiorespiratory fitness has declined among French children since 1999, although the decline appears to be getting smaller. *Scand J Med Sci Sports* 2024;**34**:e14641. doi:10.1111/sms.14641.
 55. Poon ET, Tomkinson GR, Lang JJ, Huang WY, Wong SH. Temporal trends in the physical fitness of Hong Kong children aged 6–12 years between 2003–04 and 2015–16. *J Sports Sci* 2023;**41**:1271–8.
 56. Tomkinson GR, Kaster T, Dooley FL, et al. Temporal trends in the standing broad jump performance of 10,940,801 children and adolescents between 1960 and 2017. *Sports Med* 2021;**51**:531–48.
 57. Pinoniemi BK, Tomkinson GR, Walch TJ, Roemmich JN, Fitzgerald JS. Temporal trends in the standing broad jump performance of united states children and adolescents. *Res Q Exerc Sport* 2021;**92**:71–81.
 58. Dooley FL, Kaster T, Fitzgerald JS, et al. A systematic analysis of temporal trends in the handgrip strength of 2,216,320 children and adolescents between 1967 and 2017. *Sports Med* 2020;**50**:1129–44.
 59. Fraser BJ, Blizzard L, Tomkinson GR, et al. The great leap backward: Changes in the jumping performance of Australian children aged 11–12 years between 1985 and 2015. *J Sports Sci* 2019;**37**:748–54.
 60. Tomkinson GR, Lang JJ, Tremblay MS. Temporal trends in the cardiorespiratory fitness of children and adolescents representing 19 high-income and upper middle-income countries between 1981 and 2014. *Br J Sports Med* 2019;**53**:478–86.
 61. Tomkinson GR, Léger LA, Olds TS, Cazorla G. Secular trends in the performance of children and adolescents (1980–2000): An analysis of 55 studies of the 20m shuttle run test in 11 countries. *Sports Med* 2003;**33**:285–300.
 62. Poon ET, Tomkinson GR, Huang WY, Wong SH. Temporal trends in the physical fitness of Hong Kong adolescents between 1998 and 2015. *Int J Sports Med* 2023;**44**:728–35.
 63. Ortega FB, Leskošek B, Blagus R, et al. European fitness landscape for children and adolescents: Updated reference values, fitness maps and country rankings based on nearly 8 million test results from 34 countries gathered by the FitBack network. *Br J Sports Med* 2023;**57**:299–310.
 64. Tomkinson GR, Lang JJ, Tremblay MS, et al. International normative 20 m shuttle run values from 1,142,026 children and youth representing 50 countries. *Br J Sports Med* 2017;**51**:1545–54.
 65. Klemm K, Bös K, Kron H, Eberhardt T, Woll A, Niessner C. Development and introduction of a disciplinary data repository for sport scientists based on the example MO|RE Data: eResearch infrastructure for motor research data. *Bausteine Forschungsdatenmanag* 2024. doi:10.17192/bfdm.2024.1.8615.
 66. Boivin N, Täuber S, Beisiegel U, Keller U, Hering JG. Sexism in academia is bad for science and a waste of public funding. *Nat Rev Mater* 2024;**9**:1–3.
 67. Shannon G, Jansen M, Williams K, et al. Gender equality in science, medicine, and global health: Where are we at and why does it matter? *The Lancet* 2019;**393**:560–9.

- 1701 68. Dimitris MC, Gittings M, King NB. How global is global health
1702 research? A large-scale analysis of trends in authorship. *BMJ Glob*
1703 *Health* 2021;**6**:e003758. doi:10.1136/bmjgh-2020-003758.
- 1704 69. Martínez-Rosales E, Hernández-Martínez A, Sola-Rodríguez S, Esteban-
1705 Cornejo I, Soriano-Maldonado A. Representation of women in sport
1706 sciences research, publications, and editorial leadership positions: Are we
1707 moving forward? *J Sci Med Sport* 2021;**24**:1093–7.
- 1708 70. Cowan SM, Kemp JL, Ardern CL, et al. Sport and exercise medicine/
1709 physiotherapy publishing has a gender/sex equity problem: We need
1710 action now!. *Br J Sports Med* 2023;**57**:401–7.
- 1711 71. Sánchez-Aranda L, Fernández-Ortega J, Martín-Fuentes I, et al.
1712 Reliability and criterion validity of a low-cost handgrip
1713 dynamometer: The Camry. *medRxiv* 2024. doi:10.1101/2024.06.25.
1714 24309304.
- 1715 72. España-Romero V, Ortega FB, Vicente-Rodríguez G, Artero EG, Rey JP,
1716 Ruiz JR. Elbow position affects handgrip strength in adolescents: Validity
1717 and reliability of Jamar, DynEx, and TKK dynamometers. *J Strength*
1718 *Cond Res* 2010;**24**:272–7.
- 1719 73. Wouters M, Evenhuis HM, Hilgenkamp TI. Systematic review of field-
1720 based physical fitness tests for children and adolescents with intellectual
1721 disabilities. *Res Dev Disabil* 2017;**61**:77–94.
- 1722 74. Hutzler Y, Barak S, Aubert S, et al. WOT” do we know and do about phys-
1723 ical activity of children and adolescents with disabilities? A SWOT-oriented
1724 synthesis of para report cards. *Adapt Phys Activ Q* 2023;**40**:431–55.
- 1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757