



The interplay of gut microbiota, obesity, and depression: insights and interventions

Iryna Halabitska¹ · Pavlo Petakh² · Iryna Kamyshna³ · Valentyn Oksenysh⁴ · Denis E. Kainov⁵ · Oleksandr Kamyshnyi⁶

Received: 30 July 2024 / Revised: 8 September 2024 / Accepted: 7 October 2024
© The Author(s) 2024

Abstract

The gut microbiome, body weight, and related comorbidities are intricately linked through a complex interaction of microbial, genetic, environmental, and psychological factors. Alterations in gut microbiota can contribute to the development of weight disorders and depressive symptoms, with the potential for these relationships to be bidirectional. Effective management of these interconnected conditions often involves a combination of lifestyle modifications and psychological support. Medical interventions, including treatments for obesity, antidiabetic drugs, antidepressants, antibiotics, and probiotics, can have beneficial and detrimental effects on gut microbiota and mental health. Further research is needed to better understand their impact on gut microbiome and mental health in the context of obesity.

Keywords Obesity · Depression · Gut microbiota · Mental health · Antidepressants · Probiotics

Introduction

Genetic predispositions, socio-demographic variables, environmental conditions, psychological states, lifestyle choices, infections, and medication use influence the

interplay between microbiome and health (Fig. S1a). Disruptions in the gut microbiota can lead to metabolic and psychological dysregulation, contributing to obesity and depression through mechanisms such as chronic inflammation, altered metabolism, and hormonal imbalances (Fig. S1b). Understanding these interactions is crucial for advancing our health knowledge and developing targeted prevention and treatment strategies for associated diseases [1, 2]. This review examines the interconnections between gut microbiota, obesity, and depression.

✉ Pavlo Petakh
pavlo.petakh@uzhnu.edu.ua

✉ Denis E. Kainov
denis.kainov@ntnu.no

- ¹ Department of Therapy and Family Medicine, I. Horbachevsky Ternopil National Medical University, Voli Square, 1, Ternopil 46001, Ukraine
- ² Department of Biochemistry and Pharmacology, Uzhhorod National University, Uzhhorod 88000, Ukraine
- ³ Department of Medical Rehabilitation, I. Horbachevsky Ternopil National Medical University, Ternopil 46001, Ukraine
- ⁴ Department of Clinical Science, University of Bergen, Bergen 5020, Norway
- ⁵ Department of Clinical and Molecular Medicine (IKOM), Norwegian University of Science and Technology, Trondheim 7028, Norway
- ⁶ Department of Microbiology, Virology, and Immunology, I. Horbachevsky Ternopil National Medical University, Ternopil 46001, Ukraine

Intestinal microbiota

The gut microbiota, composed of over 1,000 species of bacteria, archaea, fungi, protozoa, and viruses significantly impacts human health [3, 4]. It plays a crucial role in immune system maturation, maintaining intestinal integrity, producing vitamins K and B, preventing infections, and regulating metabolism [5].

The bacterial genome content in the gut is 150 times larger than the human genome [4]. The main microbial phyla in the gut are *Actinobacteria*, *Bacteroidetes*, *Firmicutes*, *Fusobacteria*, *Proteobacteria*, and *Verrucomicrobia*, with *Bacteroidetes* and *Firmicutes* making up about 90% of the gut microbiome [6, 7]. A balanced gut microbiota, or eubiosis,

enhances immunity, nutrient absorption, blood glucose and lipid balance, and energy metabolism, while also producing essential vitamins and protecting against pathogens. Imbalanced gut microbiota, or dysbiosis, can result from antibiotics, poor diet, stress, and other factors, leading to health issues [8, 9].

Gut microbiota composition varies across the gastrointestinal tract due to differences in environment and flow rates. Factors like age, diet, antibiotics, and exercise influence the microbiome. Gut bacteria metabolize dietary polysaccharides into short-chain fatty acids (SCFAs), providing about 70% of the energy for colon cells and contributing to glucose homeostasis, lipid metabolism, immune function, and overall health [10].

Obesity

Obesity, affecting over 650 million people globally, significantly contributes to morbidity and mortality [11]. Defined by a body mass index (BMI) over 30 kg/m² (overweight is 25–30 kg/m²), obesity results from an imbalance between energy intake and expenditure, leading to excessive fat accumulation. It is linked to diseases like type 2 diabetes, cardiovascular diseases, certain cancers, and high blood pressure [12].

This complex disease is influenced by diet, antibiotic use, environment, genetics, lifestyle, socioeconomic factors, and gut microbiota composition. High-calorie diets contribute to obesity and alter gut microbiome function. Modern lifestyle factors, hormonal imbalances, and genetic influences drive the obesity crisis. Gut hormones communicate food intake and energy stores to the brain, regulating appetite [13].

Depression

Depression is a common chronic mental disorder characterized by symptoms such as insomnia, melancholy, lack of enjoyment in life, and low energy. Its prevalence is estimated at 11% in low- to middle-income countries and 15% in high-income countries, affecting about 280 million people globally [14, 15]. Annually, depression and anxiety cost an estimated US\$ 1 trillion due to the loss of 12 billion productive workdays [16].

Depression is significantly correlated with various physical health conditions, including suicide, cancers, respiratory diseases, diabetes, and cardiovascular diseases. In the USA alone, depression-related suicides claim about 40,000 lives annually, among older men [17]. According to WHO reports from 2016, around 785,000 suicides occur worldwide each year, with up to 60% attributed to depression [18, 19].

Interplay between gut microbiota, obesity and depression

There is a link between gut microbiome, obesity, and depression (Fig. 1). For example, an increased *Firmicutes/Bacteroidetes* ratio is associated with obesity. However, this ratio alone cannot fully explain obesity due to the gut microbiome's complexity [20, 21]. For instance, mutations in the leptin gene in mice have been linked to altered bacterial proportions [8, 20]. The gut microbiome affects energy intake and metabolism, influencing weight and fat storage [22, 23].

Obese individuals often have microbiomes that increase energy extraction and fat accumulation. Gut microbiome composition, such as higher *Prevotella* levels, is associated with greater weight loss compared to *Bacteroides* dominance [24, 25]. Inflammation caused by microbiota can affect leptin expression and thermogenesis, contributing to obesity. Chronic low-grade inflammation, driven by lipopolysaccharides (LPS), impairs energy expenditure (Fig. 2) [9].

The gut microbiome also impacts vitamin synthesis, xenobiotic metabolism, and obesity regulation. Studies using germ-free mice have highlighted the microbiome's role in energy metabolism and mental health. The gut-brain axis connects microbiome composition with brain function and mental disorders, such as depression [26–28]. Previous studies indicate alterations in the gut microbiota of patients with mental disorders, with the most studied phyla being *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Proteobacteria*, and *Verrucomicrobia*, and key genera including *Prevotella*, *Coprococcus*, *Parabacteroides*, *Phascolarctobacterium*, *Escherichia-Shigella*, *Bacteroides*, *Alistipes*, and *Veillonella* [29].

Specific bacteria, like *Alistipes* and *Veillonella*, are associated with depression, while others, like *Prevotella*, are less prevalent [27, 30]. Dysbiosis, including imbalances in gut microbiota, is linked to depression and systemic inflammation. Microbial lactate accumulation may also impair brain function and contribute to mental disorders [31, 32].

The link between gut microbiota and mental health, particularly depression, is increasingly recognized through the concept of the gut-brain axis. This bidirectional communication pathway allows gut bacteria to influence brain function via neurotransmitter production, immune responses, and signaling molecules. Disruptions in gut microbiota, or dysbiosis, can lead to imbalances in neurotransmitters like serotonin, increase systemic inflammation, and impair gut barrier function, all of which may contribute to depressive symptoms [28]. Dysbiosis can lead to an increase in pro-inflammatory cytokines and a disruption in the synthesis of key neurotransmitters like serotonin, dopamine, and




Fig. 1 Gut-brain axis signaling pathways and their influence on mood and emotion regulation. The diagram depicts the interactions between the gut lumen, gut microbiota, and the central nervous system through the vagus nerve, immune cells, and neurotransmitter signaling. Various gut-derived signals, including active neurotransmitters (dopamine, serotonin, glutamate, and gamma-aminobutyric acid (GABA)), as well as tryptophan metabolites and short-chain fatty acids (SCFAs), influ-

ence mood and emotional regulation by transmitting signals to the brain. This communication occurs through the vagus nerve, immune modulation, and the hypothalamic-pituitary-adrenal (HPA) axis, which involves the secretion of corticotropin-releasing hormone (CRH), adrenocorticotropic hormone (ACTH), and cortisol. Dysregulation in these pathways can impact mood and emotional states, contributing to stress, depression, and anxiety

gamma-aminobutyric acid (GABA), contributing to the development of metabolic and mood disorders [33]. Obesity is often associated with mental health issues, particularly depression. High body weight is linked to poorer mental health due to systemic inflammation, hypothalamic-pituitary-adrenal axis dysregulation, and psychological stress from weight stigma and discrimination, exacerbating mental health problems and hindering weight control efforts [34].

Furthermore, imbalances in gut microbiota can impair the production of crucial metabolites like SCFAs, disrupting gut-brain signaling and leading to dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis. This dysregulation can result in elevated cortisol levels and altered serotonin production, which are associated with stress, anxiety, and mood disorders [35–37].

Obesity is influenced by genetics, metabolism, and environmental factors, with growing evidence highlighting the role of gut microbiota in these interactions. Dysbiosis in the gut microbiome has been linked to obesity through pathways that affect metabolism, insulin resistance, and satiety, offering potential therapeutic targets for prevention and treatment [38, 39].

ence mood and emotional regulation by transmitting signals to the brain. This communication occurs through the vagus nerve, immune modulation, and the hypothalamic-pituitary-adrenal (HPA) axis, which involves the secretion of corticotropin-releasing hormone (CRH), adrenocorticotropic hormone (ACTH), and cortisol. Dysregulation in these pathways can impact mood and emotional states, contributing to stress, depression, and anxiety


Interventions of gut microbiota, obesity and depression

Lifestyle modifications, including diet, stress management, exercise, and avoiding harmful substances, are commonly recommended for managing altered gut microbiota, obesity, and depression (Table S1). While these strategies can improve quality of life, long-term results are often modest, with only 61% of individuals completing such programs successfully. In severe cases, medical interventions, surgical options, and psychotherapy may be needed to address these complex issues (Table S2, S3). Effective management can enhance health outcomes and reduce the prevalence of related conditions like type 2 diabetes, cardiovascular disease, and certain cancers, thereby improving overall public health [40].

Antibiotics and probiotics

Antibiotics like amoxicillin and ciprofloxacin can disrupt gut microbiota by killing both harmful and beneficial bacteria. This imbalance may lead to altered neurotransmitter levels, increased inflammation, and gastrointestinal issues, potentially contributing to mood disturbances and depressive symptoms.

Fig. 2 Interconnection between obesity, diet, inactivity, gut microbiota, and the hypothalamic-pituitary-adrenal (HPA) axis contributing to inflammation and depression. The diagram illustrates the complex relationship between inactivity and diet changes, which interacts with hormonal regulators like ghrelin, insulin, and leptin. These factors influence gut microbiota, leading to reduced short-chain fatty acids (SCFAs) and increased gut barrier disruption. Such disturbances in gut health elevate cortisol and adrenocorticotropic hormone (ACTH) levels through the activation of the HPA axis, fueling inflammation, pain, fatigue, and depressive symptoms, including anhedonia. This bidirectional feedback loop highlights the role of chronic inflammation in exacerbating psychological and metabolic health issues.



Probiotics, such as *Lactobacillus rhamnosus* and *Bifidobacterium longum*, can help restore a balanced gut microbiota. They may improve mental health by enhancing gut barrier function, modulating neurotransmitter levels, and reducing stress and anxiety. These effects vary by strain and individual response, but these probiotics are commonly studied for their potential mental health benefits [41, 42].

GLP-1 receptor agonists

In 2016, the American Association of Clinical Endocrinology reviewed five anti-obesity medications (orlistat, lorcaserin, naltrexone–bupropion, liraglutide, and phentermine–topiramate) for long-term use. However, the FDA withdrew lorcaserin in 2020 due to cancer risks. In 2021, studies on semaglutide, a new weekly Glucagon-Like Peptide-1 (GLP-1) receptor agonist, showed promising results that could impact clinical practice. Other diabetes medications, such as Sodium-Glucose Cotransporter 2 (SGLT2) inhibitors, metformin, and pramlintide, also show potential for obesity management [43, 44].


Anti-obesity drugs like liraglutide and semaglutide affect gut microbiota, enhancing bacterial diversity and influencing metabolic disorders. Semaglutide significantly alters gut microbiota, impacting mental health and metabolic pathways. Liraglutide has shown therapeutic effects on fatty liver and modifies gut bacteria linked to metabolism and inflammation (Fig. 3) [45–47].

Both liraglutide and semaglutide impact the gut microbiota in conditions like Polycystic Ovary Syndrome (PCOS). They influence the abundance of various bacterial species and promote beneficial bacteria, altering the gut microbiome's diversity and composition. These changes may optimize obesity treatment outcomes [46].

Studies on GLP-1 receptor agonists reveal mixed effects on mental health. Semaglutide and liraglutide have been associated with depressive symptoms, anxiety, and, in some cases, suicidal thoughts. However, there's no clear causal link between GLP-1R agonism and severe depression or suicide. Monitoring for mood changes is advised for patients using these medications [48–50].

Recent social media discussions also reflect concerns about GLP-1 receptor agonists affecting mood and mental

Fig. 3 Effect of semaglutide on mental health and gut microbiome. This diagram illustrates how Semaglutide influences mental health and changes the gut microbiota, showing its combined effects on managing obesity and related conditions



health, indicating the need for ongoing vigilance and assessment [51].

Antidiabetic drugs

Metformin affects gut microbiota by altering its composition, enhancing SCFA-producing bacteria like *Butyricimonas* and beneficial probiotics such as *Lactobacillus*, while reducing harmful pathogens like *Prevotella* [52, 53]. It improves glucose control in obese mice and enhances SCFA-producing bacteria, and in T2DM and COVID-19 patients, it boosts beneficial bacteria like *Bacteroides* while lowering harmful ones [54]. Metformin might also impact mental health and depressive symptoms, potentially offering benefits for treatment-resistant depression. However, the exact mechanisms of its effects on mental health need further research [55, 56].

Anti-obesity pharmacological agents

Naltrexone-Bupropion has been shown to influence gut microbiome diversity [57]. Orlistat treatment results in a reduction of *Firmicutes* and an increase in *Bacteroidetes*, with associated improvements in the diversity indices and elevated levels of *Lactobacillus*, particularly *Lactobacillus gasseri* [58]. Topiramate increases *Lactobacillus johnsonii* but significantly reduces the overall abundance of *Lactobacillus* [59]. SGLT2 Inhibitors alter the prevalence of specific bacteria, including the LPS-producing *Oscillibacter* and SCFA-producing *Bacteroides* and *Odoribacter*, while

boosting *Ruminococci*, which are beneficial for SCFA production [60].

Regarding mental health outcomes, behavioral weight loss (BWL) and Naltrexone-Bupropion have demonstrated efficacy in treating binge-eating disorder, with BWL showing superior results compared to no intervention [61]. Orlistat has been linked to increased physical activity and reduced depression in Latinx patients with obesity, although these effects were not directly associated with weight loss [62]. Phentermine-Topiramate has been shown to alleviate depressive symptoms in obese patients and improve quality of life, though it poses risks such as hypomania in bipolar patients and potential psychosis when combined with venlafaxine [63].

SGLT2 inhibitors have improved quality of life in diabetes patients and show promise for adjunctive treatment of major depressive disorder (MDD) [64]. These findings highlight the complex relationship between weight loss medications, gut microbiota changes, and psychiatric outcomes.

Antidepressants

Antidepressants are pharmacological agents used primarily to treat major depressive disorder (MDD) and other mood disorders by modulating neurotransmitter systems in the brain. The most widely prescribed antidepressants include Selective Serotonin Reuptake Inhibitors (SSRIs), such as fluoxetine and sertraline, which enhance serotonergic neurotransmission by inhibiting the reuptake of serotonin. Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs), including venlafaxine and duloxetine, act on both serotonin

and norepinephrine, aiming to alleviate depressive symptoms through dual neurotransmitter modulation. Tricyclic Antidepressants (TCAs) and Monoamine Oxidase Inhibitors (MAOIs), though less commonly used as first-line treatments due to their broader side effect profiles, remain effective for treatment-resistant cases and specific mood disorders. TCAs, such as amitriptyline, affect multiple neurotransmitter systems, while MAOIs, such as phenelzine, inhibit monoamine oxidase, leading to increased levels of serotonin, norepinephrine, and dopamine [65, 66].

Atypical antidepressants, including bupropion and mirtazapine, exhibit diverse mechanisms and are employed based on individual patient profiles and specific clinical needs. Bupropion, for instance, primarily affects norepinephrine and dopamine systems, while mirtazapine enhances neurotransmitter release through its antagonistic effects on certain receptors. The therapeutic efficacy of antidepressants is often accompanied by a range of side effects, which vary by drug class and individual patient response. Monitoring and managing these effects are crucial, as abrupt discontinuation of antidepressants can precipitate withdrawal symptoms. Overall, the choice of antidepressant and treatment regimen should be tailored to the patient's specific condition and response to therapy, with ongoing assessment to optimize outcomes and minimize adverse effects [65, 66].

The studies demonstrate that fluoxetine significantly affected the growth of *E. coli*, *E. faecalis*, and *S. aureus*, with minimal impact on *C. albicans*. In contrast, escitalopram notably influenced the growth of *E. coli*, *E. faecalis*, *B. bifidum*, *L. rhamnosus*, and *C. albicans*, while mirtazapine exhibited the greatest activity against *L. rhamnosus* and *C. albicans* [67]. Antidepressants can modify the abundance and composition of intestinal microbiota, which in turn affects treatment outcomes for depression by influencing drug metabolism, absorption, and blood-brain barrier permeability [68].

Conclusions

Changes in gut microbiota and the presence of depressive disorders are commonly noted in obesity, with these conditions potentially aggravating each other. The influence of different treatments—such as antibiotics, probiotics, anti-obesity medications, antidiabetic agents, and antidepressants—on gut microbiota and mental health is often intricate and multifaceted. These interventions can produce conflicting effects and may differ considerably among individuals. Consequently, thorough research is required to clarify how these treatments affect gut microbiota and mental health.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00018-024-05476-w>.

Acknowledgements We thank Payam Behzadi for critical reading of the manuscript.

Author contributions Conceptualization, O.K.; software and data curation, I.H., P.P., O.K.; writing—original draft preparation, I.H., V.O., I.K.; writing—review and editing, all authors.; visualization, P.P., O.K., I.K., V.O.; supervision, O.K., D.K., V.O.; project administration, D.K., O.K. All authors have read and agreed to the published version of the manuscript.

Funding This research was funded by Helse Midt-Norge, Felles forskningsutvalg (FFU) grant Nr. 34155 (to D.E.K.). Open access funding provided by NTNU Norwegian University of Science and Technology (incl St. Olavs Hospital - Trondheim University Hospital)

Data availability All data generated or analyzed during this study are included in this published article and its supplementary information files.

Declarations

Ethical approval Not applicable.

Competing interests The authors have no relevant financial or non-financial interests to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Carmody RN, Bisanz JE (2023) Roles of the gut microbiome in weight management. *Nat Rev Microbiol* 21(8):535–550
2. Gou W, Miao Z, Deng K, Zheng JS (2023) Nutri-Microbiome epidemiology, an emerging field to disentangle the interplay between nutrition and microbiome for human health. *Protein Cell* 14(11):787–806
3. Behzadi P, Dodero VI, Golubnitschaja O (2024) Systemic inflammation as the Health-Related Communication Tool between the Human Host and gut microbiota in the Framework of Predictive, Preventive, and Personalized Medicine. *All around Suboptimal Health: Advanced approaches by Predictive, Preventive and Personalised Medicine for healthy populations*. Springer, pp 203–241
4. Lathakumari RH, Vajravelu LK, Satheesan A, Ravi S, Thulukanam J (2024) Antibiotics and the gut microbiome: understanding the impact on human health. *Med Microecology*. :100106

5. Mukherjee S, Patra R, Behzadi P, Masotti A, Paolini A, Sarshar M (2023) Toll-like receptor-guided therapeutic intervention of human cancers: molecular and immunological perspectives. *Front Immunol* 14:1244345
6. Rinninella E, Raoul P, Cintoni M, Franceschi F, Miggiano GAD, Gasbarrini A, Mele MC (2019) What is the healthy gut microbiota composition? A changing ecosystem across age, environment, diet, and diseases. *Microorganisms* 7(1):14
7. Behzadi P, Najafi A, Behzadi E, Ranjbar R (2016) Microarray long oligo probe designing for *Escherichia coli*: an in-silico DNA marker extraction. *Cent Eur J Urol* 69(1):105
8. Muscogiuri G, Cantone E, Cassarano S, Tuccinardi D, Barrea L, Savastano S, Colao A (2019) Obesity programs of nutrition E, Research, group A. Gut microbiota: a new path to treat obesity. *Int J Obes Supplements* 9(1):10–19
9. Montenegro J, Armet AM, Willing BP, Deehan EC, Fassini PG, Mota JF, Walter J, Prado CM (2023) Exploring the influence of gut microbiome on energy metabolism in humans. *Adv Nutr* 14(4):840–857
10. Li Z, Yi C-X, Katiraei S, Kooijman S, Zhou E, Chung CK, Gao Y, van den Heuvel JK, Meijer OC, Berbée JF (2018) Butyrate reduces appetite and activates brown adipose tissue via the gut-brain neural circuit. *Gut* 67(7):1269–1279
11. de Wit DF, Hanssen NMJ, Wortelboer K, Herrema H, Rampantelli E, Nieuwdorp M (2023) Evidence for the contribution of the gut microbiome to obesity and its reversal. *Sci Transl Med* 15(723):eadg2773
12. Van Hul M, Cani PD (2023) The gut microbiota in obesity and weight management: microbes as friends or foe? *Nat Reviews Endocrinol* 19(5):258–271
13. Kemp JVA, Kumar V, Saleem A, Hashman G, Hussain M, Taylor VH (2023) Examining associations between women's Mental Health and obesity. *Psychiatr Clin North Am* 46(3):539–549
14. Bromet E, Andrade LH, Hwang I, Sampson NA, Alonso J, De Girolamo G, De Graaf R, Demyttenaere K, Hu C, Iwata N (2011) Cross-national epidemiology of DSM-IV major depressive episode. *BMC Med* 9:1–16
15. Radjabzadeh D, Bosch JA, Uitterlinden AG, Zwinderman AH, Ikram MA, van Meurs JB, Luik AI, Nieuwdorp M, Lok A, van Duijn CM (2022) Gut microbiome-wide association study of depressive symptoms. *Nat Commun* 13(1):7128
16. (WHO) WHO, Depression, WHO (2024); [https://www.who.int/health-topics/depression#tab=tab_2]
17. Ferrari S, Mulè S, Parini F, Galla R, Ruga S, Rosso G, Brovero A, Molinari C, Uberti F (2024) The influence of the gut-brain axis on anxiety and depression: a review of the literature on the use of probiotics. *J Traditional Complement Med*
18. Luqman A, He M, Hassan A, Ullah M, Zhang L, Rashid Khan M, Din AU, Ullah K, Wang W, Wang G (2024) Mood and microbes: a comprehensive review of intestinal microbiota's impact on depression. *Front Psychiatry* 15
19. Turecki G, Brent DA, Gunnell D, O'Connor RC, Oquendo MA, Pirkis J, Stanley BH (2019) Suicide and suicide risk. *Nat Reviews Disease Primers* 5(1):74
20. Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD, Gordon JI (2005) Obesity alters gut microbial ecology. *Proceedings of the national academy of sciences* 102(31):11070–5
21. Ryan PM, Patterson E, Carafa I, Mandal R, Wishart DS, Dinan TG, Cryan JF, Tuohy KM, Stanton C, Ross RP (2020) Metformin and Dipeptidyl Peptidase-4 inhibitor differentially modulate the intestinal microbiota and plasma metabolome of metabolically dysfunctional mice. *Can J Diabetes* 44(2):146–55e2
22. Rigoulet M, Bouchez CL, Paumard P, Ransac S, Cuvellier S, Duvezin-Caubet S, Mazat JP, Devin A (2020) Cell energy metabolism: an update. *Biochim et Biophys Acta (BBA)-Bioenergetics* 1861(11):148276
23. Galgani J, Ravussin E (2008) Energy metabolism, fuel selection and body weight regulation. *Int J Obes* 32(7):S109–S119
24. Hjorth MF, Roager HM, Larsen TM, Poulsen S, Licht TR, Bahl MI, Zohar Y, Astrup A (2018) Pre-treatment microbial Prevotella-to-Bacteroides ratio, determines body fat loss success during a 6-month randomized controlled diet intervention. *Int J Obes* 42(3):580–583
25. Hjorth MF, Blædel T, Bendtsen LQ, Lorenzen JK, Holm JB, Kiil-erich P, Roager HM, Kristiansen K, Larsen LH, Astrup A (2019) Prevotella-to-Bacteroides ratio predicts body weight and fat loss success on 24-week diets varying in macronutrient composition and dietary fiber: results from a post-hoc analysis. *Int J Obes* 43(1):149–157
26. Behzadi P, Kim C-H, Pawlak EA, Algammal A (2023) The innate and adaptive immune system in human urinary system. *Front Immunol* 14:1294869
27. Wang M, Song Z, Lai S, Tang F, Dou L, Yang F (2024) Depression-associated gut microbes, metabolites and clinical trials. *Front Microbiol* 15:1292004
28. McGuinness A, Davis JA, Dawson S, Loughman A, Collier F, O'hely M, Simpson C, Green J, Marx W, Hair C (2022) A systematic review of gut microbiota composition in observational studies of major depressive disorder, bipolar disorder and schizophrenia. *Mol Psychiatry* 27(4):1920–1935
29. Zang Y, Lai X, Li C, Ding D, Wang Y, Zhu Y (2023) The role of gut microbiota in various Neurological and Psychiatric Disorders-An evidence mapping based on quantified evidence. *Mediat Inflamm* 2023:5127157
30. George F, Daniel C, Thomas M, Singer E, Guilbaud A, Tessier FJ, Revol-Junelles A-M, Borges F, Foligné B (2018) Occurrence and dynamism of lactic acid bacteria in distinct ecological niches: a multifaceted functional health perspective. *Front Microbiol* 9:2899
31. Giménez-Palomo A, Dodd S, Anmella G, Carvalho AF, Scaini G, Quevedo J, Pacchiarotti I, Vieta E, Berk M (2021) The role of mitochondria in mood disorders: from physiology to pathophysiology and to treatment. *Front Psychiatry* 12:546801
32. Larrea A, Sánchez-Sánchez L, Diez-Martin E, Elexpe A, Torrecilla M, Astigarraga E, Barreda-Gómez G (2024) Mitochondrial metabolism in major depressive disorder: from early diagnosis to Emerging Treatment options. *J Clin Med* 13(6):1727
33. Steffen J, Focken N, Çalışkan G (2024) Recognizing depression as an inflammatory disease: the search for endotypes. *Am J Physiol Cell Physiol* 327(1):C205–c12
34. Gardiner JV, Jayasena CN, Bloom SR (2008) Gut hormones: a weight off your mind. *J Neuroendocrinol* 20(6):834–841
35. Steptoe A, Frank P (2023) Obesity and psychological distress. *Philosophical Trans Royal Soc Lond Ser B Biol Sci* 378(1888):20220225
36. Song EJ, Shin NR, Jeon S, Nam YD, Kim H (2022) Lorcaserin and phentermine exert anti-obesity effects with modulation of the gut microbiota. *Front Microbiol* 13:1109651
37. Pati B, Sendh S, Sahu B, Pani S, Jena N, Bal NC (2023) Recent advancements in pharmacological strategies to modulate energy balance for combating obesity. *RSC Med Chem* 14(8):1429–1445
38. Lee CJ, Sears CL, Maruthur N (2020) Gut microbiome and its role in obesity and insulin resistance. *Ann N Y Acad Sci* 1461(1):37–52
39. Wu L, Ye S, Deng X, Fu Z, Li J, Yang C (2024) Conjugated linoleic acid ameliorates high Fat-Induced insulin resistance via regulating gut microbiota-host metabolic and immunomodulatory interactions. *Nutrients*; 16(8)
40. Shi Q, Wang Y, Hao Q, Vandvik PO, Guyatt G, Li J, Chen Z, Xu S, Shen Y, Ge L, Sun F, Li L, Yu J, Nong K, Zou X, Zhu S, Wang C, Zhang S, Qiao Z, Jian Z, Li Y, Zhang X, Chen K, Qu F, Wu Y, He Y, Tian H, Li S (2024) Pharmacotherapy for adults

- with overweight and obesity: a systematic review and network meta-analysis of randomised controlled trials. *Lancet* (London England) 403(10434):e21–e31
41. Boggio Marzet C, Burgos F, Del Compare M, Gerold I, Tabacco O, Vinderola G (2022) Approach to probiotics in pediatrics: the role of *Lactobacillus rhamnosus* GG. *Arch Argentinos De Pediatría* 120(1):e1–e7
 42. Mills S, Yang B, Smith GJ, Stanton C, Ross RP (2023) Efficacy of *Bifidobacterium longum* alone or in multi-strain probiotic formulations during early life and beyond. *Gut Microbes* 15(1):2186098
 43. Davies M, Færch L, Jeppesen OK, Pakseresht A, Pedersen SD, Perreault L, Rosenstock J, Shimomura I, Viljoen A, Wadden TA, Lingvay I (2021) Semaglutide 2.4 mg once a week in adults with overweight or obesity, and type 2 diabetes (STEP 2): a randomised, double-blind, double-dummy, placebo-controlled, phase 3 trial. *Lancet* (London England) 397(10278):971–984
 44. Rubino D, Abrahamsson N, Davies M, Hesse D, Greenway FL, Jensen C, Lingvay I, Mosenzon O, Rosenstock J, Rubio MA, Rudofsky G, Tadayon S, Wadden TA, Dicker D (2021) Effect of continued Weekly Subcutaneous Semaglutide vs Placebo on Weight loss maintenance in adults with overweight or obesity: the STEP 4 Randomized Clinical Trial. *JAMA* 325(14):1414–1425
 45. Mao T, Zhang C, Yang S, Bi Y, Li M, Yu J (2024) Semaglutide alters gut microbiota and improves NAFLD in db/db mice. *Biochem Biophys Res Commun* 710:149882
 46. Xiong C, Wu J, Ma Y, Li N, Wang X, Li Y, Ding X (2024) Effects of Glucagon-Like Peptide-1 receptor agonists on Gut Microbiota in Dehydroepiandrosterone-Induced Polycystic Ovary Syndrome mice: compared evaluation of Liraglutide and Semaglutide intervention. *Diabetes Metabolic Syndrome Obesity: Targets Therapy* 17:865–880
 47. de Paiva IHR, da Silva RS, Mendonça IP, de Souza JRB, Peixoto CA (2024) Semaglutide attenuates anxious and depressive-like behaviors and reverses the cognitive impairment in a type 2 diabetes Mellitus Mouse Model Via the Microbiota-Gut-Brain Axis. *J Neuroimmune Pharmacology: Official J Soc NeuroImmune Pharmacol* 19(1):36
 48. Ying X, Rongjiong Z, Kahaer M, Chunhui J, Wulashan M (2023) Therapeutic efficacy of liraglutide versus metformin in modulating the gut microbiota for treating type 2 diabetes mellitus complicated with nonalcoholic fatty liver disease. *Front Microbiol* 14:1088187
 49. Tobaigy M, Elkout H (2024) Psychiatric adverse events associated with semaglutide, liraglutide and tirzepatide: a pharmacovigilance analysis of individual case safety reports submitted to the EudraVigilance database. *Int J Clin Pharm* 46(2):488–495
 50. Tagliapietra GA, Cantrell MA, Lund BC (2024) Glucagon-like peptide receptor agonists and risk for depression. *Prim Care Diabetes*
 51. Arillotta D, Floresta G, Guirguis A, Corkery JM, Catalani V, Martinotti G, Sensi SL, Schifano F (2023) GLP-1 receptor agonists and related Mental Health issues; insights from a range of Social Media platforms using a mixed-methods Approach. *Brain Sci.*; 13(11)
 52. Mueller NT, Differding MK, Zhang M, Maruthur NM, Juraschek SP, Miller ER 3rd, Appel LJ, Yeh HC (2021) Metformin affects gut microbiome composition and function and circulating short-chain fatty acids: a Randomized Trial. *Diabetes Care* 44(7):1462–1471
 53. Rosell-Díaz M, Petit-Gay A, Molas-Prat C, Gallardo-Nuell L, Ramió-Torrentà L, Garre-Olmo J, Pérez-Brocal V, Moya A, Jové M, Pamplona R, Puig J, Ramos R, Bäckhed F, Mayneris-Perxachs J, Fernández-Real JM (2024) Metformin-induced changes in the gut microbiome and plasma metabolome are associated with cognition in men. *Metab Clin Exp* 157:155941
 54. Petakh P, Kobyljak N, Kamyshnyi A (2023) Gut microbiota in patients with COVID-19 and type 2 diabetes: a culture-based method. *Front Cell Infect Microbiol* 13:1142578
 55. Kessing LV, Rytgaard HC, Ekstrøm CT, Knop FK, Berk M, Gerds TA (2020) Antidiabetes agents and Incident Depression: a Nationwide Population-based study. *Diabetes Care* 43(12):3050–3060
 56. Yu H, Yang R, Wu J, Wang S, Qin X, Wu T, Hu Y, Wu Y (2022) Association of metformin and depression in patients with type 2 diabetes. *J Affect Disord* 318:380–385
 57. Raineri S, Sherriff JA, Thompson KSJ, Jones H, Pfluger PT, Illott NE, Mellor J (2022) Pharmacologically induced weight loss is associated with distinct gut microbiome changes in obese rats. *BMC Microbiol* 22(1):91
 58. Uehira Y, Ueno H, Miyamoto J, Kimura I, Ishizawa Y, Iijima H, Muroga S, Fujita T, Sakai S, Samukawa Y, Tanaka Y, Murayama S, Sakoda H, Nakazato M (2023) Impact of the lipase inhibitor orlistat on the human gut microbiota. *Obes Res Clin Pract* 17(5):411–420
 59. Thai K, Taylor MW, Fernandes T, Akinade EA, Campbell SL (2023) Topiramate alters the gut microbiome to aid in its anti-seizure effect. *Front Microbiol* 14:1242856
 60. Deng L, Yang Y, Xu G (2022) Empagliflozin ameliorates type 2 diabetes mellitus-related diabetic nephropathy via altering the gut microbiota. *Biochim et Biophys acta Mol cell Biology Lipids* 1867(12):159234
 61. Grilo CM, Lydecker JA, Fineberg SK, Moreno JO, Ivezaj V, Gueorguieva R (2022) Naltrexone-bupropion and behavior therapy, alone and combined, for binge-eating disorder: Randomized double-blind placebo-controlled trial. *Am J Psychiatry* 179(12):927–937
 62. Grilo CM, Kerrigan SG, Lydecker JA, White MA (2021) Physical activity changes during behavioral weight loss treatment by Latinx patients with obesity with and without binge eating disorder. *Obes (Silver Spring Md)* 29(12):2026–2034
 63. Kolotkin RL, Gadde KM, Peterson CA, Crosby RD (2016) Health-related quality of life in two randomized controlled trials of phentermine/topiramate for obesity: what mediates improvement? *Qual life Research: Int J Qual life Aspects Treat care Rehabilitation* 25(5):1237–1244
 64. Şahin S, Haliloğlu Ö, Polat Korkmaz Ö, Durcan E, Rekalı Şahin H, Yumuk VD, Damcı T, İlkova HM, Oşar Siva Z (2020) Does treatment with sodium-glucose co-transporter-2 inhibitors have an effect on sleep quality, quality of life, and anxiety levels in people with type 2 diabetes mellitus? *Turk J Med Sci* 51(2):735–742
 65. He L, Fu Y, Tian Y, Wang X, Zhou X, Ding RB, Qi X, Bao J (2023) Antidepressants as Autophagy modulators for Cancer Therapy. *Molecules*; 28(22)
 66. Chen T, Cheng L, Ma J, Yuan J, Pi C, Xiong L, Chen J, Liu H, Tang J, Zhong Y, Zhang X, Liu Z, Zuo Y, Shen H, Wei Y, Zhao L (2023) Molecular mechanisms of rapid-acting antidepressants: new perspectives for developing antidepressants. *Pharmacol Res* 194:106837
 67. Rukavishnikov G, Leonova L, Kasyanov E, Leonov V, Neznanov N, Mazo G (2023) Antimicrobial activity of antidepressants on normal gut microbiota: results of the in vitro study. *Front Behav Neurosci* 17:1132127
 68. Xu F, Xie Q, Kuang W, Dong Z (2023) Interactions between antidepressants and Intestinal Microbiota. *Neurotherapeutics: J Am Soc Experimental Neurother* 20(2):359–371

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com